

Confinement, deconfinement and chiral symmetry breaking in QCD

Yuri Simonov*

Institute of Theoretical and Experimental Physics

E-mail: simonov@itep.ru

Confinement is ensured by quadratic colorelectric field correlators of special type, denoted by $D^E(x)$. The latter together with four other quadratic correlators, D^H, D_1^E, D_1^H and D^{EH} support most of perturbative and nonperturbative dynamics, while contribution of higher order terms is shown to be small. These correlators, on the other hand, can be calculated via gluelump Green's function, where dynamics is defined by the correlators themselves. In this way one obtains a self-consistent scheme, where string tension can be expressed in terms of Λ_{QCD} . This confinement mechanism ensures also chiral symmetry breaking and disappears at $T \geq T_c$ where $D^E(x)$ vanishes.

*International Workshop on QCD Green's Functions, Confinement and Phenomenology
September 7-11, 2009
ECT Trento, Italy*

*Speaker.

1. Introduction

Confinement is the most important property of the QCD vacuum (see [1, 2] for reviews) and as shown on the lattice and analytically it can be quite naturally explained as arising from properties of quantum stochastic ensemble of nonperturbative (np) fields filling QCD vacuum. The development of this approach in a systematic way started in 1987 [3] (see, e.g. [4] as example of earlier investigations concerning stochasticity of QCD vacuum). The nontrivial structure of np vacuum can be described by a set of nonlocal gauge-invariant field strength correlators (FC). We discuss this scenario in the present talk (see [5] for an extended version of some results of this talk).

QCD sum rules [6] were suggested as an independent approach to np QCD dynamics, not addressing directly confinement mechanism. The key role is played by gluon condensate G_2 , which is defined as np average of the following type:

$$G_2 = \frac{\alpha_s}{\pi} \langle F_{\mu\nu}^a(0) F_{\mu\nu}^a(0) \rangle \quad (1.1)$$

However the relation of G_2 to confinement is rather tricky, since G_2 is not an order parameter and, in particular, neither perturbative nor np contributions to this quantity vanish in either phase. On the other hand one can show (see details in [7]) that the scale of deconfinement temperature is set by the condensate, or, more precisely, by its electric part: $T_c \sim G_2^{1/4}$. One clearly sees some tension between this "small" scale and a "large" mass scale in QCD given by, e.g. the mass of the lightest 0^{++} glueball, which is about 1.5 GeV.

The question about the origin of this mismatch turned out to be rather deep. It was suggested in [3], that there is another important dimensionfull parameter characterizing np dynamics of vacuum fields: correlation length λ (also denoted as T_g in some papers). The crucial feature of stochastic picture found in [9] is that the lowest, quadratic, nonlocal FC, $\langle F_{\mu\nu}(x) F_{\lambda\sigma}(y) \rangle$ describes all np dynamics with very good accuracy. It was also shown that a simple exponential form of quadratic correlators found on the lattice [9] allows to calculate the correlation length λ entering these exponents, and since λ is small, potential relativistic quantum-mechanical picture is applicable and all QCD spectrum is defined mostly by string tension σ (which is an integral characteristic of the non-local correlator, see below) and not by its exact profile (see [3, 10] for reviews of meson, glueball hybrid spectra).

However to establish the confinement mechanism unambiguously, one should be able to calculate vacuum field distributions, i.e. field correlators, self-consistently. In the long run it means that one is to demonstrate that it is essential property of QCD vacuum fields ensemble to be characterized by correlators, which support confinement for temperatures T below some critical value T_c , and deconfinement at $T > T_c$.

The main aim of this talk is to present this set of equations as a self-consistent mechanism of confinement and to clarify qualitative details of the FC - gluelump connection.

2. Wegner-Wilson loop, Field Correlators and Green's Functions

The Wilson loop is the basic element for discussion of confinement (area law), but also it enters as a main dynamical kernel in the Fock-Feynman-Schwinger Representation (FFSR) of Green's functions of any white QCD object [11].

It is convenient to use nonabelian Stokes theorem (see [2] for references and discussion) for the Wilson loop and to rewrite it as an integral over the surface S inside the contour $C = C(x, y)$:

$$\begin{aligned} \langle W(C) \rangle &= \left\langle \text{Tr} P \exp \left(ig \int_C A_\mu(z) dz_\mu \right) \right\rangle = \left\langle \text{Tr} \mathcal{P} \exp \left(ig \int_S F(u) ds(u) \right) \right\rangle = \\ &= \text{Tr} \mathcal{P}_x \exp \sum_{n=2}^{\infty} (ig)^n \int_S \langle \langle F(u_1) \dots F(u_n) \rangle \rangle ds_1 \dots ds_n = \exp \sum_{n=2}^{\infty} \Delta^{(n)}[S] \end{aligned} \quad (2.1)$$

The central role in the discussed method is played by quadratic (Gaussian) FC of the form

$$D_{\mu\nu, \lambda\sigma} \equiv g^2 \langle \text{Tr} F_{\mu\nu}(x) \Phi(x, y) F_{\lambda\sigma}(y) \Phi(y, x) \rangle, \quad (2.2)$$

where $F_{\mu\nu}$ is the field strength and $\Phi(x, y)$ is the parallel transporter. Correlation lengths λ_i for different channels are defined in terms of asymptotics of (2.2) at large distances: $\exp(-|x-y|/\lambda_i)$. The physical role of λ_i is very important since it distinguishes two regimes: one expects validity of potential-type approach describing the structure of hadrons of spatial size R and at temporal scale T_q for $\lambda_i \ll R, T_q$, while in the opposite case, when $\lambda_i \gg R, T_q$ the nonpotential description in terms of spatially homogeneous condensates can be applied.

At zero temperature the $O(4)$ invariance of Euclidean space-time holds¹ and FC (2.2) is represented through two scalar functions $D(z)$, $D_1(z)$ (where $z \equiv x - y$) as follows

$$\begin{aligned} D_{\mu\nu, \lambda\sigma}(z) &= (\delta_{\mu\lambda} \delta_{\nu\sigma} - \delta_{\mu\sigma} \delta_{\nu\lambda}) D(z) + \frac{1}{2} \left[\frac{\partial}{\partial z_\mu} (z_\lambda \delta_{\nu\sigma} - z_\sigma \delta_{\nu\lambda}) + \right. \\ &\quad \left. + \frac{\partial}{\partial z_\nu} (z_\sigma \delta_{\mu\lambda} - z_\lambda \delta_{\mu\sigma}) \right] D_1(z). \end{aligned} \quad (2.3)$$

One has to distinguish from the very beginning perturbative and np parts of the correlators $D(z)$, $D_1(z)$. Beginning with the former, one easily finds at tree level

$$D^{p,0}(z) = 0 ; D_1^{p,0}(z) = C_2(f) \frac{4\alpha_s}{\pi} \frac{1}{z^4} \quad (2.4)$$

where $C_2(f)$ is fundamental Casimir $C_2(f) = (N_c^2 - 1)/2N_c$. At higher orders situation becomes more complicated. Namely, one has at n -loop order

$$D^{p,n}(z) = D_1^{p,0}(z) \cdot G^{(n)}(z) ; D_1^{p,n}(z) = D_1^{p,0}(z) \cdot G_1^{(n)}(z) \quad (2.5)$$

where the gauge-invariant functions $G^{(n)}(z)$, $G_1^{(n)}(z)$ has the following general structure:

$$G^{(n)}(z), G_1^{(n)}(z) \sim \alpha_s^n [c_n (\log \mu z)^n + \dots]$$

Thus we assume (and confirm a posteriori) that the following decomposition takes place (vanishing of the perturbative part of $D(z)$ is shown in [12])

$$D(z) = D^{np}(z) ; D_1(z) = D_1^p(z) + D_1^{np}(z) \quad (2.6)$$

¹All treatment in this paper as well as averaging over vacuum fields is done in the Euclidean space. Notice that only after all Green's functions are computed, analytic continuation to Minkowskii space-time can be accomplished.

and $D(z)$ has a smooth limit when $z \rightarrow 0$. Notice that by $D_1(0)$ we always understand np "condensate" part in what follows:

$$G_2 = \frac{6N_c}{\pi^2} (D^{np}(0) + D_1^{np}(0)) \quad (2.7)$$

As is shown below, the correlation lengths λ and λ_1 of $D(z)$ and $D_1^{np}(z)$ respectively appear to be just inverse masses of the corresponding gluelumps $\lambda_j = 1/M_j$.

Having analytic expressions for FCs one might ask how to check them versus experimental and lattice data. On experimental side in hadron spectroscopy one measures masses and transition matrix elements, which are defined by dynamical equation and the latter can be used of potential type due to smallness of λ_j . In static potential only integrals over distance enter and the spin-independent static potential can be written as [3]

$$V_{Q\bar{Q}}(r) = 2 \int_0^r (r-\lambda) d\lambda \int_0^\infty dv D^E(\sqrt{\lambda^2+v^2}) + \int_0^r \lambda d\lambda \int_0^\infty D_1^E(\sqrt{\lambda^2+v^2}). \quad (2.8)$$

Making use of (2.4) together with definition of the string tension [3],

$$\sigma = \frac{1}{2} \int \int d^2z D(z), \quad (2.9)$$

one obtains from (2.8) the standard form of the static potential for $N_c = 3$ at distances $r \gg \lambda$

$$V_{Q\bar{Q}}(r) = \sigma r - \frac{4\alpha_s(r)}{3r} + \mathcal{O}(\lambda/r, \alpha_s^2). \quad (2.10)$$

As is argued below, $\lambda \approx 0.1$ Fm, so that the form (2.10) is applicable in the whole range of distances $r > 0.1$ fm provided asymptotic freedom is taken into account in $\alpha_s(r)$ in (2.10).

3. Gluelumps and FC

In this section we establish a connection of FC $D(z)$ and $D_1(z)$ with the gluelump Green's functions.

To proceed one can use the background field formalism [13][14], where the notions of valence gluon field a_μ and background field B_μ are introduced, so that total gluonic field A_μ is written as

$$A_\mu = a_\mu + B_\mu. \quad (3.1)$$

The main idea we are going to adopt here is suggested in [15].

Now we turn to the analytic calculation of FC in terms of gluelump Green's function. To this end we insert (3.1) into (2.2) and have for $F_{\mu\nu}(x)$:

$$F_{\mu\nu}(x) = \partial_\mu A_\nu - \partial_\nu A_\mu - ig[A_\mu, A_\nu] = \hat{D}_\mu a_\nu - \hat{D}_\nu a_\mu - ig[a_\mu, a_\nu] + F_{\mu\nu}^{(B)}. \quad (3.2)$$

Here, the term $F_{\mu\nu}^{(B)}$ contains only the field B_μ^b . It is clear that when one averages over field a_μ^a and sums finally over all color indices a , one actually exploits all the fields with color indices from

$F_{\mu\nu}^{(B)}$, so that the term $F_{\mu\nu}^{(B)}$ can be omitted, if summing over all indices a is presumed to be done at the end of calculation (see [15] for more discussion).

As a result $D_{\mu\nu,\lambda\sigma}$ can be written as

$$D_{\mu\nu,\lambda\sigma}(x,y) = D_{\mu\nu,\lambda\sigma}^{(0)} + D_{\mu\nu,\lambda\sigma}^{(1)} + D_{\mu\nu,\lambda\sigma}^{(2)} \quad (3.3)$$

where the superscript 0,1,2 refers to the power of g coming from the term $ig[a_\mu, a_\nu]$.

We can address now an important question about the relation between FCs and gluelump Green's functions. We begin with 1-gluon gluelump, whose Green's function reads

$$G_{\mu\nu}^{(1g)}(x,y) = \langle \text{Tr}_a a_\mu(x) \hat{\Phi}(x,y) a_\nu(y) \rangle. \quad (3.4)$$

As shown in [16], the first term in (3.5) is connected to the functions D_1 and it can be written as follows

$$D_{\mu\nu,\lambda\sigma}^{(0)}(x,y) = \frac{g^2}{2N_c^2} \left\{ \frac{\partial}{\partial x_\mu} \frac{\partial}{\partial y_\lambda} G_{\nu\sigma}^{(1g)}(x,y) + perm \right\} + \Delta_{\mu\nu,\lambda\sigma}^{(0)}, \quad (3.5)$$

where $\Delta_{\mu\nu,\lambda\sigma}^{(0)}$ contains contribution of higher FCs, which we systematically discard.

On the other hand one can find $G_{\mu\nu}^{(1g)}(x,y)$ from the FFSR for $G_{\mu\nu}^{(1g)}$ [11] expression [?] written as

$$G_{\mu\nu}^{(1gl)}(x,y) = \text{Tr}_a \int_0^\infty ds (Dz)_{xy} \exp(-K) \langle W_{\mu\nu}^F(C_{xy}) \rangle, \quad (3.6)$$

where the spin-dependent W-loop is

$$W_{\mu\nu}^F(C_{xy}) = PP_F \left\{ \exp(ig \int B_\lambda dz_\lambda) \exp F \right\}_{\mu\nu} \quad (3.7)$$

and the gluon spin factor is $\exp F \equiv \exp(2ig \int_0^s d\tau \hat{F}_B(z(\tau)))$ with \hat{F}_B made of the background field B_μ only.

Analogous expression can be constructed for Green's function of 2-gluon gluelump. It is given by the following expression

$$\begin{aligned} G_{\mu\nu,\lambda\sigma}^{(2gl)}(x,y) &= \langle \text{Tr}_a (f^{abc} f^{def} a_\mu^a(x) a_\nu^b(x) T^c \hat{\Phi}(x,y) T^f \times \\ &\times a_\lambda^d(y) a_\sigma^e(y) \rangle \equiv N_c^2 (N_c^2 - 1) (\delta_{\mu\lambda} \delta_{\nu\sigma} - \delta_{\mu\sigma} \delta_{\nu\lambda}) G^{(2gl)}(z). \end{aligned} \quad (3.8)$$

At small distances $G^{(2gl)}(x,y)$ is dominated by perturbative expansion terms,

$$G^{(2gl)}(z) \sim \frac{1}{z^4}, \quad (3.9)$$

however, as we already discussed in details, all these perturbative terms are canceled by those from higher FCs [12] (triple, quartic, etc...), therefore expansion in fact starts with np terms of dimension four.

To identify the np contribution to $G^{(2gl)}(z)$ we rewrite it as follows:

$$G^{(2gl)}(z) = \int_0^\infty ds_1 \int_0^\infty ds_2 (Dz_1)_{0x} (Dz_2)_{0x} \text{Tr} W_\Sigma(C_1, C_2) \quad (3.10)$$

where the two-gluon gluelump W-loop $W_\Sigma(C_1, C_2)$ can be written as (in the Gaussian approximation)

$$\text{Tr}W_\Sigma(C_1, C_2) = \exp \left\{ -\frac{1}{2} \int_S \int d\pi_{\mu\nu}(u) d\pi_{\lambda\sigma}(v) \langle \hat{F}_{\mu\nu}(u) \hat{\Phi} \hat{F}_{\lambda\sigma}(v) \rangle \right\}. \quad (3.11)$$

and the total surface S consists of 3 pieces, as shown in Fig.1

$$F_{\mu\nu} d\pi_{\mu\nu}(u) = F_{\mu\nu} ds_{\mu\nu}(u) - 2ig d\tau(\hat{F}(u)) \quad (3.12)$$

where (\hat{F}) has Lorentz tensor and adjoint color indices F_{ij}^{ab} and lives on gluon trajectories, i.e. on the boundaries of S_i . In full analogy with (3.5) we have

$$D(z) = \frac{g^4(N_c^2 - 1)}{2} G^{(2gl)}(z) \quad (3.13)$$

The crucial point is that the Green's function (3.6), (3.8) can be calculated in terms of the same FCs $D(z)$, $D_1(z)$. Indeed one has

$$G^{(w)}(z) = \langle f | \exp(-H_w|z) | i \rangle \quad (3.14)$$

where the index w stays for 1-gluon or 2-gluon gluelump Hamiltonians. The latter are expressed via the same FCs $D(z)$, $D_1(z)$ (see [17] and references therein):

$$H_w = H_0[\mu] + \Delta H_L[\mu, \nu] + \Delta H_{Coul}[D_1] + \Delta H_{string}[D, \nu] \quad (3.15)$$

where the last term $H_{string}[D, \nu]$ depends on $D(z)$ via adjoint string tension

$$\sigma_{adj} = \frac{9}{4} \int_0^\infty d^2z D(z) \quad (3.16)$$

and Hamiltonian depends on einbein fields μ and ν .

The self-consistent regimes correspond to different asymptotics of the solutions to these equations. In Coulomb phase of a gauge theory both H_{1g} , H_{2g} exhibit no mass gap, i.e. large z asymptotic of (3.14) is power-like. The function $D(z)$ vanishes in this phase and W-loop obeys perimeter law. In the confinement phase realized in Yang-Mills theory at low temperatures a typical large- z pattern is given by

$$D(z), D_1(z) \sim \exp(-|z|/\lambda_i)$$

i.e. there is a mass gap for both H_{1g} , H_{2g} .

Confining solutions are characterized by W-loops obeying area law. In other words, Hamiltonians expressed in terms of interaction kernels depending on $D(z)$, $D_1(z)$ exhibit mass gap if these kernels are confining. On the other hand the same mass gap plays a role of inverse correlation length of the vacuum. This should be compared with well known mean-field technique. In our case the role of mean-field is played by quadratic FC, which develops non-trivial Gaussian term $D(z)$. Notice that the exponential form of its large distance asymptotics $\exp(-|z|/\lambda)$ (and not, let's say, $\exp(-z^2/\lambda^2)$) is dictated by spectral expansion of the corresponding Green's function at large distances.

Full solution of the above equations is a formidable task not addressed by us here. Instead, as a necessary prerequisite we check below different asymptotic regimes and demonstrate self-consistency of the whole picture.

We begin with small distance region. The np part of contribution to $D_1(z)$ at small distances comes from the small distance (small loop) contribution of the Wilson loop with the area S [3], $W(C) = \exp(-\frac{\pi^2}{8}G_2S^2)$.

As a result one obtains

$$D_1(z) = \frac{4C_2(f)\alpha_s}{\pi} \frac{1}{z^4} + \frac{g^2}{12}G_2. \quad (3.17)$$

It is remarkable that the sign of the np correction is positive.

At large distances one can use the gluelump Hamiltonian for one-gluon gluelump from [17] to derive the asymptotics

$$D_1(z) = \frac{2C_2(f)\alpha_s M_0^{(1)} \sigma_{adj}}{|z|} e^{-M_0^{(1)}|z|}, \quad |z|M_0^{(1)} \gg 1. \quad (3.18)$$

where $M_0^{(1)} = (1.2 \div 1.4)$ GeV for $\sigma_f = 0.18$ GeV² [17, 18].

We now turn to the FC $D(z)$ as was studied in [16]. The relation (3.13) connects $D(z)$ to the two-gluon gluelump Green's function and we shall write it in the form

$$G^{(2gl)}(z) = G_p^{(2gl)}(z) + G_{np}^{(2gl)}(z) \quad (3.19)$$

where $G_p^{(2gl)}(z)$ contains purely perturbative contributions which are subtracted by higher-order FCs, while $G_{np}^{(2gl)}(z)$ contains np and possible perturbative - np interference terms. We are interested in the contribution of the FC $\langle FF \rangle$ to $G^{(2gl)}(z)$, when z tends to zero.

One can envisage two types of contributions:

a) In line with the treatment of D_1 , Eq. (3.17), one can write the term, representing two-gluon gluelump as two nearby one-gluon gluelumps which yields for $D(z)$,

$$\Delta D_a(z) = -\frac{g^4 N_c G_2}{4\pi^2} \quad (3.20)$$

b) in this case one should consider the diagram given in Fig. 1, which yields the answer (for details see Appendix of [5]).

$$\Delta D_b(z) = 2N_c^2 g^4 h(z). \quad (3.21)$$

where at small z , $h(z) \approx \frac{D(\lambda_0)}{64\pi^4} \log^2\left(\frac{\lambda_0\sqrt{e}}{z}\right)$, λ_0 is of the order of correlation length λ .

At large distances one use the two-gluon gluelump Hamiltonian as in [17] and find the corresponding spectrum and wave functions, see [17] and appendix 5 of [16] for details. As a result one obtains in this approximation

$$D(z) = \frac{g^4(N_c^2 - 1)}{2} 0.1 \sigma_f^2 e^{-M_0^{(2)}|z|}, \quad M_0^{(2)}|z| \gg 1 \quad (3.22)$$

where $M_0^{(2)}$ is the lowest two-gluon gluelump mass found in [17, 18] to be about $M_0^{(2)} = (2.5 \div 2.6)$ GeV.

We shall discuss the resulting properties of $D(x)$ and $D_1(x)$ in the next section.

4. Discussion of Consistency

We start with the check consistency for $D(z)$. As is shown above, $D(z)$ has the following behavior at small z

$$D(z) \approx -4N_c \alpha_s^2(\mu(z)) G_2 + N_c^2 \frac{\alpha_s^2(\mu(z))}{2\pi^2} D(\lambda_0) \log^2 \left(\frac{\lambda_0 \sqrt{e}}{z} \right) \quad (4.1)$$

Since $\alpha_s(\mu(z)) \sim 2\pi/\beta_0 \log(\Lambda z)^{-1}$ the first term is subleading at $z \rightarrow 0$ and the last term on the r.h.s. tends to a constant:

$$D(0) = \frac{N_c^2}{2\pi^2} D(\lambda_0) \left(\frac{2\pi}{\beta_0} \right)^2 \quad (4.2)$$

Since from (4.2) one can infer, that $D(0) \approx 0.15D(\lambda_0)$ for $N_c = 3$, where $\lambda_0 \gtrsim \lambda^E$. So $D(z)$ is an increasing function of z at small z , $z \lesssim \lambda_0$, and for $z \gg \lambda$ one observes exponential falloff. The qualitative picture illustrating this solution for $D(z)$ is shown in Fig.2.

This pattern may solve qualitatively the contradiction between the values of $D(0)$ estimated from the string tension $D_\sigma(0) \simeq \frac{\sigma}{\pi\lambda^2} \approx 0.35 \text{ GeV}^4$ and the value obtained in naive way from the gluon condensate $D_{G_2}(0) = \frac{\pi^2}{18} G_2 \approx (0.007 \div 0.012) \text{ GeV}^4$. One can see that $D_\sigma(0) \approx (30 \div 54)D_{G_2}(0)$. This seems to be a reasonable explanation of the mismatch discussed in the introduction.

As was shown in [16], the large distance exponential behavior is selfconsistent, since (assuming that it persists for all z , while small z region contributes very little) from the equality $\sigma = \pi\lambda^2 D_\sigma(0)$, comparing with (3.22) one obtains

$$0.1 \cdot 8\pi^2 \alpha_s^2(N_c^2 - 1) \sigma_f^2 = \frac{\sigma_f}{\pi\lambda^2} \quad (4.3)$$

where in $\alpha_s(\mu)$ the scale μ corresponds roughly to the gluelump average momentum (inverse radius) $\mu_0 \approx 1 \text{ GeV}$. Thus (4.3) yields $\alpha_s(\mu_0) \approx 0.4$ which is in reasonable agreement with α_s from other systems .

We end this section with discussion of three points:

1) $D(z)$ and $D_1(z)$ have been obtained here in the leading approximation, when gluelumps of minimal number of gluons contribute: 2 for $D(z)$ and 1 for $D_1(z)$. In the higher orders of $O(\alpha_s)$ one has an expansion of the type

$$\begin{aligned} D(z) &= D^{(2gl)}(z) + c_1 \alpha_s^3 D^{(1gl)}(z) + c_2 \alpha_s^3 D^{(3gl)}(z) + \dots \\ D_1(z) &= D^{(1gl)}(z) + c'_1 \alpha_s^3 D^{(2gl)}(z) + \dots \end{aligned} \quad (4.4)$$

Hence the asymptotic behavior for $D(z)$ will contain exponent of $M_0^{(1)}|z|$ too, but with a small preexponent coefficient.

2) The behavior of $D(z), D_1^{(np)}(z)$ at small z is defined by NP terms of dimension four, which are condensate G_2 , and the similar term from the expansion of $\exp F$, namely,

$$\langle \exp F \rangle = 1 + 4g^2 \int_0^s d\tau \int_0^{s'} d\tau' \langle F(u(\tau)) F(u(\tau')) \rangle + \dots \quad (4.5)$$

therefore one does not encounter mixed terms like $O(\frac{m^2}{z^2})$.

3) correlation lengths λ, λ_1 are found from gluelump masses,

$$\lambda_1 = 1/M_0^{(1)}, \quad M_0^{(1)} \approx 1.2 \div 1.4 \text{ GeV}, \quad \lambda_1 \cong 0.2 \div 0.15 \text{ fm}. \quad (4.6)$$

The value $M_0^{(1)}$ in (4.6) is taken from the calculations in [17]. The same is true for D , as it is seen from (3.8), where $G^{(2gl)}$ corresponds to two-gluon subsystem angular momentum $L = 0$ independently of $\mu\nu, \lambda\sigma$.

Hence one obtains

$$\lambda \equiv \frac{1}{M_0^{(2)}}; \quad \lambda \cong 0.08 \text{ fm}, \quad M_0^{(2)} \approx 2.5 \text{ GeV}. \quad (4.7)$$

where the value $M_0^{(2)} \approx 2.5 \text{ GeV}$ is taken from [17].

5. Conclusions

We have derived, following the method of [13, 14] the expressions for FC $D(z), D_1(z)$ in terms of gluelump Green's functions. This is done in Gaussian approximation. The latter are calculated using Hamiltonian where np dynamics is given by $D^{np}(z), D_1^{np}(z)$. In this way one obtains selfcoupled equations for these functions, which allow two types of solutions: 1) $D^{np}(z) = 0, D_1^{np}(z) = 0$, i.e. no np effects at all; 2) $D^{np}(z), D_1^{np}(z)$ are nonzero and defined by the only scale, which should be given in QCD, e.g. string tension σ or Λ_{QCD} . All other quantities are defined in terms of these basic ones. We have checked consistency of selfcoupled equations at large and small distances and found that to the order $\mathcal{O}(\alpha_s)$ no mixed perturbative- np terms appear. The function $D_1(z)$ can be represented as a sum of perturbative and np terms, while $D(z)$ contains only np contributions.

We have found a possible way to explain the discrepancy between the average values of field strength taken from G_2 and from σ by showing that $D(z)$ has a local minimum at $z = 0$ and grows at $z \sim \lambda$. Small value of λ and large value of the gluelump mass $M_{gl} = 1/\lambda \approx 2.5 \text{ GeV}$ explains the lattice data for $\lambda \approx 0.1 \text{ Fm}$. Thus the present paper argues that relevant degrees of freedom ensuring confinement are gluelumps, described self-consistently in the language of FCs.

Acknowledgments

The work is supported by the grant for support of scientific schools NS-4961.2008.2.

References

- [1] Yu.A.Simonov, Phys. Uspekhi, **39** (1996) 313.
- [2] D.S.Kuzmenko, V.I.Shevchenko, Yu.A.Simonov, Phys. Uspekhi, **47** (2004) 1.
- [3] H.G.Dosch, Phys. Lett. **B 190** (1987) 177; H.G.Dosch and Yu.A.Simonov, Phys. Lett. **B 205** (1988) 339; Yu.A.Simonov, Nucl. Phys. **B 307** (1988) 512; A.Di. Giacomo, H.G.Dosch, V.I.Shevchenko and Yu. A.Simonov, Phys.Rept., **372** (2002) 319, hep-ph/0007223.
- [4] J. Ambjorn and P. Olesen, Nucl. Phys. **B 170** (1980) 60.
- [5] V.I.Shevchenko, Yu.A.Simonov, arXiv: 0902.1405

- [6] M.A.Shifman, A.I.Vainshtein, V.I.Zakharov, Nucl. Phys. **B 147** (1979) 385, 448.
- [7] A.V.Nefediev, Yu.A.Simonov and M.A.Trusov, Int. Journ. Mod. Phys. **E 18** (2009) 549.
- [8] V.I.Shevchenko, Yu.A.Simonov, Phys. Rev. Lett. **85** (2000) 1811; Int. J. Phys. **A 18** (2003) 127; G.Bali, Phys. Rev. **D 62** (2000) 114503.
- [9] A.Di Giacomo and H.Panagopoulos, Phys. Lett. **B 285** (1992) 133; M.D'Elia, A.Di Giacomo, and E.Meggiolaro, Phys. Lett. **B 408** (1997) 315; A.Di Giacomo, E.Meggiolaro and H.Panagopoulos, Nucl. Phys. **B 483** (1997) 371; M.D'Elia, A.Di Giacomo, and E.Meggiolaro, Phys. Rev. **D 67** (2003) 114504; G.S.Bali, N.Brambilla and A Vairo, Phys. Lett. **B 42** (1998) 265.
- [10] Yu.A.Simonov, Yad. Fiz. **54** (1991) 192; Yu.A.Simonov, in: "QCD: Perturbative or nonperturbative?" eds. L.Ferreira., P.Nogueira, J.I.Silva-Marcos, World Scientific, 2001, hep-ph/9911237.
- [11] Yu.A.Simonov, J.A.Tjon, Ann. Phys. **300** (2002) 54.
- [12] V.I.Shevchenko, Yu.A.Simonov, Phys. Lett. **437** (1998) 146.
- [13] B.S.De Witt, Phys. Rev. **162** (1967) 1195, 1239; J.Honerkamp, Nucl. Phys. **B 48** (1972) 269; G.'t Hooft Nucl. Phys. **B 62** (1973) 444, Lectures at Karpacz, in : Acta Univ. Wratislaviensis **368** (1976) 345; L.F.Abbot, Nucl. Phys. **B 185** (1981) 189
- [14] Yu.A.Simonov, Phys. At Nucl. **58** (1995) 107, hep-ph/9311247; Yu.A.Simonov, JETP Lett. **75** (1993) 525; Yu.A.Simonov, in: Lecture Notes in Physics v.479, p. 139; ed. H.Latal and W.Schweiger, Springer, 1996.
- [15] Yu.A.Simonov, Phys. Atom. Nucl. **61** (1998) 855, arXiv: hep-ph/9712250.
- [16] D.V.Antonov, Yu.A.Simonov, Phys. Atom Nucl. **60** (1997) 478, arXiv:hep-ph/9507254; Yu.A.Simonov, Phys. Atom. Nucl. **61** (1998) 855, arXiv: hep-ph/9712250.
- [17] Yu.A.Simonov, Nucl. Phys. **B 592** (2001) 350.
- [18] M.Foster, C.Michael, Phys. Rev. **D 59** (1999) 094509.





Fig.1



1

Fig. 2

POS(QCD-TNT09)042