

Inclusive diffraction at HERA with a measured leading proton

Florian Goebel* (on behalf of the H1 and ZEUS Collaborations)

Address: DESY, Notkestr. 85, 22607 Hamburg, Germany

E-mail: fgoebel@mail.desy.de

ABSTRACT: Diffractive dissociation of virtual photons, $\gamma^*p \rightarrow Xp$, has been studied in ep interactions at HERA with the H1 and ZEUS detectors. The scattered proton is measured in spectrometers positioned downstream around the proton beam pipe. The cross section is presented as a function of the four-momentum transfer squared t and the azimuthal angle between the electron scattering plane and the proton scattering plane. The data are analyzed in terms of the diffractive structure function $F_2^{D(3)}(x_P, \beta, Q^2)$.

1. Introduction

The diffractive dissociation of real photons, $\gamma p \rightarrow Xp$, has characteristics similar to those of the dissociation of hadrons and can be described by Regge phenomenology. Within this framework, diffractive reactions at high energy are dominated by the exchange of a trajectory with the quantum numbers of the vacuum, referred to as the Pomeron trajectory. HERA allows the investigation of the partonic nature of diffraction in deep-inelastic scattering (DIS) using virtual photons as a probes. It is expected that the diffractive dissociation of virtual photons with high virtualities Q^2 can be described in terms of perturbative QCD.

In diffractive DIS, $ep \rightarrow eXN$, the photon dissociation system X with mass M_X is usually reconstructed in the central detector and is well separated in rapidity from the proton remnant system N which, for small masses M_N , escapes through the forward beam hole of the detector. Diffractive events have been selected either by requiring an absence of particles in the forward region of the central detector (large rapidity gap method) [1] or by making use of the characteristic distribution of the mass M_X . However, for the analyses presented here, the scattered proton is measured in the H1 forward proton spectrometer (FPS) or the ZEUS leading proton spectrometer (LPS). The detection of the final state proton not only allows the measurement of its momentum but also provides the cleanest selection of diffractive events, $ep \rightarrow eXp$, independent of the hadronic final state and free from proton dissociation background.

*Speaker.

2. Analyses

Protons scattered at small angles are deflected by the magnets of the beam optics into a system of detectors which are placed close to the proton beam to measure the four-momentum of the scattered proton. The H1 FPS consists of four stations of scintillating fiber hodoscopes installed between 64 m and 90 m from the interaction point. The ZEUS LPS is a set of six silicon microstrip stations situated 24 m to 90 m downstream of the interaction point. Diffractive DIS events are selected by requiring a reconstructed positron in the central detector and a measured leading proton in the FPS or LPS. The kinematic variables Q^2 , x_{IP} , β and t are reconstructed from the scattered positron, the hadronic final state measured in the central detector and the scattered proton. The H1 analysis is based on a luminosity of 28.8 pb^{-1} and is restricted to the kinematic range $2 < Q^2 < 50 \text{ GeV}^2$, $x_{IP} < 0.09$ and $0.08 < |t| < 0.45 \text{ GeV}^2$ while the ZEUS analysis uses 12.8 pb^{-1} and is restricted to $4 < Q^2 < 100 \text{ GeV}^2$, $x_{IP} < 0.03$ and $0.075 < |t| < 0.3 \text{ GeV}^2$.

3. Results

3.1 t -dependence




Figure 1: *Left:* The differential cross section $d\sigma/dt$ measured with the H1 FPS in four different x_{IP} bins. Results of the fit with a function $d\sigma/dt \propto e^{bt}$ are shown. *Right:* The slope parameter b is plotted as a function of x_{IP} . Results obtained with the H1 FPS and the ZEUS LPS are shown.

A typical feature of diffractive events is an exponential fall of the differential cross section with $|t|$ at small values of $|t|$. In Fig. 1, the cross section is parameterized as $d\sigma/dt \propto e^{b|t|}$ in bins of x_{IP} , and the slope parameter b is plotted as a function of x_{IP} . The results of H1[2] and ZEUS[3] are consistent within the experimental uncertainties and no significant dependence on x_{IP} is visible.




Figure 2: The diffractive structure function $x_{IP}F_2^{D(3)}$ as obtained by H1 and ZEUS is plotted as a function of x_{IP} in bins of β and Q^2 .

3.2 Diffractive structure function

The diffractive cross section for the process $ep \rightarrow eXp$ can be expressed in terms of the diffractive structure function $F_2^{D(3)}$ as $\frac{d\sigma^D}{d\beta dQ^2 dx_{IP}} = \frac{2\pi\alpha^2}{\beta Q^4} [1 + (1-y)^2] F_2^{D(3)}(\beta, Q^2, x_{IP})$, where $x_{IP} = (Q^2 + M_X^2)/(Q^2 + W^2)$ is the fraction of the proton momentum carried by the Pomeron and $\beta = Q^2/(Q^2 + M_X^2)$ is the fraction of the Pomeron momentum carried by the struck quark. In Fig. 2, the results for $x_{IP}F_2^{D(3)}$ obtained by H1[2] and ZEUS[3] are shown as a function of x_{IP} . The data are extrapolated to the full t range using the t dependence measured by each experiment. Good agreement is observed within uncertainties.

In Fig. 3, the values of $x_{IP}F_2^{D(3)}$ obtained with the requirement of a leading proton are compared to the recent H1 measurement [4] using a large rapidity gap in the central detector. Within uncertainties, the leading proton data are in good agreement with the rapidity gap data indicating that the contribution of proton dissociation to the latter analysis is small.

In factorisable models, $F_2^{D(3)}$ factorises into a Pomeron flux $f_{IP}(x_{IP})$ and the structure function $F_2^{D(2)}(\beta, Q^2)$. Assuming a Regge-inspired power law dependence of $f_{IP}(x_{IP}) \propto (1/x_{IP})^{2\alpha_{IP}(t)-1}$, ZEUS found consistency with factorization and obtained $\alpha_{IP}(0) = 1.13 \pm 0.03(stat)_{-0.01}^{+0.03}(syst)$. In Fig. 4, $F_2^{D(2)}$ is shown as a function of β and Q^2 . Clear scaling violations are seen in the low β bins of Fig. 4 (right).




Figure 3:
 $x_{IP} F_2^{D(3)}$ obtained with the FPS is compared to the recent rapidity gap measurement by H1.

3.3 Azimuthal asymmetry

For unpolarised positrons and protons, the diffractive cross section can be decomposed as $d\sigma^D/d\Phi \propto \sigma_T^D + \epsilon\sigma_L^D - 2\sqrt{\epsilon(1+\epsilon)}\sigma_{LT}^D \cos\Phi - \epsilon\sigma_{TT}^D \cos 2\Phi$, where Φ is the angle between the positron and proton scattering planes in the γ^*p rest frame, σ_T^D and σ_L^D are the cross sections for transverse and longitudinal photons, σ_{LT}^D and σ_{TT}^D are interference terms and ϵ is the polarisation parameter. If $\sigma_{LT}^D \neq 0$, the ϕ distribution is non-uniform. A measurement of this non-uniformity would furthermore imply a non-zero value of $R^D = \sigma_{LT}^D/\sigma_T^D$ [5]. Figure 5 (left) shows $d\sigma^D/d\Phi$ as measured by the ZEUS collaboration together with a fit to the form $d\sigma^D/d\Phi \propto 1 + A_{LT} \cos\Phi$ which yields $A_{LT} = -0.049 \pm 0.058(stat.)_{-0.009}^{+0.056}(syst.)$, compatible with zero indicating that the interference term σ_{LT}^D is small. Figure 5 (right) shows a first measurement of A_{LT} in bins of x_{IP} , β , $-t$ and Q^2 . Theoretical predictions for the behavior of A_{LT} [5, 6] mostly concern the high β region, which was not accessible in this analysis. No strong dependence of A_{LT} on x_{IP} , β , t and Q^2 is observed in the data.

References

- [1] P. Newmannn, these proceedings.
- [2] H1 Collaboration, Abstract 809, Paper submitted to *EPS Conf. on HEP*, Budapest 2001.




Figure 4: The values of $F_2^{D(2)}(\beta, Q^2)$, obtained from a fit to $F_2^D(3)$ are shown as a function of β (Left) and Q^2 (Right).




Figure 5: Left: The cross section $d\sigma/d\Phi$ together with the fit described in the text. Right: The azimuthal asymmetry A_{LT} determined as a function of x_{IP} , β , t and Q^2 .

- [3] ZEUS Collaboration, Abstract 566, Paper submitted to *EPS Conf. on HEP*, Budapest 2001.
- [4] H1 Collaboration, Abstract 808, Paper submitted to *EPS Conf. on HEP*, Budapest 2001.
- [5] N.N. Nikolaev et al., *Phys. Rev. D* **59**, 091501 (1999).
- [6] T. Arens et al., *Z. Phys. C* **74**, 651 (1997); T. Gehrman and W.J. Stirling, *Z. Phys. C* **70**, 89 (1996); N.N. Nikolaev and B.G. Zakharov, hep-ph/9706343.