

Determining α_s at NNLO from Event Shape Data

Gionata Luisoni*

Institut für Theoretische Physik, Universität Zürich, CH-8057 Zürich, Switzerland

E-mail: luisonig@physik.uzh.ch

We report on several methodologically different NNLO determinations of the strong coupling constant from event shapes and related observables in e^+e^- annihilation. The event shape distributions are analyzed within a combined framework of NNLO and NNLA resummation. We also investigate the role of hadronisation corrections, using both Monte Carlo generator predictions and analytic models to parametrise non-perturbative power corrections.

XVIII International Workshop on Deep-Inelastic Scattering and Related Subjects, DIS 2010

April 19-23, 2010

Firenze, Italy

*Speaker.

1. Introduction

The reaction of e^+e^- annihilation into 3 jets has played historically a very prominent role for phenomenology and allows a precise determination of the strong coupling constant α_s , since the deviation from two-jet configurations is proportional to it. The phenomenologically interesting observables for the purpose of determining the strong coupling constant can be divided into two big categories. On one side there are jet rates, which rely on a jet algorithm and are based on the number of jets observed in an event, on the other side also the topology of the single events can be studied in a systematic fashion by means of so-called event-shape observables. The popularity of the latter is mainly due to the fact that they are well suited both for experimental measurement and for theoretical description, since many of them are infrared and collinear safe. They describe topological properties of hadronic final states by parameterizing the energy-momentum flow of an event. This class of observables is also interesting for the study of hadronisation effects. Hadronisation corrections usually result in a distortion of the event-shape distributions and are thus not easily disentangled from partonic predictions. However, the study of mean values and higher moments of event shapes allows a clear separation of perturbative and non-perturbative predictions. Furthermore, the comparison of hadronization corrections computed by general purpose Monte Carlo programs with predictions from analytical models permits to shed some light on the effects of hadronization corrections in the determination of α_s .

We have studied jet rates, the distributions and the first five moments of six event-shape observables: thrust T (respectively, $\tau = 1 - T$), heavy jet mass ρ , wide and total jet broadening B_W and B_T , C -parameter and the two-to-three-jet transition parameter in the Durham algorithm, Y_3 . For the definitions of these variables and their historical origin we refer to [1] and references therein. We will denote the variables collectively as y in the following, such that their two jet limit is $y \rightarrow 0$. Recently, with the appearance of the NNLO results, several extractions of α_s have been performed using event-shape observables [2, 3, 4, 5, 6]. Their results are summarized in Fig. 1. In the following we present three of them [3, 5, 4].

2. α_s from Event-Shape Distributions

The fixed-order QCD description of event-shape distributions is given by a perturbative expansion of the form

$$\frac{1}{\sigma_{\text{had}}} \frac{d\sigma}{dy}(y, Q, \mu) = \bar{\alpha}_s(\mu) \frac{dA}{dy}(y) + \bar{\alpha}_s^2(\mu) \frac{dB}{dy}(y, x_\mu) + \bar{\alpha}_s^3(\mu) \frac{dC}{dy}(y, x_\mu) + \mathcal{O}(\bar{\alpha}_s^4), \quad (2.1)$$

where $\bar{\alpha}_s = \frac{\alpha_s}{2\pi}$ and $x_\mu = \frac{\mu}{Q}$, and where A , B and C are the perturbatively calculated coefficients [7] at LO, NLO and NNLO. The distribution is normalised to the total hadronic cross section σ_{had} in e^+e^- annihilation. The dependence of (2.1) on the collision energy is only through α_s and x_μ and the scale dependence of α_s is determined according to the three-loop running of $\alpha_s(\mu)$.

In order to obtain reliable predictions over the full kinematical range, the perturbative fixed-order prediction (2.1) has to be matched with resummation, which is taken into account at next-to-leading logarithmic (NLL) accuracy in the $\ln R$ -matching scheme. For more details about the NLLA+NNLO matching we refer to Ref. [8] and references therein.

We have used the six event-shape observables listed in Section 1 for our fits. The measurements we use have been carried out by the ALEPH collaboration [9] at eight different centre-of-mass (CM) energies between 91.2 and 206 GeV. The perturbative QCD prediction is corrected for hadronisation and resonance decays by means of a transition matrix, which is computed with the MC generators PYTHIA [10], HERWIG [11] and ARIADNE [12], all tuned to global hadronic observables at M_Z [13]. Corrected measurements of event-shape distributions are compared to the theoretical calculation at particle level. For a detailed description of the determination and treatment of experimental systematic uncertainties we refer to Refs. [9, 2, 3].

The combined results of six event-shape variables and eight LEP1/LEP2 CM energies is

$$\alpha_s(M_Z) = 0.1224 \pm 0.0009 (\text{stat}) \pm 0.0009 (\text{exp}) \pm 0.0012 (\text{had}) \pm 0.0035 (\text{theo}) .$$

For the fitted values of the coupling constant as found from event-shape variables calculated at various orders we refer to the figures and tables of [3]. The central value of the result is slightly lower than the central value of 0.1228 obtained from a fit using purely fixed-order NNLO predictions [2]. Furthermore the dominant theoretical uncertainty on $\alpha_s(M_Z)$, as estimated from scale variations, is reduced by 20% compared to NLO+NLLA and the scatter among the values of $\alpha_s(M_Z)$ extracted from the six different event-shape variables is substantially reduced. However, compared to the fit based on purely fixed-order NNLO predictions, the perturbative uncertainty is *increased* in the NNLO+NLLA fit. The reason is that in the two-jet region the NLLA+NLO and NLLA+NNLO predictions agree by construction and therefore the renormalisation scale uncertainty is dominated by the resummation in this region, which results in a larger overall scale uncertainty in the α_s fit.

Apart from the α_s determination using the standard MC generators mentioned above, we used HERWIG++ [14] version 2.3 together with the MCNLO [15] and POWHEG [16] schemes for investigating hadronization corrections. From this study it appears that there are two “classes” of variables. With standard hadronisation corrections from PYTHIA we obtain $\alpha_s(M_Z)$ values some 5% higher for the first class, consisting of T , C and B_T , which still suffer from sizable missing higher order corrections, than for the second class consisting of the ρ , B_W and Y_3 , which have a better perturbative stability [17]. This means that the PYTHIA hadronisation corrections, applied in the α_s fit, might be too small for the first class of variables, resulting in a larger $\alpha_s(M_Z)$ value. For further details of our analysis we refer to Ref. [3].

3. α_s from Moments of Event Shapes

The n th moment of an event-shape observable y is defined by

$$\langle y^n \rangle = \frac{1}{\sigma_{\text{had}}} \int_0^{y_{\text{max}}} y^n \frac{d\sigma}{dy} dy , \quad (3.1)$$

where y_{max} is the kinematically allowed upper limit of the observable. For moments of event shapes, one expects the hadronisation corrections to be additive, such that the cross section can be divided into a perturbative and a non-perturbative contribution, where the non-perturbative contribution accounts for hadronisation effects.

In ref. [4], the dispersive model derived in Refs. [18, 19, 20] has been used and extended to NNLO to estimate hadronisation corrections to event-shape moments by calculating analytical

predictions for power corrections. It introduces only a single new parameter α_0 , which can be interpreted as the average strong coupling in the non-perturbative region:

$$\frac{1}{\mu_I} \int_0^{\mu_I} dQ \alpha_{\text{eff}}(Q^2) = \alpha_0(\mu_I), \quad (3.2)$$

where below the IR cutoff μ_I the strong coupling is replaced by an effective coupling. This dispersive model for the strong coupling leads to a shift in the distributions

$$\frac{d\sigma}{dy}(y) = \frac{d\sigma_{\text{pt}}}{dy}(y - a_y P), \quad (3.3)$$

where the numerical factor a_y depends on the event shape, while P is believed to be universal and scales with the CM energy like μ_I/Q . Insertion of eq. (3.3) into (3.1) leads to

$$\langle y^n \rangle = \int_{-a_y P}^{y_{\text{max}} - a_y P} dy (y + a_y P)^n \frac{1}{\sigma_{\text{tot}}} \frac{d\sigma_{\text{pt}}}{dy}(y) \approx \int_0^{y_{\text{max}}} dy (y + a_y P)^n \frac{1}{\sigma_{\text{tot}}} \frac{d\sigma_{\text{pt}}}{dy}(y). \quad (3.4)$$

From this expression one can extract the non-perturbative predictions for the moments of y .

The expressions derived in [4] match the dispersive model with the perturbative prediction at NNLO QCD. Comparing these expressions with experimental data on event-shape moments, a combined determination of the perturbative strong coupling constant α_s and the non-perturbative parameter α_0 has been performed [4], based on data from the JADE and OPAL experiments [21]. The data consist of 18 points at CM energies between 14.0 and 206.6 GeV for the first five moments of T , C , Y_3 , ρ , B_W and B_T , and have been taken from [22]. For each moment the NLO as well as the NNLO prediction was fitted with $\alpha_s(M_Z)$ and α_0 as fit parameters, except for the moments of Y_3 , which have no leading $\frac{1}{Q}$ power correction and thus are independent of α_0 .

Compared to previous results at NLO, inclusion of NNLO effects results in a considerably improved consistency in the parameters determined from different shape variables, and in a substantial reduction of the error on α_s . Furthermore the theoretical error on the extraction of $\alpha_s(M_Z)$ from ρ , Y_3 and B_W is considerably smaller than from τ , C and B_T . As mentioned above, the moments of the former three shape variables receive moderate NNLO corrections for all n , while the NNLO corrections for the latter three are large already for $n = 1$ and increase with n . Consequently, the theoretical description of the moments of ρ , Y_3 and B_W displays a higher perturbative stability, which is reflected in the smaller theoretical uncertainty on $\alpha_s(M_Z)$ derived from those variables.

In a second step, we combine the $\alpha_s(M_Z)$ and α_0 measurements obtained from different event-shape variables. Taking the weighted mean over all values except B_W and B_T , we obtain at NNLO:

$$\alpha_s(M_Z) = 0.1153 \pm 0.0017(\text{exp}) \pm 0.0023(\text{th}), \quad \alpha_0 = 0.5132 \pm 0.0115(\text{exp}) \pm 0.0381(\text{th}).$$

The moments of B_W and B_T have been excluded here since their theoretical description requires an additional contribution to the non-perturbative coefficient P [4] which is unknown to NNLO.

The average of $\alpha_s(M_Z)$ is dominated by the measurements based on ρ and Y_3 , which have the smallest theoretical uncertainties. From NLO to NNLO [4], the error on $\alpha_s(M_Z)$ is reduced by a factor of two. The error on $\alpha_s(M_Z)$ is clearly dominated by the x_μ variation, while the largest contribution to the error on α_0 comes from the uncertainty on the Milan factor \mathcal{M} [19], which has not been improved in the current study.




Figure 1: Recent determinations of α_s based NNLO predictions taken from Refs. [2, 3, 4, 5, 6].

To quantify the difference of the dispersive model to hadronisation corrections from the legacy generators, we analysed the moments of $(1 - T)$ with hadronisation corrections from PYTHIA. As a result, we obtained fit results for $\alpha_s(M_Z)$ which are typically 4% higher than by using the dispersive model, with a slightly worse quality of the fit. Comparing perturbative and non-perturbative contributions at $\sqrt{s} = M_Z$, we observed that PYTHIA hadronisation corrections amount to less than half the power corrections obtained in the dispersive model, thereby explaining the tendency towards a larger value of $\alpha_s(M_Z)$, since the missing numerical magnitude of the power corrections must be compensated by a larger perturbative contribution.

4. α_s from Jet Rates

Very recently a new determination of α_s was performed using jet rates [5]. Theoretical NNLO predictions for jet rates [23] as a function of the jet resolution parameter y_{cut} are compared to ALEPH data [9] using the Durham jet algorithm, for which the distance measure is given by

$$y_{ij,D} = 2 \min(E_i^2, E_j^2) (1 - \cos \theta_{ij}) / E_{\text{vis}}^2 \quad (4.1)$$

where E_{vis} denotes the energy sum of all particles in the final state. In Ref. [23] it was shown that NNLO predictions for jet rates have only very small hadronization corrections and the theoretical error for $10^{-1} < y_{\text{cut}} < 10^{-2}$ drops below the per-cent relative uncertainty. This motivates a dedicated extraction of α_s . The corrected ALEPH measurements for the three-jet rate are compared to the theoretical calculation at particle level. Values for $\alpha_s(M_Z)$ are obtained by a least-squares fit, performed separately for each y_{cut} value in the range . A nice stability of the result is found up to values of $\ln y_{\text{cut}} \approx -4.5$. As final result the value for $y_{\text{cut}} = 0.02$ is taken, which represents an optimal compromise between minimal systematic uncertainty and stability. The following values for α_s is found:

$$\alpha_s(M_Z) = 0.1175 \pm 0.0020(\text{exp}) \pm 0.0015(\text{th}), \quad (4.2)$$

Results from LEP2 energies give similar central values but larger statistical uncertainties. Combining the errors in quadrature yields $\alpha_s(M_Z) = 0.1175 \pm 0.0025$, which is in excellent agreement

with the latest world average [24]. This verifies the expectations that the three-jet rate is an excellent observable for this kind of analysis, thanks to the good behaviour of its perturbative and non-perturbative contributions over a sizable range of jet-resolution parameters.

5. Acknowledgments

This research was supported in part by the Swiss National Science Foundation (SNF) under contracts PP0022-118864 and 200020-126691.

References

- [1] R. W. L. Jones *et al.*, JHEP **0312** (2003) 007 [arXiv:hep-ph/0312016].
- [2] G. Dissertori *et al.*, JHEP **0802** (2008) 040 [arXiv:0712.0327];
S. Bethke *et al.*, Eur. Phys. J. C **64** (2009) 351 [arXiv:0810.1389].
- [3] G. Dissertori *et al.*, JHEP **0908** (2009) 036 [arXiv:0906.3436].
- [4] T. Gehrmann, M. Jaquier and G. Luisoni, Eur. Phys. J. C **67** (2010) 57 [arXiv:0911.2422].
- [5] G. Dissertori *et al.*, Phys. Rev. Lett. **104** (2010) 072002 [arXiv:0910.4283].
- [6] T. Becher and M. D. Schwartz, JHEP **0807** (2008) 034 [arXiv:0803.0342];
Y. T. Chien and M. D. Schwartz, [arXiv:1005.1644];
R. Abbate, M. Fickinger, A. H. Hoang, V. Mateu and I. W. Stewart, [arXiv:1006.3080].
- [7] A. Gehrmann-De Ridder *et al.*, JHEP **0712** (2007) 094 [arXiv:0711.4711].
- [8] T. Gehrmann, G. Luisoni and H. Stenzel, Phys. Lett. B **664** (2008) 265 [arXiv:0803.0695].
- [9] A. Heister *et al.* [ALEPH Collaboration], Eur. Phys. J. C **35** (2004) 457.
- [10] T. Sjostrand *et al.*, Comput. Phys. Commun. **135** (2001) 238 [hep-ph/0010017].
- [11] G. Corcella *et al.* JHEP **0101** (2001) 010 [hep-ph/0011363].
- [12] L. Lönnblad, Comput. Phys. Commun. **71** (1992) 15.
- [13] R. Barate *et al.* [ALEPH Collaboration], Phys. Rep. **294** (1998) 1.
- [14] O. Latunde-Dada, JHEP **0711** (2007) 040 [arXiv:0708.4390];
O. Latunde-Dada, S. Gieseke and B. Webber, JHEP **0702** (2007) 051 [hep-ph/0612281].
- [15] S. Frixione and B. R. Webber, JHEP **0206** (2002) 029 [hep-ph/0204244].
- [16] P. Nason, JHEP **0411** (2004) 040 [hep-ph/0409146].
- [17] A. Gehrmann-De Ridder *et al.*, JHEP **0905** (2009) 106 [arXiv:0903.4658].
- [18] Y. L. Dokshitzer *et al.*, Nucl. Phys. B **469** (1996) 93 [hep-ph/9512336];
Y. L. Dokshitzer and B. R. Webber, Phys. Lett. B **404** (1997) 321 [hep-ph/9704298].
- [19] Y. L. Dokshitzer *et al.*, JHEP **9805** (1998) 003 [hep-ph/9802381].
- [20] Y. L. Dokshitzer *et al.*, Eur. Phys. J. direct C **1** (1999) 3 [hep-ph/9812487].
- [21] C. Pahl *et al.*, Eur. Phys. J. C **60** (2009) 181, (E) C **62** (2009) 451 [arXiv:0810.2933].
- [22] C. J. Pahl, Ph.D. thesis, MPI Munich 2007.
- [23] A. Gehrmann-De Ridder *et al.*, Phys. Rev. Lett. **100** (2008) 172001 [arXiv:0802.0813].
- [24] S. Bethke, Eur. Phys. J. C **64** (2009) 689 [arXiv:0908.1135].