

Direct photon observables from hydrodynamics and implications on the initial temperature and EoS

Máté Csanád*

Eötvös University, Department of Atomic Physics, Pázmány Péter s. 1/A, H-1117 Budapest, Hungary

E-mail: csanad@elte.hu

The expansion of the strongly interacting quark gluon plasma (sQGP) created in Au+Au collisions at RHIC can be described by hydrodynamical models. Hadrons are created after a freeze-out, thus their distribution describes the final state of the evolution. The earlier stages can be analyzed via penetrating probes like photon observables. These were measured in 2010 and 2011 by the PHENIX experiment. Here we analyze an analytic, 1+3 dimensional perfect relativistic hydrodynamic solution and calculate hadron and photon observables, such as transverse momentum spectra, elliptic flow and correlation (HBT) radii. We find that our model is not incompatible with the data, not even with the direct photon elliptic flow. From fitting the data, we find that early temperatures of the sQGP were well above the quark-hadron transition temperature, in the hottest point, the center of the fireball the temperature may have reached 507 ± 12 MeV. The equation of state of this quark matter can be described by an average speed of sound of 0.36 ± 0.02 . We also predict a photon source that is significantly larger in the *out* direction than in the *side* direction.

*The Seventh Workshop on Particle Correlations and Femtoscopy
September 20 - 24 2011
University of Tokyo, Japan*

*Speaker.

1. A hydrodynamic model

It is a well established picture [1] that a strongly interacting quark gluon plasma is created in relativistic Au+Au collisions of the Relativistic Heavy Ion Collider, and the evolution of this medium can be described by perfect hydrodynamics. The equations of hydrodynamics can be solved numerically, which has the advantage of having arbitrary initial conditions. It is also possible to find analytic solutions with realistic properties. However, very few truly 1+3 dimensional (and not spherically symmetric), relativistic models were compared to data yet. In this paper we analyze the relativistic, ellipsoidally symmetric model of Ref. [2]. Hadronic observables were calculated in Ref. [3], while photonic observables in Ref. [4].

The picture used in hydro models is that the pre freeze-out (FO) medium is described by hydrodynamics, and the post FO medium is that of observed hadrons. Hadronic observables can be calculated via the phase-space distribution at the FO. This will correspond to the hadronic final state or source distribution $S(x, p)$, as detailed in Ref. [3]. The same final state can be achieved with different equations of state or initial conditions [5]. However, photon observables are sensitive to the whole time evolution, thus to initial conditions and equation of state as well.

For the direct photon calculations [4], the key assumption is, that the strongly interacting QCD plasma is thermalized and that its photon spectrum corresponds to a thermal, black-body type of radiation. Such a radiation from sQGP is an approximation, valid if the volume elements are in thermal equilibrium, and needs to be cross-checked against more detailed subsequent calculations. Black-body radiation is however a frequent first approximation for the photon radiation of complex bodies like stars and planets, even though these bodies are frequently not in thermal equilibrium with their surroundings. The application of this approximation for the fireballs created in high energy heavy ion collisions is supported by the simple, near exponential experimentally observed direct photon spectrum [6]. Thus the phase-space distribution of the photons is characterized by the temperature of the medium (at a given space-time cell), while the expansion of the fireball also effects the observed spectrum. This is a macroscopic model, and in the following we will calculate photon observables from it and compare to RHIC data. The most important assumption is, that the spectrum of direct photons is thermal because macroscopically, the photon radiation is thermal.

The analyzed solution [2] assumes self-similarity and ellipsoidal symmetry, as described also in Refs. [3, 4]. The ellipsoidal symmetry means that at a given proper time the thermodynamical quantities are constant on the surface of expanding ellipsoids. The ellipsoids are given by constant values of the scale variable s , defined as

$$s = \frac{r_x^2}{\dot{X}^2 t^2} + \frac{r_y^2}{\dot{Y}^2 t^2} + \frac{r_z^2}{\dot{Z}^2 t^2}, \quad (1.1)$$

where the constants \dot{X} , \dot{Y} , and \dot{Z} describe the expansion rate of the fireball in the three spatial directions. Spatial coordinates are r_x , r_y , and r_z . The velocity-field is described by an isotropic Hubble-type expansion, $u^\mu(x) = x^\mu / \tau$, where x^μ means the four-vector (t, r_x, r_y, r_z) and $\tau = \sqrt{t^2 - r_x^2 - r_y^2 - r_z^2}$ is the proper-time coordinate.

The temperature distribution $T(x)$ is given as

$$T(x) = T_0 \left(\frac{\tau_0}{\tau} \right)^{3/\kappa} \exp \left(\frac{bs}{2} \right), \quad (1.2)$$

where τ is the proper time, s is the above scaling variable, while $T_0 = T|_{s=0, \tau=\tau_0}$, and τ_0 is an arbitrary proper time, but practically we choose it to be the time of the freeze-out, thus T_0 is the central freeze-out temperature. Parameter b is proportional to the temperature gradient, i.e. if the fireball is the hottest in the center, then $b < 0$. If there is a conserved charge in the system e.g. the baryon number density, then charge number density $n(x)$ can be utilized in the solution. As described in Refs. [2, 3], such a number density can be introduced as

$$n(x) = n_0 \left(\frac{\tau_0}{\tau} \right)^3 \exp\left(-\frac{bs}{2}\right). \quad (1.3)$$

For the momentum distribution of direct photons, this will not be needed, as the only the temperature of the medium (the strongly interacting plasma) governs the creation of photons, not the density (which however plays an important role also in the case of hadron creation). The equation of state (EoS) we use here is $\varepsilon = \kappa p$, with ε being the energy density and p the pressure. Here $\kappa = c_s^{-2}$ (one over speed of sound squared) is the main parameter describing the EoS.

From the above hydrodynamic quantities, source functions can be created. For bosonic hadrons, it takes the following form [3]:

$$S(x, p)d^4x = \mathcal{N} \frac{p_\mu d^3\Sigma^\mu(x) H(\tau) d\tau}{n(x) \exp(p_\mu u^\mu(x)/T(x)) - 1}, \quad (1.4)$$

where $\mathcal{N} = g/(2\pi)^3$ (with g being the degeneracy factor), $H(\tau)$ is the proper-time probability distribution of the FO. It is assumed to be a δ function or a narrow Gaussian centered at the freeze-out proper-time τ_0 . Furthermore, $\mu(x)/T(x) = \ln n(x)$ is the fugacity factor and $d^3\Sigma_\mu(x)p^\mu$ is the Cooper-Frye factor [7] describing the flux of the particles, and $d^3\Sigma_\mu(x)$ is the vector-measure of the FO hyper-surface. Here the source distribution is normalized such as $\int S(x, p)d^4x d^3\mathbf{p}/E = N$, i.e. one gets the total number of particles N (using $c=1, \hbar=1$ units). Note that one has to change variables from τ to t , and so a Jacobian of $d\tau/dt = t/\tau$ has to be taken into account.

For the source function of photon creation we have [4]:

$$S(x, p)d^4x = \mathcal{N}' \frac{p_\mu d^3\Sigma^\mu(x) dt}{\exp(p_\mu u^\mu(x)/T(x)) - 1} = \mathcal{N}' \frac{p_\mu u^\mu}{\exp(p_\mu u^\mu(x)/T(x)) - 1} d^4x \quad (1.5)$$

where $p_\mu d^3\Sigma^\mu$ is again the Cooper-Frye factor of the emission hyper-surfaces. Similarly to Ref. [3] we assume that the hyper-surfaces are parallel to u^μ , thus $d^3\Sigma^\mu(x) = u^\mu d^3x$. This yields then $p_\mu u^\mu$ which is the energy of the photon in the co-moving system. The photon creation is assumed to happen from an initial time t_i until a point sufficiently near the freeze-out.

Experimental observables can then be calculated from the source function, using a second order saddle-point approximation. In this approximation the point of maximal emissivity is

$$r_{0,i} = \rho_i t \frac{p_i}{E}, \quad \text{for } i = x, y, z \quad (1.6)$$

while the widths of the particle emitting source are

$$R_i^2 = \rho_i \tau_0^2 \frac{T_0}{E} \left(\frac{t}{\tau_0} \right)^{-3/\kappa+2}, \quad \text{for } i = x, y, z \quad (1.7)$$

where we introduced the auxiliary quantities $\rho_i = \kappa/(\kappa - 3 - \kappa b/\dot{R}_i^2)$, where again $\kappa = c_s^{-2}$ is describing the EoS, and $\dot{R}_i = \dot{X}, \dot{Y}, \dot{Z}$ for $i = x, y, z$, respectively.

2. Calculated observables

The invariant one-particle momentum distribution is defined as $N_1(p) = \int S(x, p) d^4x$. It depends on the three-momentum $p = (p_x, p_y, p_z)$. We will introduce the (p_t, φ, p_z) cylindrical coordinates (z being the beam direction) and use the longitudinal rapidity y (for which $E dy = dp_z$ is true). As usual, we will restrict our calculations to $y = 0$ (note that in this case $E = p_t$ is true for photons). Our calculated quantities will then be the elliptic flow v_2 , and the transverse momentum distribution $N_1(p_t)$. These can be calculated from $N_1(p)$ as

$$N_1(p_t) = \int_0^{2\pi} N_1(p)|_{y=0} d\varphi \quad (2.1)$$

$$v_2(p_t) = \frac{\frac{1}{2\pi} \int_0^{2\pi} N_1(p)|_{y=0} \cos(2\varphi) d\varphi}{N_1(p_t)} \quad (2.2)$$

We also calculated Bose-Einstein correlation radii. As usual, the two-particle correlation function for identical particles can be calculated from the single particle source function $S(x, p)$ as

$$C_2(q) = 1 + \lambda \left| \frac{\tilde{S}(q)}{\tilde{S}(0)} \right|^2 \quad (2.3)$$

where q is the momentum difference of the two particles and $\tilde{S}(q)$ is the Fourier-transformed of the source $S(x, p)$ in the variable x (momentum p is usually dropped). This correlation function has, as usual, a shape with a peak, the width of which is characterized by the HBT radii R_{out} , R_{side} and R_{out} . We calculated these radii for different average momenta p .

Here we do not detail the analytic result of these calculation, but will show a comparison of the model to the data in the next section. The detailed results are given in Refs. [3, 4]. It is important to note however, that in the final formulas, we use transverse expansion (u_t) and eccentricity (ε) instead of expansion rates \dot{X} and \dot{Y} : $u_t^{-2} = 0.5(1/\dot{X}^2 + 1/\dot{Y}^2)$, $\varepsilon = (\dot{X}^2 - \dot{Y}^2)/(\dot{X}^2 + \dot{Y}^2)$.

3. Comparison to the measured direct photon spectrum

The freeze-out parameters were determined from hadronic fits in Ref. [3]. These properties include the expansion rates, the freeze-out proper-time and freeze-out temperature (in the center of the fireball), as shown in Table 1. When describing direct photon data, we used the parameters of the hadronic fit and left only the remaining as free parameters [4]. The free parameters will be κ (the equation of state parameter) and t_i , the initial time of the evolution.

We compared the above results to PHENIX 200 GeV Au+Au hadron and photon data. We fitted our formulas to PHENIX invariant transverse momentum distributions [8], HBT radii [9] and elliptic flow data [10]. We used direct photon data also from PHENIX [6, 11]. Results are shown in Figs. 1 and 2, while the model parameters are detailed in Table. 1.

The EoS result from the photon fit is $\kappa = 7.9 \pm 0.7$, or alternatively, using $\kappa = 1/c_s^2$:

$$c_s = 0.36 \pm 0.02_{\text{stat}} \pm 0.04_{\text{sys}} \quad (3.1)$$

which is in agreement with lattice QCD calculations [12] and measured hadronic data [13, 14]. This represents an average EoS as it may vary with temperature. There may be solutions with a

Dataset		N_1 and HBT	elliptic flow	N_1
		0-30% cent. hadrons	0-30% cent. hadrons	0-92% cent. photons
Central FO temperature	T_0 [MeV]	199 ± 3	204 ± 7	204 MeV (fixed)
Eccentricity	ε	0.80 ± 0.02	0.34 ± 0.03	0.34 (fixed)
Transverse expansion	u_t^2/b	-0.84 ± 0.08	-0.34 ± 0.01	-0.34 (fixed)
FO proper-time	τ_0 [fm/c]	7.7 ± 0.1	-	7.7 (fixed)
Longitudinal expansion	Z_0^2/b	-1.6 ± 0.3	-	-1.6 (fixed)
Equation of State	κ	-	-	7.9 ± 0.7
Initial time	t_i [fm/c]	-	-	0 – 0.7 fm/c
Fit quality				
Degrees of freedom	NDF	41	34	3
Chisquare	χ^2	24	66	7
Confidence level		98%	0.1%	7.2%

Table 1: Parameters of the solution, describing the expanding sQGP. The first five were determined from hadronic fits [3], the remaining from direct photon data [4]. See details in the text and in these references.




Figure 1: Fits to invariant momentum distribution of pions [8] (top left), HBT radii [10] (top right) and elliptic flow [9] (bottom). See the obtained parameters in Table 1. In the middle plot the lower curve is the fit to R_{out} and R_{side} .




Figure 2: Fit to direct photon invariant transverse momentum data [6] (top left), comparison to elliptic flow data [11] (top right) and direct photon HBT predictions (bottom). See the model parameters in Table 1.

$\kappa(T)$ function [15], but for the sake of simplicity we assumed here an average, fixed κ . As detailed in Ref. [4], we determined an “interval of acceptability” for t_i . The maximum value for t_i within 95% probability is 0.7 fm/c. This can then be used to determine a lower bound for the initial temperature, using the eq. 1.2. Thus the initial temperature of the fireball (in its center) is:

$$T_i > 507 \pm 12_{stat} \pm 90_{syst} \text{ MeV} \quad (3.2)$$

at 0.7 fm/c. This is in accordance with other hydro models as those values are in the 300 – 600 MeV interval [6]. Note that a systematic uncertainty was determined by using a prefactor of $(T/T_0)^N$, with $N = 0, 1, 2, 3$. This factor arises if the photon creation can be described by a microscopic process, as detailed in Ref. [4]. This gives a systematic uncertainty to our parameters. In detail, we multiplied the source function with each of these prefactors (with $N = 0, 1, 2, 3$ being separate cases) and calculated the spectra with this modified source. Then we compared the spectrum to data and determined c_s and T_i . The difference between the $N = 0$ and 3 cases was taken as a systematic error. This causes only a minor change in the resulting spectrum, as it is dominated by the exponential factors in it. However, the equation of state parameter κ changes from 7.9 to 6.5 as we increase the exponent in the prefactor, and thus the speed of sound changes by 0.04, and the initial temperature by 90 MeV.

A measurement of direct photon elliptic flow was also performed recently at PHENIX [11]. Using the previously determined fit parameters we can calculate the elliptic flow of direct photons in Au+Au collisions at RHIC. Due to the low number of points in the desired range, a fit could not be performed here, but we used the average value ε in case of the two fits of Ref. [3]. The resulting curve, where the value $\varepsilon = 0.59$ was used, is shown on Fig. 2.

In case of hadronic HBT, correlation radii in the side and out directions are almost equal, as for the hadronic transition is of cross-over type (i.e. the transition time is short), see details in Ref. [3]. However, in case of photons, the creation spans the whole evolution of the fireball, thus R_{out} will be significantly larger than R_{side} . Indeed this was observed in our model, as shown on Fig. 2.

To summarize, we find that thermal radiation is consistent with direct photon data, and our result on the equation of state is $c_s = 0.36 \pm 0.02_{\text{stat}} \pm 0.04_{\text{syst}}$. We set a lower bound on the initial temperature of the sQGP to $507 \pm 12_{\text{stat}} \pm 90_{\text{syst}}$ MeV at 0.7 fm/c. We also find that the thermal photon elliptic flow from this mode is not incompatible with measurements. We also predicted photon HBT radii from the model, and discovered a significantly larger R_{out} than R_{side} for photons, which is in contrast to the behaviour observed for hadrons.

Acknowledgments

The author would like to thank the invitation to the WPCF 2011 conference, and the kind hospitality of the organizers, in particular T. Hirano. The author also would like to thank T. Csörgő and S. Pratt for valuable discussions and important insights on the topic of this paper. M. Csanád gratefully acknowledges the support of the Bolyai Scholarship of the Hungarian Academy of Sciences and the OTKA grant NK 101438.

References

- [1] PHENIX, K. Adcox *et al.*, Nucl. Phys. **A757**, 184 (2005), nucl-ex/0410003.
- [2] T. Csörgő, L. P. Csernai, Y. Hama, and T. Kodama, Heavy Ion Phys. **A21**, 73 (2004), nucl-th/0306004.
- [3] M. Csanád and M. Vargyas, Eur. Phys. J. **A44**, 473 (2010), 0909.4842.
- [4] M. Csanád and I. Májer, (2011), 1101.1279.
- [5] M. Csanád, Acta Phys. Polon. **B40**, 1193 (2009), 0903.1278.
- [6] PHENIX, A. Adare *et al.*, Phys. Rev. Lett. **104**, 132301 (2010), 0804.4168.
- [7] F. Cooper and G. Frye, Phys. Rev. **D10**, 186 (1974).
- [8] PHENIX, S. S. Adler *et al.*, Phys. Rev. **C69**, 034909 (2004), nucl-ex/0307022.
- [9] PHENIX, S. S. Adler *et al.*, Phys. Rev. Lett. **93**, 152302 (2004), nucl-ex/0401003.
- [10] PHENIX, S. S. Adler *et al.*, Phys. Rev. Lett. **91**, 182301 (2003), nucl-ex/0305013.
- [11] PHENIX Collaboration, A. Adare *et al.*, (2011), 1105.4126.
- [12] S. Borsányi *et al.*, JHEP **11**, 077 (2010), 1007.2580.
- [13] PHENIX, A. Adare *et al.*, Phys. Rev. Lett. **98**, 162301 (2007), nucl-ex/0608033.
- [14] R. A. Lacey and A. Taranenko, PoS **CFRNC2006**, 021 (2006), nucl-ex/0610029.
- [15] M. Csanád, Talk at the Zimányi School 2010.