

Gluon saturation and final states

Krzysztof Kutak*

*Instytut Fizyki Jądrowej im. H. Niewodniczańskiego
Radzikowskiego 152, 31-342 Krakow, Poland
E-mail: krzysztof.kutak@ifj.edu.pl*

In order to study such effects like parton saturation in final states at the LHC one of the approaches is to combine physics of the BK and the CCFM evolution equations. We report on obtained resummed form of the BK equation and nonlinear extension of the CCFM equation.

*Xth Quark Confinement and the Hadron Spectrum,
October 8-12, 2012
TUM Campus Garching, Munich, Germany*

*Speaker.

1. Introduction

Quantum Chromodynamics (QCD) is a theory which is used to set up the initial conditions for the collisions at the LHC as well as to calculate properties of hadronic observables. In order to apply perturbative QCD to scattering process one needs to decompose the cross section into long distance part, called parton density, and a short distance part, called matrix element. In particular here we will focus on high energy factorization [1, 2]. The evolution equations of high energy factorization resum logarithms of energy accompanied by a strong coupling constant, i.e. terms proportional to $\alpha_s^n \ln^m s/s_0$. This framework applies if the total energy of a scattering process is much bigger than any other hard scale involved in a scattering process.

With the LHC one entered into a region of phase space where both the energy and momentum transfers are high and partons form dense system allowing in principle for parton saturation [3–6]. Recently a framework has been provided in [7–11] where both dense systems and hard processes at high energies can be studied.

2. Exclusive form of the Balitsky-Kovchegov equation

At the leading order in $\ln 1/x$ the Balitsky-Kovchegov [12, 13] equation for the WW gluon density in the momentum space is written as an integral equation reads [7, 9]:

$$\begin{aligned} \Phi(x, k^2) = & \Phi_0(x, k^2) \\ & + \bar{\alpha}_s \int_{x/x_0}^1 \frac{dz}{z} \int_0^\infty \frac{dl^2}{l^2} \left[\frac{l^2 \Phi(x/z, l^2) - k^2 \Phi(x/z, k^2)}{|k^2 - l^2|} + \frac{k^2 \Phi(x/z, k)}{\sqrt{(4l^4 + k^4)}} \right] \\ & - \frac{\bar{\alpha}_s}{\pi R^2} \int_x^1 \frac{dz}{z} \Phi^2(x/z, k) \end{aligned} \quad (2.1)$$

where the lengths of transverse vectors lying in transversal plane to the collision axis are $k \equiv |\mathbf{k}|$, $l \equiv |\mathbf{l}|$ (\mathbf{k} is a vector sum of transversal momenta of emitted gluons during evolution), $z = x/x'$ (see Fig. (1)), $\bar{\alpha}_s = N_c \alpha_s / \pi$. The strength of the nonlinear term is controlled by the targets radius R . The linear term in eq. (2.1) can be linked to the process of creation of gluons while the nonlinear term can be linked to fusion of gluons and therefore introduces gluon saturation effects.

The unintegrated gluon density obeying the high energy factorization theorem [1] is obtained from [14, 15]:

$$\mathcal{F}_{BK}(x, k^2) = \frac{N_c}{\alpha_s \pi^2} k^2 \nabla_k^2 \Phi(x, k^2) \quad (2.2)$$

where the angle independent Laplace operator is given by $\nabla_k^2 = 4 \frac{\partial}{\partial k^2} k^2 \frac{\partial}{\partial k^2}$.

As explained in [7] this equation can be rewritten in a resummed form:

$$\begin{aligned} \Phi(x, k^2) = & \check{\Phi}^0(x, k^2) + \bar{\alpha}_s \int_{x/x_0}^1 dz \int \frac{d^2 \mathbf{q}}{\pi q^2} \theta(q^2 - \mu^2) \frac{\Delta_R(z, k, \mu)}{z} \left[\Phi\left(\frac{x}{z}, |\mathbf{k} + \mathbf{q}|^2\right) \right. \\ & \left. - \frac{1}{\pi R^2} q^2 \delta(q^2 - k^2) \Phi^2\left(\frac{x}{z}, q^2\right) \right]. \end{aligned} \quad (2.3)$$




Figure 1: Plot explaining meaning of variables in BK and CCFM evolution equations.

where $\mathbf{q} = \mathbf{1} - \mathbf{k}$ and $\Delta_R(z, k, \mu) \equiv \exp\left(-\bar{\alpha}_s \ln \frac{1}{z} \ln \frac{k^2}{\mu^2}\right)$ is a Regge form factor.

Eq. (2.3) is a new form of the BK equation in which the resummed terms in a form of Regge form factor are the same for the linear and nonlinear part. This form will serve as a guiding equation to generalize the CCFM [16–18] equation to include nonlinear effects which allow for recombination of partons with constraint on angle of emission. As has been shown in [9] an analogous equation to (2.3) can be written for the high energy factorizable gluon density and it reads:

$$\mathcal{F}(x, k^2) = \tilde{\mathcal{F}}_0(x, k^2) + \bar{\alpha}_s \int_{x/x_0}^1 \frac{dz}{z} \Delta_R(z, k, \mu) \left\{ \int \frac{d^2 q}{\pi q^2} \theta(q^2 - \mu^2) \mathcal{F}\left(\frac{x}{z}, |k+q|^2\right) - \frac{\pi \alpha_s^2}{4N_c R^2} k^2 \nabla_k^2 \left[\int_{k^2}^{\infty} \frac{dl^2}{l^2} \ln \frac{l^2}{k^2} \mathcal{F}(x, l^2) \right]^2 \right\} \quad (2.4)$$

3. CCFM evolution equation

The CCFM equation sums up gluonic emissions with the condition of strong ordering in emission angle which allow for smooth interpolation between BFKL limit when $z \rightarrow 0$ and the DGLAP limit (in gluonic channel) when $z \rightarrow 1$. The BFKL or the BK are generally applied for the calculation of elastic scattering and total cross-sections while due to the dependence on the hard scale (which is expressed as the maximal angle) the CCFM equation allows for studies of exclusive observables. It is the following

$$\mathcal{A}(x, k^2, p) = \mathcal{A}_0(x, k^2, p) + \bar{\alpha}_s \int_{x/x_0}^1 dz \int \frac{d^2 \bar{q}}{\pi \bar{q}^2} \theta(p - z\bar{q}) \Delta_s(p, z\bar{q}) \left(\frac{\Delta_{ns}(z, k, q)}{z} + \frac{1}{1-z} \right) \mathcal{A}\left(\frac{x}{z}, k'^2, \bar{q}\right). \quad (3.1)$$

The gluon density obtained from the CCFM equation, usually denoted $\mathcal{A}(x, k^2, p)$, on a level of linear equation has interpretation of the gluon density describing parton with longitudinal momentum fraction x , and transverse momentum (squared) k^2 which is probed by a hard system at scale

p .

The momentum vector associated with i -th emitted gluon is

$$q_i = \alpha_i p_P + \beta_i p_e + q_{ti} \quad (3.2)$$

from which the rapidity and angle of emitted gluon with respect to incoming parent proton (beam direction) can be obtained as

$$\eta_i = \frac{1}{2} \ln(\xi_i) \equiv \frac{1}{2} \ln\left(\frac{\beta_i}{\alpha_i}\right) = \ln\left(\frac{|\mathbf{q}_i|}{\sqrt{s} \alpha_i}\right), \quad \tan \frac{\theta_i}{2} = \frac{|\mathbf{q}_i|}{\sqrt{s} \alpha_i}. \quad (3.3)$$

The variable p in (3.1) is defined via $\bar{\xi} = p^2/(x^2 s)$ where $\frac{1}{2} \ln(\bar{\xi})$ is a maximal rapidity which is determined by the kinematics of hard scattering, \sqrt{s} is the total energy of the collision. For example, for the electron proton scattering $s = (p_P + p_e)^2$ and $k' = |\mathbf{k} + (1-z)\bar{\mathbf{q}}|$. Using the variables ξ the angular ordering can be conveniently expressed as: $\bar{\xi} > \xi_i > \xi_{i-1} > \dots > \xi_1 > \xi_0$, where $\xi_0 \equiv \mu$ with μ being infrared cut off. The momentum $\bar{\mathbf{q}}$ is the transverse rescaled momentum of the real gluon, and is related to \mathbf{q} by $\bar{\mathbf{q}} = \mathbf{q}/(1-z)$ and $\bar{q} \equiv |\bar{\mathbf{q}}|$.

The Sudakov form factor which screens the $1-z$ singularity is given by

$$\Delta_s(p, z\bar{q}) = \exp\left(-\bar{\alpha}_s \int_{(z\bar{q})^2}^{p^2} \frac{dq^2}{q^2} \int_0^{1-\mu/q} \frac{dz}{1-z}\right). \quad (3.4)$$

The form factor Δ_{ns} screens the $1/z$ singularity, in a similar form as the Regge form factor but also accounts for angular ordering:

$$\Delta_{ns}(z, k, q) = \exp\left(-\bar{\alpha}_s \int_z^1 \frac{dz'}{z'} \int_{z'^2 q^2}^{k^2} \frac{dq'^2}{q'^2}\right) = \exp\left(-\alpha_s \ln \frac{1}{z} \ln \frac{k^2}{zq^2}\right). \quad (3.5)$$

where for the lowest value of zq^2 we use a cut off μ . In the literature [19] different forms of the non-Sudakov form factor were considered, where the differences came from the approximations taken, the one we are using follows results obtained in [17]. For studies of effects of various forms of non-Sudakov form factors we refer the reader to [23].

3.1 Nonlinear extension of the CCFM equation - the KGBJS equation

As it has already been stated the motivation to extend the CCFM to account for nonlinearity is to be able to study the impact of saturation of partons on exclusive observables. There are indications that such effects might be significant in for instance production of charged particles at HERA [20, 21] or in forward production of di-jets [3, 5].

The simplest nonlinear extension of CCFM has been recently proposed in [7] and its extension changes the interpretation of the quantity for which the equation is written. It is not longer high energy factorizable gluon density but should be interpreted as the dipole amplitude in momentum space or Weizsacker-Williams gluon density Φ , denoted from now on by \mathcal{E} , which besides x and k^2 depends also on a hard scale p . The peculiar structure of the nonlinear term of the equation written below is motivated by the following requirements:

- the second argument of the \mathcal{E} should be k^2 as motivated by the analogy to BK

- the third argument should reflect locally the angular ordering

The KGBJS equation reads:

$$\mathcal{E}(x, k^2, p) = \mathcal{E}_0(x, k^2, p) + \bar{\alpha}_s \int_x^1 dz \int \frac{d^2 \bar{q}}{\pi \bar{q}^2} \theta(p - z\bar{q}) \Delta_s(p, z\bar{q}) \left(\frac{\Delta_{ns}(z, k, q)}{z} + \frac{1}{1-z} \right) \times \quad (3.6)$$

$$\left[\mathcal{E} \left(\frac{x}{z}, k'^2, \bar{q} \right) - \frac{1}{\pi R^2} \bar{q}^2 \delta(\bar{q}^2 - k^2) \mathcal{E}^2 \left(\frac{x}{z}, \bar{q}^2, \bar{q} \right) \right].$$

For recent numerical study and incorporation of higher order effects to equation (3.6) we refer the Reader to [24]. Similarly as in case of the BK equation in order to obtain high energy factorizable unintegrated gluon density one applies relation (2.2). The nonlinear term in (3.6), apart from allowing for recombination of gluons might be understood as a way to introduce the decoherence into the emission pattern of gluons. This is because the gluon density is build up due to coherent gluon emission and since the nonlinear term comes with the negative sign it slows down the growth of gluon density and therefore it introduces the decoherence. We expect the nonlinear term to be of main importance at low x similarly as in case of the BK equation. In this limit it will be of special interest to check whether in our formulation of the nonlinear extension of the CCFM equation one obtains an effect of saturation of the saturation scale as observed in [23] (for other formulations to include unitarity corrections in the CCFM evolution equation we refer the reader to [21, 25]). This effect is of great importance since it has a consequences for example for imposing a bound on amount of produced entropy from saturated part of gluon density as observed in [26].

4. Conclusions and outlook

We reported on recently obtained new form of the BK equation written in a resummed form and on extension of the CCFM equation to account for nonlinearity. The obtained extension of the CCFM equation will be useful for studies of impact of saturation of gluons on exclusive observables.

5. Acknowledgements

Part of the research presented has been obtained in collaboration with Krzysztof Golec-Biernat, Stanislaw Jadach and Maciej Skrzypek. This research has been supported by grant LIDER/02/35/L-2/10/NCBiR/2011

6. Bibliography

References

- [1] S. Catani, M. Ciafaloni, F. Hautmann, Nucl. Phys. **B366** (1991) 135-188.
- [2] L. V. Gribov, E. M. Levin and M. G. Ryskin, Phys. Rept. **100** (1983) 1.
- [3] J. L. Albacete, C. Marquet, Phys. Rev. Lett. **105** (2010) 162301.

- [4] A. Dumitru, K. Dusling, F. Gelis *et al.*, Phys. Lett. **B697** (2011) 21-25.
- [5] K. Kutak and S. Sapeta, arXiv:1205.5035 [hep-ph].
- [6] K. Dusling and R. Venugopalan, arXiv:1210.3890 [hep-ph].
- [7] K. Kutak, K. Golec-Biernat, S. Jadach and M. Skrzypek, JHEP **1202** (2012) 117.
- [8] K. Kutak, arXiv:1206.1223 [hep-ph].
- [9] K. Kutak, arXiv:1206.5757 [hep-ph].
- [10] A. van Hameren, P. Kotko and K. Kutak, arXiv:1207.3332 [hep-ph].
- [11] A. van Hameren, P. Kotko and K. Kutak, arXiv:1211.0961 [hep-ph].
- [12] I. Balitsky, Nucl. Phys. B **463** (1996) 99.
- [13] Y. V. Kovchegov, Phys. Rev. D **60** (1999) 034008.
- [14] M. Braun, Eur. Phys. J. C **16**, 337 (2000).
- [15] K. Kutak and J. Kwiecinski, Eur. Phys. J. C **29** (2003) 521.
- [16] M. Ciafaloni, Nucl. Phys. B **296**, 49 (1988).
- [17] S. Catani, F. Fiorani and G. Marchesini, Nucl. Phys. B **336**, 18 (1990).
- [18] S. Catani, F. Fiorani and G. Marchesini, Phys. Lett. B **234**, 339 (1990).
- [19] J. Kwiecinski, A. D. Martin and P. J. Sutton, Phys. Rev. D **52** (1995) 1445.
- [20] C. Adloff *et al.* [H1 Collaboration], Nucl. Phys. **B485** (1997) 3-24.
- [21] K. Kutak, H. Jung, Acta Phys. Polon. **B40** (2009) 2063-2070.
- [22] E. Avsar and E. Iancu, Nucl. Phys. A **829** (2009) 31.
- [23] E. Avsar and A. M. Stasto, JHEP **1006** (2010) 112.
- [24] M. Deak, arXiv:1209.6092 [hep-ph].
- [25] E. Avsar and E. Iancu, Nucl. Phys. A **829** (2009) 31 [arXiv:0906.2683 [hep-ph]].
- [26] K. Kutak, Phys. Lett. B **705** (2011) 217.