

$\Upsilon(5S)$ spectroscopy at Belle

A.Bondar*, A.Garmash and P.Krokovny (on behalf of the Belle Collaboration)

We report the first observation of the three-body $\Upsilon(5S) \rightarrow \Upsilon(1,2S)\pi^0\pi^0$ decays. Strong evidence for the $Z_b^0(10610)$ with 4.9σ significance is found in a Dalitz plot analysis of the $\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^0\pi^0$ decays. First results on the analysis of the three-body $\Upsilon(10860) \rightarrow [B\bar{B}^* + \text{c.c.}]^\pm\pi^\mp$ and $\Upsilon(10860) \rightarrow [B^*\bar{B}^*]^\pm\pi^\mp$ including first observation of $Z_b^\pm(10610) \rightarrow [B\bar{B}^* + \text{c.c.}]^\pm$ and $Z_b^\pm(10650) \rightarrow [B^*\bar{B}^*]^\pm$ are also reported. The results are obtained with a 121.4fb^{-1} data sample collected with the Belle detector at the $\Upsilon(5S)$ resonance at the KEKB asymmetric-energy e^+e^- collider.

ICHEP 2012

*Speaker.

1. Introduction

Two new charged bottomonium-like resonances, $Z_b(10610)$ and $Z_b(10650)$, have recently been observed by the Belle Collaboration in decays of $\Upsilon(10860)$ to five different final states: $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ and $h_b(mP)\pi^+\pi^-$, $m = 1, 2$ [1, 2]. The analysis of the quark composition of the initial and final states allows to assert that these hadronic objects are the first unambiguous examples of states of an exotic nature: Z_b should be comprised of (at least) four quarks. Several models have been proposed to describe the internal structure of these states. One suggests [3] that $Z_b(10610)$ and $Z_b(10650)$ states might be a loosely bound $B\bar{B}^*$ and $B^*\bar{B}^*$ systems, respectively. The proximity of the $Z_b(10610)$ and $Z_b(10650)$ masses to those of the sum of the B and B^* mesons and the sum of the two B^* mesons, respectively, supports this hypothesis. In this case, it would be natural to expect that the $Z_b(10610)$ and $Z_b(10650)$ states decay respectively to $B\bar{B}^*$ and $B^*\bar{B}^*$ final states with substantial rates. In this analysis we use 121.4 fb^{-1} of data accumulated by the Belle detector at a center-of-mass (c.m.) energy near the $\Upsilon(10860)$ to study three-body $\Upsilon(10860) \rightarrow [B^{(*)}\bar{B}^{(*)}]^\pm \pi^\mp$ decays and to search for $\Upsilon(10860) \rightarrow Z_b^\pm \pi^\mp \rightarrow [B^{(*)}\bar{B}^{(*)}]^\pm \pi^\mp$ decays. We also search for a neutral partner of Z_b states in the resonant substructure of the $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^0\pi^0$ decays.

2. $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^0\pi^0$ decays

We reconstruct $\Upsilon(nS)$ candidates from pairs of leptons (e^+e^- and $\mu^+\mu^-$). An additional decay channel is used for the $\Upsilon(2S)$: $\Upsilon(2S) \rightarrow \Upsilon(1S)[l^+l^-]\pi^+\pi^-$. Muon and electron candidates are required to be positively identified. Candidate π^0 mesons are selected from pairs of photons with an invariant mass within $15 \text{ MeV}/c^2$ of the nominal π^0 mass. $\Upsilon(5S) \rightarrow \Upsilon(nS)[l^+l^-]\pi^0\pi^0$ candidates are identified via the missing mass recoiling against the $\pi^0\pi^0$ system, $M_{\text{miss}}(\pi^0\pi^0)$. More details can be found in Ref. [4]. Figures 1 (a) and 1 (b) show the $M_{\text{miss}}(\pi^0\pi^0)$ distributions for $\Upsilon(5S) \rightarrow \Upsilon(nS)[l^+l^-]\pi^0\pi^0$ candidates. We fit these distributions to extract the $\Upsilon(nS)$ signal yield. Results of fits are summarized in Table 1, where signal yield, MC efficiency, measured branching fraction (only the statistical uncertainty is quoted), number of selected events and signal purity. The reconstruction efficiency is obtained using signal MC events with uniform distribution of final state particles. Branching fraction is calculated as $\mathcal{B} = \frac{N_{\text{sig}}}{\varepsilon \mathcal{L} \sigma(e^+e^- \rightarrow \Upsilon(5S))}$, where N_{sig} is number of signal events, ε is reconstruction efficiency, \mathcal{L} is integrated luminosity. Weighted averages are found to be $\mathcal{B}(\Upsilon(5S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (2.25 \pm 0.11 \pm 0.20) \times 10^{-3}$ and $\mathcal{B}(\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^0\pi^0) = (3.66 \pm 0.22 \pm 0.48) \times 10^{-3}$.

The amplitude analysis of the three-body $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^0\pi^0$ decays utilizes an unbinned maximum likelihood fit. We parameterize the three-body decay amplitude as a sum of quasi-two-body amplitudes: $M(s_1, s_2) = A_{Z_1} + A_{Z_2} + A_{f_0} + A_{f_2} + A_{\text{nr}}$, where A_{Z_1} and A_{Z_2} are amplitudes for contributions from the $Z_b^0(10610)$ and $Z_b^0(10650)$, respectively; the amplitudes A_{f_0} , A_{f_2} and A_{nr} are the contributions from the $\pi^0\pi^0$ system in an $f_0(980)$, $f_2(1275)$ and a non-resonant state, respectively. The masses and widths of Z_b resonances are fixed at the values obtained from the $\Upsilon(nS)\pi^+\pi^-$ analysis: $M(Z_1) = 10607.2 \text{ MeV}/c^2$, $\Gamma(Z_1) = 18.4 \text{ MeV}/c$, $M(Z_2) = 10652.2 \text{ MeV}/c^2$, $\Gamma(Z_2) = 11.5 \text{ MeV}/c$ [1]. We use a Flatté function for the $f_0(980)$ and a Breit-Wigner function for the $f_2(1275)$. The non-resonant amplitude A_{nr} is parameterized as $A_{\text{nr}} = A_{\text{nr}}^1 e^{i\phi_{\text{nr}}^1} + A_{\text{nr}}^2 e^{i\phi_{\text{nr}}^2} s_3$, where




Figure 1: The $\pi^0\pi^0$ missing mass distribution for $\Upsilon(nS)\pi^0\pi^0$, (a) $\Upsilon(nS) \rightarrow \mu^+\mu^-$ and (b) $\Upsilon(nS) \rightarrow e^+e^-$ candidates. The $M(\Upsilon(1S)\pi^+\pi^-)$ distribution for $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ candidates is shown in (c). Histograms represent the data. The solid curves show the fit result while the dashed curves correspond to the background contributions.

$A_{nr}^1, A_{nr}^2, \phi_{nr}^1$ and ϕ_{nr}^2 are free parameters in the fit. As the fit is sensitive only to the relative amplitudes and phases between decay modes, we fix $A_{nr}^1 = 10.0$ and $\phi_{nr}^1 = 0.0$.

The logarithmic likelihood function is defined as $\mathcal{L} = -2 \sum \log\{\varepsilon(s_1, s_2)(f_{sig}S(s_1, s_2) + (1 - f_{sig})B(s_1, s_2))\}$, where $S(s_1, s_2)$ denotes $|M(s_1, s_2)|^2$ convoluted with the detector resolution function ($6.0\text{MeV}/c^2$ for $\Upsilon(nS)\pi^0$ combinations), $\varepsilon(s_1, s_2)$ describes variation of the reconstruction efficiency over the Dalitz plot and f_{sig} is the fraction of signal events in the data sample. The fraction f_{sig} is determined separately for each $\Upsilon(nS)$ decay mode (see Table 1). The function $B(s_1, s_2)$ describes the distribution of background events over the phase space. Both products $\varepsilon(s_1, s_2) \cdot S(s_1, s_2)$ and $\varepsilon(s_1, s_2) \cdot B(s_1, s_2)$ are normalized to unity.

Results of the fits are shown in Fig. 2 as one-dimensional projections that look similar to the corresponding distributions for the $\Upsilon(nS)\pi^+\pi^-$ decays [1]. A Z_b^0 signal is most clearly seen in $M(\Upsilon\pi^0)_{max}$. The values and errors of amplitudes and phases obtained from the fit are presented in [4]. The statistical significance of the $Z_b^0(10610)$ signal in the $\Upsilon(2S)\pi^0\pi^0$ sample is 5.3σ . The signal for the $Z_b^0(10610)$ is not significant in the fit to the $\Upsilon(1S)\pi^0\pi^0$ events due to the smaller relative branching fraction. The signal of the $Z_b^0(10650)$ is not significant in either $\Upsilon(1)\pi^0\pi^0$ or $\Upsilon(2S)\pi^0\pi^0$ datasets.

We study possible uncertainties due to parameterization of the background PDF, variation of signal efficiency over the Dalitz plot and detector resolution function. The model uncertainty is estimated using various description of S -wave contribution. The significance of the $Z_b^0(10610)$ signal exceeds 4.9σ in all cases. We use this value as the final value for the $Z_b^0(10610)$ significance.

Table 1: Signal yield, MC efficiency, measured branching fraction, number of selected events and purity.

Final state	Signal yield	$\varepsilon, \%$	$\mathcal{B}, 10^{-3}$	Events	Purity
$\Upsilon(1S) \rightarrow \mu^+\mu^-$	261 ± 15	11.2	2.28 ± 0.13	247	0.95
$\Upsilon(1S) \rightarrow e^+e^-$	123 ± 13	5.61	2.15 ± 0.23	140	0.78
$\Upsilon(2S) \rightarrow \mu^+\mu^-$	241 ± 18	8.04	3.77 ± 0.28	253	0.87
$\Upsilon(2S) \rightarrow e^+e^-$	108 ± 13	3.58	3.84 ± 0.46	151	0.66
$\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$	24 ± 5	2.27	2.85 ± 0.60	28	0.86




Figure 2: Comparison of the fit results (open histograms) with experimental data (points with error bars) for the (a,b) $\Upsilon(1S)\pi^0\pi^0$ and (c) $\Upsilon(2S)\pi^0\pi^0$ events in the signal region. Red and blue open histograms show the fit with and without Z_b^0 's, respectively. Hatched histograms show the background components.

3. $\Upsilon(10860) \rightarrow B^{(*)}\bar{B}^*\pi$ Decays

B decays are reconstructed in the following channels: $B^+ \rightarrow J/\psi K^+$, $B^+ \rightarrow \bar{D}^0\pi^+$, $B^0 \rightarrow J/\psi K^{*0}$, $B^0 \rightarrow D^-\pi^+$, $B^0 \rightarrow D^{*-}\pi^+$. We identify B candidates by their invariant mass $M(B)$ and momentum $P(B)$ in the c.m. We require $M(B)$ to be within 30 to 40 MeV/c^2 (depending on the B decay mode) of the nominal B mass. Reconstructed B^+ or B^0 candidates are then combined with a π^- candidate and a recoil mass to the $B\pi$ combination, $M_r(B\pi)$, is calculated as $M_r(B\pi) = \sqrt{E_{\text{cms}}^2 - P_{B\pi}^2}$, where E_{cms} is the c.m. energy and $P_{B\pi}$ is the measured three-momentum of the $B\pi$ combination. More details can be found in Ref. [5]. The $M_r(B\pi) + M(B) - M_B$ distribution for the data is shown in Fig. 3(a), where clear peaks are visible in the $BB^*\pi$ and $B^*B^*\pi$ signal regions.

The fit to the $M_r(B\pi) + M(B) - M_B$ distribution for signal events shown in Fig. 3(a) yields $N_{BB\pi} = 1 \pm 14$, $N_{BB^*\pi} = 184 \pm 19$ and $N_{B^*B^*\pi} = 82 \pm 11$ signal events. The statistical significance of the observed $BB^*\pi$ and $B^*B^*\pi$ signal is 9.3σ and 5.7σ , respectively. For the subsequent analysis of the internal structures of the three-body decays, we require $|(M_r(B) + M(B) - M_B) - M_{B^*}| < 0.015 \text{ GeV}/c^2$ to select $\Upsilon(10860) \rightarrow BB^*\pi$ events and $|(M_r(B) + M(B) - M_B) - (M_{B^*} + E_\gamma)| < 0.015 \text{ GeV}/c^2$, where $E_\gamma = 0.049 \text{ GeV}$, to select $\Upsilon(10860) \rightarrow B^*B^*\pi$ events. For selected $B^{(*)}\bar{B}^{(*)}\pi$ candidate events, we calculate the mass recoiling against the charged pion: $M_r(\pi) = \sqrt{E_{\text{cms}}^2 - P_\pi^2}$, where $P_{B\pi}$ is the measured three-momentum of the charged pion.

The $M_r(\pi)$ distributions for right-sign $B\pi$ combinations in the $BB^*\pi$ and $B^*B^*\pi$ signal regions are shown in Fig. 3. Excesses of signal events over the expected background levels at lower mass edges of the $M_r(\pi)$ spectra are clearly visible for both final states. The distribution of signal $\Upsilon(10860) \rightarrow BB^*\pi$ events is parameterized with the following model $S_{BB^*\pi}(m) = (A_{Z_b(10610)} + A_{NR}) \times E_{BB^*\pi}(m)$, where A_{NR} is the non-resonant amplitude parameterized as a complex constant and the $Z_b(10610)$ amplitude is a Breit-Wigner function. As a variation of this nominal model, we also add a second Breit-Wigner amplitude to account for possible $Z_b(10650) \rightarrow BB^*\pi$ decay. We also fit the data with only the $Z_b(10610)$ channel included in the decay amplitude. The results of these fits are shown in Fig. 3(b). Two models give about equally good description of the data: nominal model and a model with additional non-resonant amplitude. However, we select the former one as our nominal model since adding a non-resonant amplitude does not improve the fit quality that much. The worst fit to the data is provided by a model with just a non-resonant amplitude.




Figure 3: (a) $M_r(B\pi)$ distributions for selected B candidates in data. Hatched histogram shows distribution for events in the $M(B)$ sidebands. (b) $M_r(\pi)$ distribution for right-sign $B\pi$ combinations for $\Upsilon(10860) \rightarrow BB^*\pi$ and (c) $\Upsilon(10860) \rightarrow B^*B^*\pi$ candidate events. Points with error bars are data, the solid line is the result of the fit with the nominal model, the dashed line - fit to pure non-resonant amplitude, the dotted line - fit to a single Z_b state plus a non-resonant amplitude, and the dash-dotted - two Z_b states and a non-resonant amplitude. The hatched histogram represents background component.

From this analysis, we find that the significance of the $Z_b(10610) \rightarrow BB^*$ signal is exceeding the 8σ level.

As the nominal model for the $\Upsilon(10860) \rightarrow B^*B^*\pi$ decay, we use the following parameterization: $S_{B^*B^*\pi}(m) = (A_{Z_b(10650)} + A_{NR})E_{B^*B^*\pi}(m)$. We also fit the data without a non-resonant component and with a non-resonant amplitude alone. Results of the fits are shown in Fig. 3(c); numerical values are given in [5]. The best description of the $B^*B^*\pi$ data is achieved in a model with only the $Z_b(10650)$ amplitude included. The addition of a non-resonant amplitude does not provide any significant improvement of the fit quality. The fit with a non-resonant amplitude alone gives a much worse likelihood value. From this analysis, we determine the significance of the $Z_b(10650) \rightarrow B^*B^*$ signal to be 6.8σ .

In all fits discussed above, the masses and widths of the Z_b states were fixed at the values obtained from the analysis of the $\Upsilon(nS)\pi^+\pi^-$ and $h_b(mP)\pi^+\pi^-$ final states: $M[Z_b(10610)] = 10607.2 \pm 2.0 \text{ MeV}/c^2$, $\Gamma[Z_b(10610)] = 18.4 \pm 2.4 \text{ MeV}$ and $M[Z_b(10650)] = 10652.2 \pm 1.5 \text{ MeV}/c^2$, $\Gamma[Z_b(10650)] = 11.5 \pm 2.2 \text{ MeV}$. If allowed to float, the fit returns $10597 \pm 9 \text{ MeV}$ for $Z_b(10610)$ mass in the fit to $BB^*\pi$ events and $10649 \pm 12 \text{ MeV}$ for $Z_b(10650)$ mass in the fit to $B^*B^*\pi$ events. Large errors here reflect a strong negative correlation between resonance mass and its amplitude.

4. Conclusion

We report the first observation of three-body $\Upsilon(5S) \rightarrow \Upsilon(1,2S)\pi^0\pi^0$ decays. The measured branching fractions are $\mathcal{B}(\Upsilon(5S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (2.25 \pm 0.11 \pm 0.20) \times 10^{-3}$ and $\mathcal{B}(\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^0\pi^0) = (3.66 \pm 0.22 \pm 0.48) \times 10^{-3}$. Evidence for a $Z_b^0(10610) \rightarrow \Upsilon(2S)\pi^0$ decay has been obtained from the amplitude analysis of the $\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^0\pi^0$ decay. The statistical significance of the $Z_b^0(10610)$ signal is 4.9σ including model and systematic uncertainties. Its measured mass, $M(Z_b^0(10610)) = 10609_{-6}^{+8} \pm 6 \text{ MeV}/c^2$, is consistent with that measured in the analysis of $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^+\pi^-$ decays. The $Z_b^0(10650)$ signal is not significant in either $\Upsilon(1S)\pi^0\pi^0$ or $\Upsilon(2S)\pi^0\pi^0$ final decays.

Table 2: List of branching fractions for the $Z_b^+(10610)$ and $Z_b^+(10650)$ decays.

Channel	Fraction, %	
	$Z_b(10610)$	$Z_b(10650)$
$\Upsilon(1S)\pi^+$	0.32 ± 0.09	0.24 ± 0.07
$\Upsilon(2S)\pi^+$	4.38 ± 1.21	2.40 ± 0.63
$\Upsilon(3S)\pi^+$	2.15 ± 0.56	1.64 ± 0.40
$h_b(1P)\pi^+$	2.81 ± 1.10	7.43 ± 2.70
$h_b(2P)\pi^+$	4.34 ± 2.07	14.8 ± 6.22
$B^+\bar{B}^{*0} + \bar{B}^0B^{*+}$	86.0 ± 3.6	–
$B^{*+}\bar{B}^{*0}$	–	73.4 ± 7.0

We also report measurement of branching fractions for three-body decays: $\mathcal{B}(\Upsilon(10860) \rightarrow [B\bar{B}^* + c.c.]^+\pi^-) = (28.3 \pm 2.9 \pm 4.6) \times 10^{-3}$ and $\mathcal{B}(\Upsilon(10860) \rightarrow [B^*\bar{B}^*]^+\pi^-) = (14.1 \pm 1.9 \pm 2.4) \times 10^{-3}$. For the $\Upsilon(10860) \rightarrow B\bar{B}\pi$ decay, we calculate a 90% confidence level upper limit of $\mathcal{B}(\Upsilon(10860) \rightarrow [B\bar{B}]^+\pi^-) < 4.0 \times 10^{-3}$ (including systematic uncertainty). In addition, we report the ratio of the branching fractions $\frac{\mathcal{B}(Z_b(10610) \rightarrow BB^*)}{\sum_n \mathcal{B}(Z_b(10610) \rightarrow \Upsilon(nS)\pi) + \sum_m \mathcal{B}(Z_b(10610) \rightarrow h_b(mP))} = 6.2 \pm 0.7 \pm 1.3_{-1.8}^{+0.0}$ and $\frac{\mathcal{B}(Z_b(10650) \rightarrow B^*B^*)}{\sum_n \mathcal{B}(Z_b(10650) \rightarrow \Upsilon(nS)\pi) + \sum_m \mathcal{B}(Z_b(10650) \rightarrow h_b(mP))} = 2.8 \pm 0.4 \pm 0.6_{-0.4}^{+0.0}$.

We also find it useful to calculate the relative fractions for Z_b decays assuming that they are saturated by the already observed $\Upsilon(nS)$ ($n = 1, 2, 3$), $h_b(mP)$ ($m = 1, 2$), and $B^*B^{(*)}$ channels. Combining results reported here with results on amplitude analysis from Ref. [5] one calculate relative fractions summarized in Table 2. We do not include the $Z_b(10650) \rightarrow BB^*$ channel in the table as this decay mode has marginal significance. However, if the central value is used, its fraction would be $25.4 \pm 10.2\%$. All other fractions would be reduced by a factor of 1.33.

All presented results are preliminary.

5. Acknowledgments

This work is partially supported by the Russian Foundation for Basic Research Grants RFBR 12-02-01296, 12-02-01032 and by the grant of the Russian Federation government 11.G34.31.0047

References

- [1] A. Bondar, A. Garmash, R. Mizuk *et al.* (Belle Collaboration), Phys. Rev. Lett. **108**, 122001 (2012).
- [2] I. Adachi *et al.* (Belle Collaboration), arXiv:1105.4583 [hep-ex].
- [3] A.E. Bondar, A. Garmash, A.I. Milstein, R. Mizuk, M.B. Voloshin, Phys. Rev. D **84**, 054010 (2011).
- [4] I. Adachi *et al.*, (Belle Collaboration), [arXiv:1207.4345].
- [5] I. Adachi *et al.*, (Belle Collaboration), [arXiv:1209.6450].