

## On the pair correlations of neutral $K$ , $D$ , $B$ and $B_s$ mesons with close momenta produced in inclusive multiparticle processes

**Valery V. LYUBOSHITZ\***

*Joint Institute for Nuclear Research (Dubna, Russia)*

*E-mail: Valery.Lyuboshitz@jinr.ru*

**Vladimir L. Lyuboshitz**

*Joint Institute for Nuclear Research (Dubna, Russia)*

The phenomenological structure of inclusive cross-sections of the production of two neutral  $K$  mesons in hadron–hadron, hadron–nucleus and nucleus–nucleus collisions is theoretically investigated taking into account the strangeness conservation in strong and electromagnetic interactions. Relations describing the dependence of the correlations of two short-lived and two long-lived neutral kaons  $K_S^0 K_S^0$ ,  $K_L^0 K_L^0$  and the correlations of "mixed" pairs  $K_S^0 K_L^0$  at small relative momenta upon the space-time parameters of the generation region of  $K^0$  and  $\bar{K}^0$  mesons have been obtained. It is shown that under the strangeness conservation the correlation functions of the pairs  $K_S^0 K_S^0$  and  $K_L^0 K_L^0$ , produced in the same inclusive process, coincide, and the difference between the correlation functions of the pairs  $K_S^0 K_S^0$  and  $K_S^0 K_L^0$  is conditioned exclusively by the production of the pairs of non-identical neutral kaons  $K^0 \bar{K}^0$ .

For comparison, analogous correlations for the pairs of neutral heavy mesons  $D^0$ ,  $B^0$  and  $B_s^0$ , generated in multiple inclusive processes with charm ( beauty ) conservation, are also theoretically analyzed . These correlations are described by quite similar expressions: in particular, just as for  $K^0$  mesons, the correlation functions for the pairs of states with the same  $CP$  parity ( $R_{SS} = R_{LL}$ ) and with different  $CP$  parity ( $R_{SL}$ ) do not coincide, and the difference between them is conditioned exclusively by the production of pairs  $D^0 \bar{D}^0$ ,  $B^0 \bar{B}^0$  and  $B_s^0 \bar{B}_s^0$  . However, contrary to the case of  $K^0$  mesons, here the distinction of  $CP$ -even and  $CP$ -odd states encounters difficulties – due to the insignificant differences of their lifetimes and the relatively small probability of purely  $CP$ -even and  $CP$ -odd decay channels. Nevertheless, one may hope that it will become possible at future colliders.

*XXVII International Symposium on Lepton Photon Interactions at High Energies*

*17-22 August 2015*

*Ljubljana, Slovenia*

---

\* Speaker.

## 1. Consequences of the strangeness conservation for neutral kaons

In the work [1] the properties of the density matrix of two neutral  $K$  mesons, following from the strangeness conservation in strong and electromagnetic interactions, have been investigated. By definition, the diagonal elements of the non-normalized two-particle density matrix coincide with the two-particle structure functions, which are proportional to the double inclusive cross-sections.

Taking into account the strangeness conservation, the pairs of neutral kaons  $K^0 K^0$ ,  $\bar{K}^0 \bar{K}^0$  and  $K^0 \bar{K}^0$  are produced incoherently. This means that in the  $K^0$ - $\bar{K}^0$ - representation the non-diagonal elements of the density matrix between the states  $K^0 K^0$  and  $\bar{K}^0 \bar{K}^0$ ,  $K^0 K^0$  and  $K^0 \bar{K}^0$ ,  $\bar{K}^0 \bar{K}^0$  and  $K^0 \bar{K}^0$  are equal to zero. However, the non-diagonal elements of the two-kaon density matrix between the two states  $|K^0\rangle(\mathbf{p}_1)|\bar{K}^0\rangle(\mathbf{p}_2)$  and  $|\bar{K}^0\rangle(\mathbf{p}_1)|K^0\rangle(\mathbf{p}_2)$  with the zero strangeness are not equal to zero, in general ( here  $\mathbf{p}_1$  and  $\mathbf{p}_2$  are the momenta of the first and second kaons ).

The internal states of  $K^0$  meson (strangeness  $S = 1$ ) and  $\bar{K}^0$  meson ( $S = -1$ ) are as follows :

$$|K^0\rangle = \frac{1}{\sqrt{2}}(|K_S^0\rangle + |K_L^0\rangle), \quad |\bar{K}^0\rangle = \frac{1}{\sqrt{2}}(|K_S^0\rangle - |K_L^0\rangle).$$

where  $K_S^0$  is the short-lived neutral kaon and  $K_L^0$  is the long-lived one. Neglecting the small effect of  $CP$  non-invariance, the  $CP$  parity of the state  $K_S^0$  is equal to  $(+1)$ , and the  $CP$  parity of the state  $K_L^0$  is equal to  $(-1)$ . Meantime, the  $CP$  parity of the system  $K^0 \bar{K}^0$  is always positive [2].

The system of two non-identical neutral kaons  $K^0 \bar{K}^0$  in the symmetric internal state, corresponding to even orbital momenta, is decomposed into the schemes  $|K_S^0\rangle|K_S^0\rangle$  and  $|K_L^0\rangle|K_L^0\rangle$  [2]; meantime, the system  $K^0 \bar{K}^0$  in the antisymmetric internal state, corresponding to odd orbital momenta, is decomposed into the scheme  $|K_S^0\rangle|K_L^0\rangle$  [2].

The strangeness conservation leads to the fact that all the double inclusive cross-sections of production of pairs  $K_S^0 K_S^0$ ,  $K_L^0 K_L^0$  and  $K_S^0 K_L^0$  (two-particle structure functions) prove to be symmetric with respect to the permutation of momenta  $\mathbf{p}_1$  and  $\mathbf{p}_2$ , and, besides, prove to be invariant with respect to the replacement of the short-lived state  $K_S^0$  by the long-lived state  $K_L^0$ , and *vice versa* [1]:  $f_{SS}(\mathbf{p}_1, \mathbf{p}_2) = f_{LL}(\mathbf{p}_1, \mathbf{p}_2)$ ,  $f_{SL}(\mathbf{p}_1, \mathbf{p}_2) = f_{LS}(\mathbf{p}_1, \mathbf{p}_2)$ . In doing so,

$$f_{SS}(\mathbf{p}_1, \mathbf{p}_2) - f_{SL}(\mathbf{p}_1, \mathbf{p}_2) = \text{Re} \rho_{K^0 \bar{K}^0 \rightarrow \bar{K}^0 K^0}(\mathbf{p}_1, \mathbf{p}_2), \quad (1.1)$$

where  $\rho_{K^0 \bar{K}^0 \rightarrow \bar{K}^0 K^0}(\mathbf{p}_1, \mathbf{p}_2) = (\rho_{\bar{K}^0 K^0 \rightarrow K^0 \bar{K}^0}(\mathbf{p}_1, \mathbf{p}_2))^*$  are the non-diagonal elements of the two-kaon density matrix.

## 2. Structure of pair correlations of identical and non-identical neutral kaons with close momenta

Now let us consider, within the model of one-particle sources [2-7], the correlations of pairs of neutral  $K$  mesons with close momenta ( see also, e.g., [8-10] ). In the case of the identical states  $K_S^0 K_S^0$  and  $K_L^0 K_L^0$  we obtain the following expressions for the correlation functions  $R_{SS}$ ,  $R_{LL}$  ( proportional to the structure functions ), normalized to unity at large relative momenta:

$$R_{SS}(\mathbf{k}) = R_{LL}(\mathbf{k}) = \lambda_{K^0 K^0} [1 + F_{K^0}(2\mathbf{k}) + 2b_{\text{int}}(\mathbf{k})] + \lambda_{\bar{K}^0 \bar{K}^0} [1 + F_{\bar{K}^0}(2\mathbf{k}) + 2\tilde{b}_{\text{int}}(\mathbf{k})] + \lambda_{K^0 \bar{K}^0} [1 + F_{K^0 \bar{K}^0}(2\mathbf{k}) + 2B_{\text{int}}(\mathbf{k})]. \quad (2.1)$$

Here  $\mathbf{k}$  is the momentum of one of the kaons in the c.m. frame of the pair, and the quantities  $\lambda_{K^0 K^0}$ ,  $\lambda_{\bar{K}^0 \bar{K}^0}$  and  $\lambda_{K^0 \bar{K}^0}$  are the relative fractions of the average numbers of produced pairs  $K^0 K^0$ ,  $\bar{K}^0 \bar{K}^0$  and  $K^0 \bar{K}^0$ , respectively ( $\lambda_{K^0 K^0} + \lambda_{\bar{K}^0 \bar{K}^0} + \lambda_{K^0 \bar{K}^0} = 1$ ). The "formfactors"  $F_{K^0}(2\mathbf{k})$ ,  $F_{\bar{K}^0}(2\mathbf{k})$  and  $F_{K^0 \bar{K}^0}(2\mathbf{k})$  appear due to the contribution of Bose statistics:

$$\begin{aligned} F_{K^0}(2\mathbf{k}) &= \int W_{K^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}, & F_{\bar{K}^0}(2\mathbf{k}) &= \int W_{\bar{K}^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}, \\ F_{K^0 \bar{K}^0}(2\mathbf{k}) &= \int W_{K^0 \bar{K}^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}. \end{aligned} \quad (2.2)$$

where  $W_{K^0}(\mathbf{r})$ ,  $W_{\bar{K}^0}(\mathbf{r})$  and  $W_{K^0 \bar{K}^0}(\mathbf{r})$  are the probability distributions of distances between the sources of emission of two  $K^0$  mesons, two  $\bar{K}^0$  mesons and the  $K^0$  and  $\bar{K}^0$  mesons, respectively, in the c.m. frame of the kaon pair. Meantime, the quantities  $b_{\text{int}}(\mathbf{k})$ ,  $\tilde{b}_{\text{int}}(\mathbf{k})$  and  $B_{\text{int}}(\mathbf{k})$  describe the respective contributions of the  $S$ -wave final-state interaction of two  $K^0$  mesons, two  $\bar{K}^0$  mesons and of the  $K^0$  and  $\bar{K}^0$  mesons.

Due to the  $CP$  invariance, the quantities  $b_{\text{int}}(\mathbf{k})$  and  $\tilde{b}_{\text{int}}(\mathbf{k})$  can be expressed by means of averaging the same function  $b(\mathbf{k}, \mathbf{r})$  over the different distributions:

$$b_{\text{int}}(\mathbf{k}) = \int W_{K^0}(\mathbf{r}) b(\mathbf{k}, \mathbf{r}) d^3\mathbf{r}, \quad \tilde{b}_{\text{int}}(\mathbf{k}) = \int W_{\bar{K}^0}(\mathbf{r}) b(\mathbf{k}, \mathbf{r}) d^3\mathbf{r}.$$

Meantime,  $B_{\text{int}}(\mathbf{k}) = \int W_{K^0 \bar{K}^0}(\mathbf{r}) B(\mathbf{k}, \mathbf{r}) d^3\mathbf{r}$ , where  $B(\mathbf{k}, \mathbf{r}) \neq b(\mathbf{k}, \mathbf{r})$ .

Let us emphasize that when the pair of non-identical neutral kaons  $K^0 \bar{K}^0$  is produced but the pair of identical quasistationary states  $K_S^0 K_S^0$  ( or  $K_L^0 K_L^0$  ) is registered over decays, the two-particle correlations at small relative momenta have the same character as in the case of usual identical bosons with zero spin [2].

For the pairs of non-identical kaon states  $K_S^0 K_L^0$ , the correlation functions at small relative momenta have the form:

$$\begin{aligned} R_{SL}(\mathbf{k}) = R_{LS}(\mathbf{k}) &= \lambda_{K^0 K^0} [1 + F_{K^0}(2\mathbf{k}) + 2b_{\text{int}}(\mathbf{k})] + \\ &+ \lambda_{\bar{K}^0 \bar{K}^0} [1 + F_{\bar{K}^0}(2\mathbf{k}) + 2\tilde{b}_{\text{int}}(\mathbf{k})] + \lambda_{K^0 \bar{K}^0} [1 - F_{K^0 \bar{K}^0}(2\mathbf{k})]. \end{aligned} \quad (2.3)$$

It follows from Eqs. (2.1) and (2.3) that the correlation functions of pairs of neutral  $K$  mesons with close momenta, which are created in inclusive processes, satisfy the relation :

$$R_{SS}(\mathbf{k}) - R_{SL}(\mathbf{k}) = 2\lambda_{K^0 \bar{K}^0} [F_{K^0 \bar{K}^0}(2\mathbf{k}) + B_{\text{int}}(\mathbf{k})]. \quad (2.4)$$

We see that the difference between the correlation functions of the pairs of identical neutral kaons  $K_S^0 K_S^0$  and pairs of non-identical neutral kaons  $K_S^0 K_L^0$  is conditioned exclusively by the generation of  $K^0 \bar{K}^0$  pairs.

Relations connecting the contribution of the  $S$ -wave strong interaction into the pair correlations of particles at small relative momenta with the parameters of low-energy scattering were obtained earlier in the papers [4-7]. It is essential that the "formfactors" (2.2) and the functions

$b_{\text{int}}(\mathbf{k})$ ,  $\tilde{b}_{\text{int}}(\mathbf{k})$  and  $B_{\text{int}}(\mathbf{k})$  depend on the space-time parameters of the generation region of neutral kaons and tend to zero at high values of the relative momentum  $q = 2|\mathbf{k}|$  of two neutral kaons. Concretely, the expression for the function  $B(\mathbf{k}, \mathbf{r})$  in the case of the  $K^0 \bar{K}^0$  system has been obtained in the paper [10], and – in the same paper – the estimate of contribution of the transition  $K^+ K^- \rightarrow K^0 \bar{K}^0$  has also been presented .

### 3. Correlations of neutral heavy mesons

Formally, analogous relations are valid also for the neutral heavy mesons  $D^0$ ,  $B^0$  and  $B_s^0$  . In doing so, the role of strangeness conservation is played, respectively, by the conservation of charm and beauty in inclusive multiple processes with production of these mesons . In these cases the quasistationary states are also states with definite  $CP$  parity ( neglecting the weak effects of  $CP$  nonconservation ). For example,

$$|B_s^0\rangle = \frac{1}{\sqrt{2}}(|B^0\rangle + |\bar{B}^0\rangle), \text{ } CP \text{ parity } (+1); \quad |B_L^0\rangle = \frac{1}{\sqrt{2}}(|B^0\rangle - |\bar{B}^0\rangle), \text{ } CP \text{ parity } (-1).$$

The difference of masses between the respective  $CP$ -odd and  $CP$ -even states is very insignificant in all the cases, ranging from  $10^{-12}$  MeV for  $K^0$  mesons up to  $10^{-8}$  MeV for  $B_s^0$  mesons . Concerning the lifetimes of these states, in the case of  $K^0$  mesons they differ by 600 times, but for  $D^0$ ,  $B^0$  and  $B_s^0$  mesons the respective lifetimes are almost the same. In connection with this, it is practically impossible to distinguish the states of  $D^0$ ,  $B^0$  and  $B_s^0$  mesons with definite  $CP$  parity by the difference in their lifetimes. These states, in principle, can be identified through the purely  $CP$ -even and purely  $CP$ -odd decay channels ; however, in fact the branching ratio for such decays is very small, for example,

$$Br(D^0 \rightarrow \pi^+ \pi^-) = 1.62 \cdot 10^{-3} \quad (CP = +1);$$

$$Br(D^0 \rightarrow K^+ K^-) = 4.25 \cdot 10^{-3} \quad (CP = +1);$$

$$Br(B_s^0 \rightarrow J/\Psi \pi^0) < 1.2 \cdot 10^{-3} \quad (CP = +1);$$

$$Br(B^0 \rightarrow J/\Psi K_s^0) = 9 \cdot 10^{-4} \quad (CP = -1).$$

Just as in the case of neutral  $K$  mesons, the correlation functions for the pairs of states of neutral  $D$ ,  $B$  and  $B_s$  mesons with the same  $CP$  parity ( $R_{SS} = R_{LL}$ ) and for the pairs of states with different  $CP$  parity ( $R_{SL}$ ) do not coincide, and the difference between them is conditioned exclusively by the production of pairs  $D^0 \bar{D}^0$ ,  $B^0 \bar{B}^0$  and  $B_s^0 \bar{B}_s^0$ , respectively. In particular, for  $B_s^0$  mesons the following relation holds:

$$R_{SS}(\mathbf{k}) - R_{SL}(\mathbf{k}) = 2\lambda_{B_s^0 \bar{B}_s^0} \left[ F_{B_s^0 \bar{B}_s^0}(2\mathbf{k}) + B_{\text{int}}(\mathbf{k}) \right]; \quad (3.1)$$

here  $\lambda_{B_s^0 \bar{B}_s^0}$  is the relative fraction of generated pairs  $B_s^0 \bar{B}_s^0$ , and the contributions of Bose statistics ( $F_{B_s^0 \bar{B}_s^0}(2\mathbf{k})$ ) and the  $S$ -wave final-state  $B_s^0 \bar{B}_s^0$ -interaction ( $B_{\text{int}}(\mathbf{k})$ ) are determined analogously to the case of  $K^0 \bar{K}^0$  pairs ( see Section 2 ) :

$$F_{B_s^0 \bar{B}_s^0}(2\mathbf{k}) = \int W_{B_s^0 \bar{B}_s^0}(\mathbf{r}) \cos(2\mathbf{k}\mathbf{r}) d^3\mathbf{r}, \quad B_{\text{int}}(\mathbf{k}) = \int W_{B_s^0 \bar{B}_s^0}(\mathbf{r}) B(\mathbf{k}, \mathbf{r}) d^3\mathbf{r},$$

$$B(\mathbf{k}, \mathbf{r}) = |A_{B_s^0 \bar{B}_s^0}(k)|^2 \frac{1}{r^2} + 2 \operatorname{Re} \left( A_{B_s^0 \bar{B}_s^0}(k) \frac{\exp(ikr) \cos \mathbf{k}\mathbf{r}}{r} \right),$$

where  $A_{B_s^0 \bar{B}_s^0}(k) \equiv A_{B_s^0 \bar{B}_s^0 \rightarrow B_s^0 \bar{B}_s^0}(k)$  is the amplitude of  $S$ -wave  $B_s^0 \bar{B}_s^0$  - scattering,  $k = |\mathbf{k}|$ ,  $r = |\mathbf{r}|$ . Let us remark that the  $B_s^0$  and  $\bar{B}_s^0$  mesons do not have charged partners ( the isotopic spin equals zero ) and, on account of that, here the transition being similar to  $(K^+ K^- \rightarrow K^0 \bar{K}^0)$  is absent .

#### 4. Summary

1. It is shown that, taking into account the strangeness conservation, the correlation functions for two short-lived neutral  $K$  mesons (  $R_{SS}$  ) and two long-lived neutral  $K$  mesons (  $R_{LL}$  ) are equal to each other. This result is the direct consequence of the strangeness conservation.
2. It is shown that just the production of  $K^0 \bar{K}^0$  pairs with the zero strangeness leads to the difference between the correlation functions  $R_{SS}$  and  $R_{SL}$  of two neutral kaons.
3. The character of analogous correlations for neutral heavy mesons  $D^0, B^0, B_s^0$  with nonzero charm and beauty is discussed . Contrary to the case of  $K^0$  mesons, here the distinction of respective  $CP$ -even and  $CP$ -odd states encounters difficulties, which are connected with the insignificant difference of their lifetimes and the relatively small probability of purely  $CP$  -even and purely  $CP$ -odd decay channels.

#### References

- [1] V.L. Lyuboshitz, *Yad. Fiz.* **23**, 1266 (1976) [ *Sov. J. Nucl. Phys.* **23**, 673 (1976) ].
- [2] V.L. Lyuboshitz and M.I. Podgoretsky, *Yad. Fiz.* **30**, 789 (1979) [ *Sov. J. Nucl. Phys.* **30**, 407 (1979) ].
- [3] M.I. Podgoretsky, *Fiz. Elem. Chast. At. Yadra* **20**, 628 (1989) [ *Sov. J. Part. Nucl.* **20**, 266 (1989) ].
- [4] R. Lednicky and V.L. Lyuboshitz, *Yad. Fiz.* **35**, 1316 (1982) [ *Sov. J. Nucl. Phys.* **35**, 770 (1982) ].
- [5] V.L. Lyuboshitz, *Yad. Fiz.* **41**, 820 (1985) [ *Sov. J. Nucl. Phys.* **41**, 523 (1985) ].
- [6] V.L. Lyuboshitz, *Yad. Fiz.* **48**, 1501 (1988) [ *Sov. J. Nucl. Phys.* **48**, 956 (1988) ].
- [7] R. Lednicky, V.V. Lyuboshitz and V.L. Lyuboshitz, *Yad. Fiz.* **61**, 2161 (1998) [ *Phys. At. Nucl.* **61**, 2050 (1998) ].
- [8] V.L. Lyuboshitz and V.V. Lyuboshitz, *Nukleonika*, v.**51**, Suppl. 3, p.S69 (2006) - in Proceedings of Quark Matter 2005 Conference ( Budapest, Hungary, August 4 - 9, 2005 ).
- [9] V.L. Lyuboshitz and V.V. Lyuboshitz, in Proceedings of the IV International Conference on Quarks and Nuclear Physics – QNP-06 ( Madrid, Spain, June 5 - 10, 2006 ), Berlin - Heidelberg, 2007, p.109 .
- [10] V.L. Lyuboshitz and V.V. Lyuboshitz, *Pis'ma v EchaYa*, **4**, 654 (2007) [ *Physics of Particles and Nuclei Letters*, **4**, 388 (2007) ].