

LHC-13 and Neutrinos

P. S. Bhupal Dev*

*Department of Physics and McDonnell Center for the Space Sciences, Washington University,
St. Louis, MO 63130, USA*

E-mail: bdev@wustl.edu

We review the current status and future prospects of testing the low-scale seesaw models of neutrino mass generation at the energy frontier, with particular emphasis on the ongoing searches at the $\sqrt{s} = 13$ TeV LHC. We mostly focus on the type-I seesaw, both within the SM gauge group and in extended gauge theories, and discuss the current limits and the future LHC reach for the extra fermion, gauge and Higgs bosons.

*XVII International Workshop on Neutrino Telescopes
13-17 March 2017
Venezia, Italy*

*Speaker.

1. Introduction

The observed phenomenon of neutrino oscillations requires at least two of the three active neutrinos to have non-zero masses, which is by far the only convincing laboratory evidence for physics beyond the Standard Model (SM). So understanding the origin of neutrino mass could shed light on the underlying new physics scale. A natural way to generate neutrino masses is by breaking the $B - L$ symmetry of the SM, parametrized by the dimension-5 operator $LLHH/\Lambda$ [1], where L and H are respectively the SM lepton and Higgs doublets, and Λ is the new physics scale. There are three tree-level realizations of this operator, commonly known as the type-I [2], type-II [3] and type-III [4] seesaw mechanisms, depending on whether the products of L and H form an $SU(2)_L$ fermion singlet $(L^\top H)(L^\top H)/\Lambda$, scalar triplet $(L^\top \sigma_a L)(H^\top \sigma_a H)/\Lambda$ or fermion triplet $(L^\top \sigma_a H)(L^\top \sigma_a H)/\Lambda$, respectively (σ_a 's being the usual Pauli matrices). All of these seesaw mechanisms generically predict lepton number violation (LNV), as well as charged lepton flavor violation (cLFV). These effects might be observable in both energy and intensity frontier experiments, provided the seesaw scale is below a few TeV or so, i.e. within an experimentally accessible range. There exists a plethora of such low-scale seesaw models; for a review, see e.g. Ref. [5]. In this proceedings, we mainly focus on the testable aspects of the simplest seesaw paradigm, viz. the type-I seesaw scenario, though other seesaw scenarios are also briefly mentioned. We summarize the current status and future prospects of the searches for the seesaw messengers at the energy frontier, with particular emphasis on the ongoing Large Hadron Collider (LHC) experiments. For details, see e.g. Ref. [6] and references therein.

2. Minimal Type-I seesaw

This is the simplest extension of the SM for understanding the small neutrino masses. It just requires the addition of SM-singlet Majorana fermions, known as sterile neutrinos N_α , to the SM particle content. The relevant piece of the Lagrangian is given by

$$-\mathcal{L} = (Y_\nu)_{\ell\alpha} \bar{L}_\ell \tilde{H} N_\alpha + \frac{1}{2} (M_N)_{\alpha\beta} \bar{N}_\alpha^c N_\beta + \text{H.c.}, \quad (2.1)$$

with $\tilde{H} = i\sigma_2 H^*$ and L, H being the $SU(2)_L$ -doublet lepton and Higgs fields, respectively. The origin of the Majorana mass term M_N can be readily explained in an ultraviolet (UV) completion of seesaw, such as the left-right (LR) symmetric model [7] or $SO(10)$ grand unified theory (GUT) [8]. After electroweak symmetry breaking by the Higgs vacuum expectation value (VEV) v , Eq. (2.1) leads to a Dirac neutrino mass term $M_D = vY_\nu$ which, together with the Majorana mass M_N , induces the tree-level active neutrino masses by the seesaw formula [2]

$$M_\nu \simeq -M_D M_N^{-1} M_D^\top. \quad (2.2)$$

In a bottom-up phenomenological approach, the mass scale of the sterile neutrinos, synonymous with the seesaw scale, is *a priori* unknown, and could be anywhere ranging from sub-eV scale up to the sub-GUT scale $\sim 10^{14}$ GeV. However, theoretical arguments based on the naturalness of the SM Higgs mass of 125 GeV against radiative effects induced by the neutrino loop suggest the

seesaw scale to be below $\sim 10^7$ GeV [9]. Of particular interest to us are TeV-scale seesaw models which are kinematically accessible to the current and foreseeable future collider energies.

In the minimal type-I seesaw with only the SM gauge group, the active-sterile neutrino mixing parameter $V_{\ell N} \equiv M_D M_N^{-1}$ is also required to be sizable, in addition to a low seesaw-scale, to get an observable production cross section at colliders [10]. In the canonical seesaw [2], the active-sterile neutrino mixing parameter is suppressed by the light neutrino mass scale $M_\nu \lesssim 0.1$ eV:

$$V_{\ell N} \simeq \sqrt{\frac{M_\nu}{M_N}} \lesssim 10^{-6} \sqrt{\frac{100 \text{ GeV}}{M_N}}. \quad (2.3)$$

Thus for a TeV-scale seesaw, the experimental effects of the active-sterile neutrino mixing are naively expected to be too small. However, there exist low-scale type-I seesaw models [11, 12], where $V_{\ell N}$ can be sizable due to specific textures of the Dirac and Majorana mass matrices in Eq. (2.2). These textures can in principle be stabilized by imposing some discrete symmetry in the leptonic sector. But most of these low-scale seesaw scenarios require the sterile neutrinos having large mixing with the active sector to be quasi-Dirac, thus suppressing all LNV effects, except when the mass splitting between the pseudo-Dirac sterile neutrino pair is comparable to their decay width, which could lead to a resonant enhancement of the LNV amplitude [13]. On the other hand, if the heavy sterile neutrinos have additional gauge interactions, e.g. when they are charged under the $SU(2)_R$ gauge group in LR seesaw models, one could get large LNV signals at colliders, irrespective of their mixing with the active neutrino sector [14]. Note that the LNV effects involving only the electron flavor are strongly constrained from neutrinoless double beta decay searches [12, 15], but those involving the muon and/or tau flavors could still allow for an observable signal at colliders, depending on the model construction.

Another natural realization of low-scale seesaw with potentially large active-sterile neutrino mixing is the inverse seesaw mechanism [16], which introduces two sets of SM-singlet pseudo-Dirac fermions (N, S) with only a small Majorana mass term μ_S for S . In this case, the magnitude of the light neutrino mass can be decoupled from the heavy neutrino mass scale:

$$M_\nu \simeq (M_D M_N^{-1}) \mu_S (M_D M_N^{-1})^\top, \quad (2.4)$$

thus allowing for large active-sterile mixing even for TeV-scale seesaw without any fine-tuning:

$$V_{\ell N} \simeq \sqrt{\frac{M_\nu}{\mu_S}} \approx 10^{-2} \sqrt{\frac{1 \text{ keV}}{\mu_S}}. \quad (2.5)$$

Note that the smallness of μ_S is technically natural, i.e. in the limit of $\mu_S \rightarrow 0$, lepton number symmetry is restored and the light neutrinos are exactly massless to all orders in perturbation theory. However, the smallness of μ_S implies that all LNV effects will be suppressed. This could be partially overcome by introducing a Majorana mass term μ_N for the N fermions [17].

The hadron collider experiments can *simultaneously* probe the Majorana nature of the neutrinos and the active-sterile neutrino mixing parameters through the ‘‘smoking gun’’ LNV signature of same-sign dilepton plus two jets: $pp \rightarrow N \ell^\pm \rightarrow \ell^\pm \ell^\pm jj$ [10]. This is in contrast with the complementary low-energy probes at the intensity frontier [18], which are mostly sensitive to only one

aspect, e.g. neutrinoless double beta decay for the Majorana nature and cLFV searches for the active-sterile neutrino mixing. The current direct search limits using the same-sign dilepton channel at $\sqrt{s} = 8$ TeV LHC range from $|V_{\ell N}|^2 \lesssim 10^{-2} - 1$ (with $\ell = e, \mu$) for $M_N = 100 - 500$ GeV [19]. These limits could be improved by roughly an order of magnitude and extended for heavy neutrino masses up to a TeV or so with the $\sqrt{s} = 14$ TeV LHC and further improvements by another order of magnitude are possible at the proposed 100 TeV pp collider [6]. It is worth emphasizing that the $W\gamma$ vector boson fusion processes [20] become increasingly important at higher center-of-mass energies and/or higher masses, and must be taken into account, along with the usual Drell-Yan production mechanism via s -channel W boson so far considered in the experimental analyses.

To test seesaw models with suppressed LNV effects, as well as to distinguish between Dirac and Majorana nature of the sterile neutrinos, one should also study the opposite-sign dilepton signal $pp \rightarrow N\ell^\pm \rightarrow \ell^\pm \ell^\mp jj$ and its ratio with the same-sign signal [21]. Although the opposite-sign dilepton signal suffers from a huge SM background, mostly from $Z \rightarrow \ell^+ \ell^-$ decays, one can exploit specific kinematic features to get a good signal-to-background ratio [22]. Another promising channel is the trilepton mode $pp \rightarrow N\ell^\pm \rightarrow \ell^\pm \ell^\mp \ell^\pm + \cancel{E}_T$ [23], which has a relatively smaller cross section, but a smaller SM background as well. In addition, there exist indirect signals for electroweak-scale sterile neutrinos which can probe large active-sterile neutrino mixing, irrespective of their Majorana nature, such as anomalous Higgs [24] and W/Z decays [25]. For sterile neutrinos lighter than the W boson, one could also look for displaced vertex signatures [26]. GeV-scale sterile neutrinos are also constrained at LHCb from searches in rare B -meson decays [27]. In future, this mass range can be more effectively probed in beam dump experiments, such as SHiP [28].

A future lepton collider, such as ILC or FCC-ee, will provide better sensitivity up to sterile neutrino masses very close to its kinematic threshold [29], mainly due to its relatively cleaner environment, as compared to hadron colliders. See also Ref. [30] for sterile neutrino studies at a future electron-proton collider, such as LHeC. For a summary of all relevant constraints and future prospects for sterile neutrinos over MeV-TeV mass range, see Refs. [6, 31]. Note that the cLFV processes (such as $\mu \rightarrow e\gamma$) put stringent constraints on the product $|V_{\ell N} V_{\ell' N}^*|$ (with $\ell \neq \ell'$), but do not restrict the individual mixing parameters $|V_{\ell N}|^2$ in a model-independent way. Similarly in the electron sector, the $0\nu\beta\beta$ constraints are the most stringent for a large range of the heavy neutrino masses, but are subject to a large uncertainty due to the unknown CP phases in the seesaw matrix, and hence, do not necessarily render the direct searches in the electron channel redundant.

3. Left-Right Seesaw

In gauge extensions of the SM seesaw, the new gauge interactions of the heavy neutrinos allow them to be produced copiously at the LHC without necessarily involving the light-heavy neutrino mixing, as long as the new gauge bosons are in the kinematically accessible range. Of particular interest is the left-right seesaw, based on the gauge group $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ [7], which provides a natural framework for low-scale type-I seesaw embedding. The two key features of seesaw, namely, the existence of right-handed (RH) neutrinos and their Majorana mass, naturally arise in this set-up and the seesaw scale is intimately connected with the $U(1)_{B-L}$ symmetry breaking, which can be realized at $\nu_R \gtrsim 5$ TeV, with many observable effects at both energy and intensity frontiers.

At the energy frontier, the smoking gun signal of LR seesaw is the LNV process of same-sign dilepton plus dijet production [14], which can probe the heavy W_R gauge boson masses upto 5-6 TeV at the high-luminosity LHC. One could also get displaced vertex signals at the LHC, if the RH neutrinos are in the few GeV range, in which case W_R masses can be probed up to about 20 TeV [32]. At the intensity frontier, the new RH current effects can also manifest in the form of enhanced cLFV or $0\nu\beta\beta$ rates [6], as well as at future beam dump experiments, such as SHiP [33, 28].

4. Type-II seesaw

Unlike the sterile neutrinos in the minimal type-I seesaw which, being SM gauge singlets, can only communicate with the SM sector through their mixing with active neutrinos, the type-II seesaw messenger, being an $SU(2)_L$ -triplet scalar ($\Delta^{++}, \Delta^+, \Delta^0$), can be *directly* produced at the LHC via its gauge interactions. The smoking gun collider signal in this case would be the detection of the doubly-charged scalars with LNV interactions. The most promising channels at the LHC are $pp \rightarrow Z^*/\gamma^* \rightarrow \Delta^{++}\Delta^{--} \rightarrow \ell^+\ell^+\ell^-\ell^-$ and $pp \rightarrow W^\pm \rightarrow \Delta^\pm\Delta^\mp \rightarrow \ell^\pm\ell^\pm\ell^\mp\nu_\ell$ using which lower limits on the triplet mass of up to 600 GeV have been derived at the LHC [34]. These limits could be marginally improved up to about 800 GeV at the $\sqrt{s} = 14$ TeV LHC, whereas a future 100 TeV pp collider could probe up to 5 TeV or so [35], depending on the model parameters.

In the case of left-right seesaw, apart from the doubly-charged scalars, there exist new CP-even and odd neutral, as well as singly-charged Higgs bosons, all associated with the spontaneous breaking of $SU(2)_R \times U(1)_{B-L}$ symmetry [35]. However, the low-energy flavor-changing neutral current (FCNC) constraints impose stringent lower bounds of $\mathcal{O}(10)$ TeV on the bidoublet CP-even, odd and singly-charged Higgs bosons [36]. So there is no hope for them at the LHC. But the CP-even neutral triplet component is largely unrestricted experimentally, primarily because of its hadrophobic nature [32]. It could in principle be very light, in the GeV range, in which case the FCNC constraints necessarily imply that it will be sufficiently long-lived at the LHC with new displaced diphoton signals [37].

5. Type-III seesaw

Similar to the type-II seesaw case, the type-III seesaw messenger, being an $SU(2)_L$ -triplet fermion ($\Sigma^+, \Sigma^0, \Sigma^-$), can be directly produced at the LHC via its gauge interactions. The most promising production channels at the LHC are $pp \rightarrow W^{\pm*} \rightarrow \Sigma^\pm\Sigma^0$ and $pp \rightarrow Z^*/\gamma^*/h^* \rightarrow \Sigma^+\Sigma^-$, which lead to multi-lepton final states. Depending on the theoretical scenario considered, the $\sqrt{s} = 8$ TeV LHC data has excluded fermion triplets of mass up to 540 GeV at 95% CL [38]. These limits could be improved up to about 1 TeV at the $\sqrt{s} = 14$ TeV LHC [39].

6. Conclusion

Neutrino oscillations have provided us with the first (and so far only) conclusive experimental evidence of physics beyond the SM. Therefore, it is very important to explore the experimental signatures of neutrino mass models, which might lead to some crucial insights into the underlying new

physics. A simple theoretical paradigm for neutrino masses is the seesaw mechanism, which also provides potentially attractive solutions to other outstanding puzzles like dark matter and baryon asymmetry of the universe. We have briefly reviewed the current status and future prospects of the direct searches for various seesaw messengers at the energy frontier, and show that colliders offer an ideal testing ground for low-scale seesaw models, which is complementary to the low-energy probes at the intensity frontier.

References

- [1] S. Weinberg, Phys. Rev. Lett. **43**, 1566 (1979).
- [2] P. Minkowski, Phys. Lett. B **67**, 421 (1977); R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980); T. Yanagida, Conf. Proc. C **7902131**, 95 (1979); M. Gell-Mann, P. Ramond and R. Slansky, Conf. Proc. C **790927**, 315 (1979); S. L. Glashow, NATO Sci. Ser. B **61**, 687 (1980).
- [3] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980); T. P. Cheng and L. F. Li, Phys. Rev. D **22**, 2860 (1980); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); R. N. Mohapatra and G. Senjanović, Phys. Rev. D **23**, 165 (1981).
- [4] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C **44**, 441 (1989).
- [5] S. M. Boucenna, S. Morisi and J. W. F. Valle, Adv. High Energy Phys. **2014**, 831598 (2014).
- [6] F. F. Deppisch, P. S. B. Dev and A. Pilaftsis, New J. Phys. **17**, 075019 (2015).
- [7] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 2558 (1975); G. Senjanović and R. N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
- [8] H. Fritzsch and P. Minkowski, Annals Phys. **93**, 193 (1975).
- [9] F. Vissani, Phys. Rev. D **57**, 7027 (1998); J. D. Clarke, R. Foot and R. R. Volkas, Phys. Rev. D **91**, 073009 (2015); G. Bambhaniya, P. S. B. Dev, S. Goswami, S. Khan and W. Rodejohann, Phys. Rev. D **95**, no. 9, 095016 (2017).
- [10] A. Datta, M. Guchait and A. Pilaftsis, Phys. Rev. D **50**, 3195 (1994); F. M. L. Almeida Jr., Y. D. A. Coutinho, J. A. Martins Simoes and M. A. B. do Vale, Phys. Rev. D **62**, 075004 (2000); O. Panella, M. Cannoni, C. Carimalo and Y. N. Srivastava, Phys. Rev. D **65**, 035005 (2002); T. Han and B. Zhang, Phys. Rev. Lett. **97**, 171804 (2006); F. del Aguila, J. A. Aguilar-Saavedra and R. Pittau, JHEP **0710**, 047 (2007); F. del Aguila and J. A. Aguilar-Saavedra, Nucl. Phys. B **813**, 22 (2009).
- [11] A. Pilaftsis, Z. Phys. C **55**, 275 (1992); D. Tommasini, G. Barenboim, J. Bernabeu and C. Jarlskog, Nucl. Phys. B **444**, 451 (1995); J. Gluza, Acta Phys. Polon. B **33**, 1735 (2002); A. Pilaftsis, Phys. Rev. Lett. **95**, 081602 (2005); J. Kersten and A. Y. Smirnov, Phys. Rev. D **76**, 073005 (2007); Z. z. Xing, Prog. Theor. Phys. Suppl. **180**, 112 (2009); M. B. Gavela, T. Hambye, D. Hernandez and P. Hernandez, JHEP **0909**, 038 (2009); X. G. He, S. Oh, J. Tandean and C. C. Wen, Phys. Rev. D **80**, 073012 (2009); R. Adhikari and A. Raychaudhuri, Phys. Rev. D **84**, 033002 (2011); F. F. Deppisch and A. Pilaftsis, Phys. Rev. D **83**, 076007 (2011); M. Mitra, G. Senjanovic and F. Vissani, Nucl. Phys. B **856**, 26 (2012); C. H. Lee, P. S. B. Dev and R. N. Mohapatra, Phys. Rev. D **88**, 093010 (2013).
- [12] A. Ibarra, E. Molinaro and S. T. Petcov, JHEP **1009**, 108 (2010); Phys. Rev. D **84**, 013005 (2011).
- [13] S. Bray, J. S. Lee and A. Pilaftsis, Nucl. Phys. B **786**, 95 (2007).

- [14] W. Y. Keung and G. Senjanović, *Phys. Rev. Lett.* **50**, 1427 (1983); A. Ferrari *et al.*, *Phys. Rev. D* **62**, 013001 (2000); M. Nemevsek, F. Nesti, G. Senjanovic and Y. Zhang, *Phys. Rev. D* **83**, 115014 (2011); S. P. Das, F. F. Deppisch, O. Kittel and J. W. F. Valle, *Phys. Rev. D* **86**, 055006 (2012); J. A. Aguilar-Saavedra and F. R. Joaquim, *Phys. Rev. D* **86**, 073005 (2012); C. Y. Chen, P. S. B. Dev and R. N. Mohapatra, *Phys. Rev. D* **88**, 033014 (2013); J. N. Ng, A. de la Puente and B. W. P. Pan, *JHEP* **1512**, 172 (2015); P. S. B. Dev, D. Kim and R. N. Mohapatra, *JHEP* **1601**, 118 (2016). M. Mitra, R. Ruiz, D. J. Scott and M. Spannowsky, *Phys. Rev. D* **94**, no. 9, 095016 (2016).
- [15] J. Lopez-Pavon, E. Molinaro and S. T. Petcov, *JHEP* **1511**, 030 (2015); E. Fernandez-Martinez, J. Hernandez-Garcia, J. Lopez-Pavon and M. Lucente, *JHEP* **1510**, 130 (2015).
- [16] R. N. Mohapatra and J. W. F. Valle, *Phys. Rev. D* **34**, 1642 (1986).
- [17] P. S. B. Dev and A. Pilaftsis, *Phys. Rev. D* **86**, 113001 (2012); *Phys. Rev. D* **87**, 053007 (2013); R. L. Awasthi, M. K. Parida and S. Patra, *JHEP* **1308**, 122 (2013); P. Pritimita, N. Dash and S. Patra, *JHEP* **1610**, 147 (2016); N. Haba, H. Ishida and Y. Yamaguchi, *JHEP* **1611**, 003 (2016).
- [18] A. de Gouvea and P. Vogel, *Prog. Part. Nucl. Phys.* **71**, 75 (2013).
- [19] V. Khachatryan *et al.*, *Phys. Lett. B* **748**, 144 (2015); *JHEP* **1604**, 169 (2016); G. Aad *et al.*, *JHEP* **1507**, 162 (2015).
- [20] P. S. B. Dev, A. Pilaftsis and U. K. Yang, *Phys. Rev. Lett.* **112**, 081801 (2014); D. Alva, T. Han and R. Ruiz, *JHEP* **1502**, 072 (2015); A. Das and N. Okada, *Phys. Rev. D* **93**, 033003 (2016); C. Degrande, O. Mattelaer, R. Ruiz and J. Turner, *Phys. Rev. D* **94**, 053002 (2016).
- [21] P. S. B. Dev and R. N. Mohapatra, *Phys. Rev. Lett.* **115**, 181803 (2015); J. Gluza, T. Jeliński and R. Szafron, *Phys. Rev. D* **93**, 113017 (2016); G. Anamiati, M. Hirsch and E. Nardi, *JHEP* **1610**, 010 (2016); A. Das, P. S. B. Dev and R. N. Mohapatra, arXiv:1709.06553 [hep-ph].
- [22] V. Khachatryan *et al.* [CMS Collaboration], *Eur. Phys. J. C* **74**, 3149 (2014).
- [23] F. del Aguila and J. A. Aguilar-Saavedra, *Phys. Lett. B* **672**, 158 (2009); C. Y. Chen and P. S. B. Dev, *Phys. Rev. D* **85**, 093018 (2012); A. Das and N. Okada, *Phys. Rev. D* **88**, 113001 (2013); A. Das, P. S. B. Dev and N. Okada, *Phys. Lett. B* **735**, 364 (2014). G. Bambhaniya, S. Goswami, S. Khan, P. Konar and T. Mondal, *Phys. Rev. D* **91**, 075007 (2015).
- [24] P. S. B. Dev, R. Franceschini and R. N. Mohapatra, *Phys. Rev. D* **86**, 093010 (2012); C. G. Cely, A. Ibarra, E. Molinaro and S. T. Petcov, *Phys. Lett. B* **718**, 957 (2013); A. Das, P. S. B. Dev and C. S. Kim, *Phys. Rev. D* **95**, no. 11, 115013 (2017).
- [25] A. Blondel *et al.*, arXiv:1411.5230 [hep-ex]; E. Izaguirre and B. Shuve, *Phys. Rev. D* **91**, 093010 (2015); C. O. Dib and C. S. Kim, *Phys. Rev. D* **92**, no. 9, 093009 (2015); C. O. Dib, C. S. Kim, K. Wang and J. Zhang, *Phys. Rev. D* **94**, 013005 (2016).
- [26] J. C. Helo, M. Hirsch and S. Kovalenko, *Phys. Rev. D* **89**, 073005 (2014); S. Antusch, E. Cazzato and O. Fischer, *JHEP* **1612**, 007 (2016); L. Duarte, J. Peressutti and O. A. Sampayo, arXiv:1610.03894.
- [27] R. Aaij *et al.* [LHCb Collaboration], *Phys. Rev. Lett.* **112**, 131802 (2014). B. Shuve and M. E. Peskin, *Phys. Rev. D* **94**, 113007 (2016); T. Asaka and H. Ishida, *Phys. Lett. B* **763**, 393 (2016).
- [28] S. Alekhin *et al.*, *Rept. Prog. Phys.* **79**, no. 12, 124201 (2016).
- [29] S. Antusch and O. Fischer, *JHEP* **1505**, 053 (2015); S. Banerjee *et al.*, *Phys. Rev. D* **92**, 075002 (2015); S. Antusch, E. Cazzato and O. Fischer, arXiv:1612.02728 [hep-ph].
- [30] S. Mondal and S. K. Rai, *Phys. Rev. D* **93**, 011702 (2016); *Phys. Rev. D* **94**, 033008 (2016); M. Lindner, F. S. Queiroz, W. Rodejohann and C. E. Yaguna, *JHEP* **1606**, 140 (2016).

- [31] A. Atre, T. Han, S. Pascoli and B. Zhang, *JHEP* **0905**, 030 (2009); A. de Gouvea and A. Kobach, *Phys. Rev. D* **93**, 033005 (2016); P. S. B. Dev and A. Ibarra, *Frascati Phys. Ser.* **61**, 40 (2016); M. Drewes, B. Garbrecht, D. Gueter and J. Klaric, arXiv:1609.09069.
- [32] P. S. B. Dev, R. N. Mohapatra and Y. Zhang, *Nucl. Phys. B* **923**, 179 (2017).
- [33] O. Castillo-Felisola, C. O. Dib, J. C. Helo, S. G. Kovalenko and S. E. Ortiz, *Phys. Rev. D* **92**, no. 1, 013001 (2015).
- [34] G. Aad *et al.*, *JHEP* **1503**, 041 (2015); CMS Collaboration, CMS-PAS-HIG-14-039.
- [35] J. F. Gunion, J. Grifols, A. Mendez, B. Kayser and F. I. Olness, *Phys. Rev. D* **40**, 1546 (1989); P. S. B. Dev, R. N. Mohapatra and Y. Zhang, *JHEP* **1605**, 174 (2016)
- [36] Y. Zhang, H. An, X. Ji and R. N. Mohapatra, *Nucl. Phys. B* **802**, 247 (2008); S. Bertolini, A. Maiezza and F. Nesti, *Phys. Rev. D* **89**, no. 9, 095028 (2014).
- [37] P. S. B. Dev, R. N. Mohapatra and Y. Zhang, *Phys. Rev. D* **95**, no. 11, 115001 (2017); *Acta Phys. Polon. B* **48**, 969 (2017).
- [38] G. Aad *et al.*, *Phys. Rev. D* **92**, 032001 (2015); CMS Collaboration, CMS-PAS-EXO-16-002.
- [39] R. Franceschini, T. Hambye and A. Strumia, *Phys. Rev. D* **78**, 033002 (2008); R. Ruiz, *JHEP* **1512**, 165 (2015).