

Direct-photon elliptic flow in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$

Mike Sas*, for the ALICE Collaboration

University of Utrecht & Nikhef, Netherlands

E-mail: mike.sas@cern.ch

The elliptic flow of inclusive and direct photons was measured by ALICE for central and semi-central Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$. The photons were reconstructed using the electromagnetic calorimeter PHOS and the central tracking system. The inclusive-photon flow reconstructed with both methods are combined and used to extract the direct-photon flow, using a decay-photon cocktail simulation and the direct-photon excess R_γ , in the transverse momentum range $0.9 < p_T < 6.2 \text{ GeV}/c$. We find that the theoretical predictions generally under-predict the results.

*ICHEP 2018, XXXIX International Conference on High Energy Physics
4-11 July 2018
Seoul, South Korea*

*Speaker.

1. Introduction

Ultrarelativistic nucleus–nucleus collisions give access to an experimental study of the Quark-Gluon Plasma (QGP) [1], which is the main goal of the ALICE collaboration. The droplet of hot QCD matter created in AA collisions expands, cools down, and transforms into ordinary matter. The measured elliptic flow of final state particles has been interpreted as being due to collective expansion, which transforms the initial spatial anisotropy into a momentum anisotropy of the final state particles. Here, the emphasis is on the elliptic flow of direct photons. Direct photons are the photons not coming from hadronic decays, and are produced during all stages of the collision. Direct photons are a unique tool to probe the QGP since they leave the system without interacting with the formed medium, because their mean free path is much larger than the size of the system. At low transverse momentum, the direct photons are mainly thermally produced by the hot matter, and their production rate can be used to estimate the temperature of the medium. Additionally, their elliptic flow provides information on the development of flow during the whole evolution of the system. Collective flow is quantified by the azimuthal distribution of particles, which is expanded as $1 + 2 \sum v_n \cos[n(\varphi - \Psi_{RP})]$, where φ is the azimuthal angle of the measured particle, and Ψ_{RP} is the orientation of the reaction plane. The theoretical calculations are predicting a smaller elliptic flow of direct photons compared to that of hadrons. This is because the direct photons probe the momentum anisotropy at the time of their emission, which occurs when the system is still expanding and developing the final momentum anisotropy. The measurement of inclusive- and direct-photon elliptic flow in Pb–Pb collisions is presented and compared to theoretical calculations. More details on the analysis and results can be found in [2].

2. Analysis method

The presented results use the Pb–Pb data recorded by ALICE in 2010. For the measurement of the inclusive-photon flow two methods are used to reconstruct the photons: the Photon Conversion Method (PCM) and the calorimeter PHOS. In the PCM, the Inner Tracking System (ITS) and the Time Projection Chamber (TPC) are used to reconstruct the e^+e^- pairs, which result from a photon converting in the detector material. For PHOS, the photons are reconstructed by measuring the energy they deposit in the calorimeter. The scintillator array detectors V0A and V0C are used for the minimum bias trigger and the calculation of the event plane orientation, which subtend the pseudorapidity ranges $2.8 < \eta < 5.1$ and $-3.7 < \eta < -1.7$, respectively. Furthermore, the events are divided into two centrality classes; central (0–20%) and semi-central (20–40%), according to the summed amplitudes of the V0A and V0C detectors.

The direct-photon elliptic flow is calculated using $v_2^{\gamma,\text{dir}} = (v_2^{\gamma,\text{inc}} R_\gamma - v_2^{\gamma,\text{dec}}) / (R_\gamma - 1)$, where $v_2^{\gamma,\text{inc}}$ is the inclusive-photon flow, $v_2^{\gamma,\text{dec}}$ the decay-photon flow, and $R_\gamma = N_{\gamma,\text{inc}} / N_{\gamma,\text{dec}}$ is the ratio quantifying the excess of direct photons. The ratio R_γ was measured and can be found in [3]. The decay-photon flow is estimated by a simulation, using the existing measurements of hadronic flow. Furthermore, the inclusive-photon elliptic flow is measured with the Scalar Product method, using a pseudorapidity gap of $|\Delta\eta| > 0.9$ between the photon and the reference flow particles. The results from the PCM and PHOS measurement are treated as independent and combined.




Figure 1: (Color online) The elliptic flow of inclusive and decay photons in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV [2], compared to hydrodynamic [4] and transport PHSD [5] model predictions, for the 0–20% (left) and 20–40% (right) centrality classes. The boxes indicate the total uncertainty, and the vertical bars indicate the statistical uncertainty.

3. Results

The elliptic flow of inclusive and decay photons, as measured in two centrality classes, are shown in Fig. 1 [2]. Over the full transverse momentum range of $0.9 < p_T < 6.2$ GeV/c, the flow of inclusive and decay photons is similar, which reflects the fact that the inclusive-photon flow is dominated by the decay photons. Interestingly, the theoretical prediction [4] and [5] overshoots the data by about 40% in the transverse momentum range $1 < p_T < 3$ GeV/c, showing that the identified hadron flow is not fully described in the given models. The direct-photon v_2 is calculated by subtracting the decay-photon flow from the inclusive-photon flow using the direct-photon excess R_γ . Since the significance of $R_\gamma > 1$ is limited, it was chosen to use a Bayesian approach to extract the direct-photon elliptic flow. The true value of R_γ is restricted to be greater than unity, corresponding to a constant prior for $R_{\gamma,\text{true}} \geq 1$. The posterior distributions are sampled for each p_T bin, from which $v_2^{\gamma,\text{dir}}$ is calculated. The resulting direct-photon elliptic flow is shown in Fig. 2, for the centrality classes 0–20% and 20–40%, respectively [2]. The data are compared to theoretical calculations using a state-of-the-art hydrodynamic model [4, 6] and the PHSD transport model [5]. The calculations in general under-predict the data, but the difference is not significant due to the large experimental uncertainties.

4. Summary

The measurement of inclusive- and direct-photon elliptic flow in central and semi-central Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV is presented. The results of the inclusive-photon elliptic flow are the combination of a photon conversion and calorimetric measurement, where both measurements cover the transverse momentum range of $0.9 < p_T < 6.2$ GeV/c. To extract the direct-photon elliptic flow, the decay-photon flow is subtracted from the inclusive-photon elliptic flow, using the previ-




Figure 2: (Color online) The elliptic flow of direct photons in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV [2], compared to hydrodynamic [4] and transport PHSD [5] model predictions in the 0–20% (left) and 20–40% (right) centrality classes. The boxes indicate the total uncertainty, and the vertical bars indicate the statistical uncertainty.

ously measured direct-photon excess R_γ . $v_2^{\gamma,inc}$ shows to be similar to $v_2^{\gamma,dec}$, which is expected since the inclusive-photon sample is dominated by the photons coming from hadronic decays. However, theoretical calculations are over-predicting the data. In addition, $v_2^{\gamma,dir}$ also appears to be close to $v_2^{\gamma,dec}$. The theoretical calculations for the direct-photon elliptic flow are, in contrast to the calculations for inclusive photons, under-predicting the data. However, the uncertainties on the measurements are still too large in order to draw strong conclusions. Future measurements, utilizing a much larger statistics dataset, will significantly improve the precision of the measurement.

References

- [1] S. Borsanyi, Z. Fodor et al., *Full result for the QCD equation of state with 2+1 flavors*, *Phys. Lett. B* **370** (2014) 99 [1309.5258].
- [2] ALICE collaboration, S. Acharya et al., *Direct photon elliptic flow in Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV*, 1805.04403.
- [3] ALICE collaboration, J. Adam et al., *Direct photon production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV*, *Phys. Lett. B* **754** (2016) 235 [1509.07324].
- [4] C. Gale, Y. Hidaka, S. Jeon, S. Lin, J.-F. Paquet, R. D. Pisarski et al., *Production and Elliptic Flow of Dileptons and Photons in a Matrix Model of the Quark-Gluon Plasma*, *Phys. Rev. Lett.* **114** (2015) 072301 [1409.4778].
- [5] O. Linnyk, W. Cassing and E. L. Bratkovskaya, *Centrality dependence of the direct photon yield and elliptic flow in heavy-ion collisions at $\sqrt{s_{NN}} = 200$ GeV*, *Phys. Rev. C* **89** (2014) 034908 [1311.0279].
- [6] R. Chatterjee, P. Dasgupta and D. K. Srivastava, *Anisotropic flow of thermal photons at energies available at the BNL Relativistic Heavy Ion Collider and at the CERN Large Hadron Collider*, *Phys. Rev. C* **96** (2017) 014911 [1702.02378].