

Major Progress in Understanding of GRBs: Discovery of TeraelectronVolt Gamma-Ray Emission

Razmik Mirzoyan

On behalf of the MAGIC Collaboration

Max-Planck-Institute for Physics
Munich, Germany

This year we celebrate the 30 year jubilee of ground-based VHE γ -ray astronomy

- The first 9σ detection of the Crab Nebula marked the birth of the VHE γ -ray astronomy as an independent branch of astronomy
- This detection was reported by the 10m diameter Whipple IACT team in Arizona, lead by the pioneer of VHE γ -ray astronomy Trevor Weekes, in 1989
- With the detection of the first gigantic signal from the GRB190114C (in the first 30 s the gamma-ray rate was x 100 Crab!) we want to celebrate the 30 years jubilee of VHE γ -ray astronomy !



The Astronomer's Telegram

First time detection of a GRB at sub-TeV energies; MAGIC detects the GRB 190114C

ATel #12390; *Razmik Mirzoyan on behalf of the MAGIC Collaboration*
on 15 Jan 2019; 01:03 UT

Credential Certification: *Razmik Mirzoyan (Razmik.Mirzoyan@mpp.mpg.de)*


Subjects: Gamma Ray, >GeV, TeV, VHE, Request for Observations, Gamma-Ray Burst

Referred to by ATel #: 12395


Offline analyses revealed signal $> 50 \sigma$

The MAGIC telescopes performed a rapid follow-up observation of GRB 190114C (Gropp et al., GCN 23688; Tyurina et al., GCN 23690, de Ugarte Postigo et al., GCN 23692, Lipunov et al. GCN 23693, Selsing et al. GCN 23695). This observation was triggered by the Swift-BAT alert; we started observing at about 50s after Swift T0: 20:57:03.19. The MAGIC real-time analysis shows a significance >20 sigma in the first 20 min of observations (starting at T0+50s) for energies >300 GeV. The relatively high detection threshold is due to the large zenith angle of observations (>60 degrees) and the presence of partial Moon. Given the brightness of the event, MAGIC will continue the observation of GRB 190114C until it is observable tonight and also in the next days. We strongly encourage follow-up observations by other instruments. The MAGIC contact persons for these observations are R. Mirzoyan (Razmik.Mirzoyan@mpp.mpg.de) and K. Noda (nodak@icrr.u-tokyo.ac.jp). MAGIC is a system of two 17m-diameter Imaging Atmospheric Cherenkov Telescopes located at the Observatorio Roque de los Muchachos on the Canary island La Palma, Spain, and designed to perform gamma-ray astronomy in the energy range from 50 GeV to

Gamma Ray Bursts



- Most powerful, violent, distant explosions in the Universe
- 2 different populations, short and long bursts
- Long GRBs: $T > \sim 2s$; massive star collapse > ultra-relativistic jet
- Recent detection of a gravitational wave signal consistent with a binary neutron star merger and associated to a short GRB
- Both long and short GRBs have been detected at $E < 100 \text{ GeV}$
- No strict division in time between prompt and afterglow



Past Hint from a GRB @ $E \geq 20$ TeV

AIROBICC & GRB 920925c

- AIROBICC in 1990's was an air Cherenkov integrating array of 7x7 40cm \varnothing PMT-based stations covering $\geq 32000\text{m}^2$
- From GRB 920925c one expected 0.93 events while 11 were observed.
- But the “signal” preceded the WATCH trigger by < 1 minute and was $\sim 9^\circ$ away from its location.
- Evidence $\sim 2.7 \sigma$ (post-trial) from AIROBICC above 20 TeV from GRB 920925c was reported

Padilla et al., 1998





Fig. 6. Map with the situation of the WATCH GRB 920925c and the IPN ring as calculated from the WATCH and ULYSSES observations. The location of the excess detected by AIROBICC nearly coincident in time with WATCH is also shown.




Past Hints from GRBs @ sub-TeV

Milagrito & GRB 970417a

- GRB 970417a was a weak, soft GRB as observed by BATSE. The fluence in the 50–300 keV range was 1.5×10^{-7} ergs cm^{-2} and the T90 period was 7.9 s.
- Milagrito, a prototype of Milagro, consisted of a planar array of 228 PMTs of 8'' size submerged in a light-tight water reservoir with a size of $\sim 42 \times 42 \text{ m}^2$. Milagrito reported evidence for emission above 650 GeV from GRB 970417a, with a (posttrials) probability of 1.5×10^{-3} of being a background fluctuation (Atkins et al. 2000; Atkins et al. 2003)

MILAGRITO and GRB 970417a



- The VHE fluence must be at least an order of magnitude greater than sub-MeV fluence measured by BATSE
- A way out: if the actual peak of the SED may lie below the energy range of BATSE ?


Fig. 9.—Spectral energy distribution for GRB 970417a, showing a single power-law fit to the BATSE data, upper limits at 1 and 10 MeV from the EGRET TASC detector, and three possible spectral forms consistent with the Milagro observations.

Observing Transients with TeV Instruments

- All the major TeV instruments like **Veritas, H.E.S.S., HAWC, MAGIC, LHASSO,...** are pursuing intense programs for following transient alerts: GRBs, Gravitational wave sources, Neutrino ToO, Fast Radio Bursts, Novae, etc.
- While for a wide angle detector like HAWC the transient just needs to happen in their field of view, the narrow field of view IACTs need to possibly fast slew and track the alerted position
- All the IACTs optimized their operation for possibly fast slewing to the alerted position and starting observations
- Despite the really large number of followed GRBs by the above collaborations over many years, until recently only upper limits were reported

The 17m \emptyset MAGIC IACT project for VHE γ astrophysics at $E \sim 30$ GeV - 100 TeV

www.magic.mpp.mpg.de



Fast Motion of the MAGIC Telescopes



We devote this detection to the memory of Eckart Lorenz

- Those who had the privilege to work with him or simply had discussions with him knew his innovative spirit, charisma and optimism
- 25 years ago we were dreaming about measuring GRBs, Pulsars, AGN, ..., at $E > 10$ GeV
- With his unquenchable energy, brilliant know-how of materials and experimental methods in physics he was the driver behind to meet the preparations for the GRB observations with MAGIC

This snapshot recorded the very happy last minute in the life of Eckart; he became member of the Polish Academy of Sciences




MAGIC is Using a Fully-Automatic System for Following the Transient Alerts

- When obtaining an alert the software checks several criteria if the alert should be followed
- If yes, the telescopes start fast rotation and within ≤ 25 s slue to any given position in the sky and start tracking it
- While the telescopes are in the fast motion, the software closes the data-taking files of the previous observation, downloads the look-up tables for the aimed for elevation angle and by using the AMC system re-adjusts the individual mirror positions, adjusts the discriminator thresholds to the estimated brightness of the aimed position

MAGIC is Using a Fully-Automatic System for Following the Transient Alerts

- A special attention is paid to the safety issues because for saving precious seconds, there is no human intervention
- All these operations follow detailed protocols, which over many years were improved, tested and debugged
- For keeping the system and the current crew on the observational shift in a good shape, during observations we used to issue fake alerts to imaginary GRB positions; typically these were chosen to be one of the known intense gamma-ray emitters – this way later on one could check if the action was successful
- Of course the initiators of the fake alerts kept information confidential


MAGIC: 180 Astro-Physicists From 12 Countries




- Armenia** ICRA Net and Alikhanian Broth. Nat. Lab.
- Bulgaria** Sofia nuclear Physics Institute
- Brazil** CBPF Rio de Janeiro
- Croatia** Consortium (Zagreb, +...)
- Finland** Consortium (Tuorla, +...)
- Germany** DESY Zeuthen, U. Dortmund, MPI Munich, U. Würzburg
- Japan** Consortium (Kyoto, +...)
- Italy** INFN & U. Padova, INFN Pisa & U. Siena, INFN Como/Milano Bicocca, INFN Udine/Trieste & U. Udine, INAF (Consortium: Rome, +...)
- Poland** U. Lodz
- Spain** U. Barcelona, UAB Barcelona, IEEC-CSIC Barcelona, IFAE Barcelona, IAA Granada, IAC Tenerife, U. Complutense Madrid, CIEMAT Madrid
- Switzerland** ETH Zurich
- India** Kolkata

GRB follow-ups


- Since 2005 we observed 105 GRBs
- On average 8-10 GRB/year
- Afterglow observations since 2013
- 24 with < 100 s delay & stereo,
4 out of which with $z < 1.5$



Red-shift distribution




GRB 160821B



List of MAGIC GRBs observed under good technical and weather conditions with $z < 1$ and $T_{\text{delay}} < 1$ h

Event	redshift	T_{delay} (s)	Zenith angle (deg)	
GRB 061217	0.83	786.0	59.9	short
GRB 100816A	0.80	1439.0	26.0	short
GRB 160821B	0.16	24.0	34.0	short
GRB 190114C	0.42	58.0	55.8	long

MAGIC Hint From GRB 160821B



MAGIC capability to observe at the presence of **partial Moon** and at **large zenith angles** were of key importance for this detection

First time detection of a GRB at sub-TeV energies; MAGIC detects the GRB 190114C

ATel #12390; *Razmik Mirzoyan on behalf of the MAGIC Collaboration*

on 15 Jan 2019; 01:03 UT

Credential Certification: *Razmik Mirzoyan (Razmik.Mirzoyan@mpp.mpg.de)*


ATel issued 4h
after GRB onset

Subjects: Gamma Ray, >GeV, TeV, VHE, Request for Observations, Gamma-Ray Burst

Referred to by ATel #: 12395



The MAGIC telescopes performed a rapid follow-up observation of GRB 190114C (Gropp et al., GCN 23688; Tyurina et al., GCN 23690, de Ugarte Postigo et al., GCN 23692, Lipunov et al. GCN 23693, Selsing et al. GCN 23695). This observation was triggered by the Swift-BAT alert; we started observing at about 50s after Swift T0: 20:57:03.19. The MAGIC real-time analysis shows a significance >20 sigma in the first 20 min of observations (starting at T0+50s) for energies >300GeV. The relatively high detection threshold is due to the **large zenith angle of observations** (>60 degrees) and the **presence of partial Moon**. Given the brightness of the event, MAGIC will continue the observation of GRB 190114C until it is observable tonight and also in the next days. We strongly encourage follow-up observations by other instruments. The MAGIC contact persons for these observations are R. Mirzoyan (Razmik.Mirzoyan@mpp.mpg.de) and K. Noda (nodak@icrr.u-tokyo.ac.jp). MAGIC is a system of two 17m-diameter Imaging Atmospheric Cherenkov Telescopes located at the Observatorio Roque de los Muchachos on the Canary island La Palma, Spain, and designed to perform gamma-ray astronomy in the energy range from 50 GeV to

The most intense, purest signal in VHE γ -ray astrophysics: GRB190114C detection by MAGIC at $E \geq 200$ GeV



γ -rate from GRB190114C & Crab Nebula by MAGIC for the first ~ 100 s after start of data taking; during the first 30s the rate is ~ 1 kCrab

Please note the Y-axes scale difference of $\times 10$ for the GRB190114C




~5 Months After MAGIC GRB Detection @ CTA Symposium in Bologna on 8th of May 2019 H.E.S.S. Reported on GRB 180721B


The GRB180720B detection

- 10 hours after the Swift trigger
- Redshift $z = 0.653$ (ESO-VLT/X-shooter)

Details in Gal10a by Quentin Piel



2 hours taken between 25° and 40°
zenith angle



Between 18 days and 25 days
after the burst




Piel - Gamma-Ray Burst observations at Very High Energy with H.E.S.S. - ICRC 2019 9

H.E.S.S. Detection of Afterglow From GRB180720B

The GRB180720B detection

- Fermi-LAT detection up to 700 s after trigger with a photon index close to -2.0.
- Coincident optical and X-ray temporal decay
- H.E.S.S. detection until 440 GeV and same level as X-ray domain



The Original Goals of the MAGIC Project: Measure GRBs, Pulsars, Distant AGN above the threshold energy 10 GeV

MAGIC Design Study, 1998



MAGIC is a pioneering IACT
paving the road down to
a few 10's of GeV energies

Introduced multiple novel
techniques & technologies
into the field

- Fast slew for transients
- Active Mirror Control
- T° controlled camera
- Analog signal via optic fibers, ~2ns fast signals
- Carbon-fiber structure,...

After 15 years of successful operation of MAGIC we thought we are ready to measure the next best GRB above the lowest threshold

- With time we improved the performance of the MAGIC telescopes and learnt to measure γ -ray signals from sources at $E \geq 20\text{-}30$ GeV;
- Interested are invited to see the presentations by *Giovanni Ceribella, Gal9b “Crab Pulsar & Nebula”* and *Marcos López, Gal2g “Geminga pulsar”*, this conference
- We succeeded to measure the spectrum of a remote *FSRQ PKS1441* ($z=0.939$) at $E \geq 40$ GeV (*Ahnen, et al, ApJL 2015*)
- So we were preparing and training ourselves to measure γ -rays at lowest energies $\geq 20\text{-}30$ GeV from the next best GRB

The Surprise with the GRB 190114C


- But the reality looked quite different
- The burst of GRB 190114C happened at a **large zenith angle of $\sim 60^\circ$** and moreover, **the Moon was up**
- Both effects together increased the threshold energy to ~ 200 - 300 GeV
- Why GRB happened at high zenith angle ? Because the observation chance is proportional to the solid angle
- Why at the time when the Moon was up ? Because the partial Moon observations seriously increase the duty cycle of IACTs
- Despite the high threshold we were surprised to measure in a shortest time a truly gigantic signal, the most intense signal in the 30 years history of VHE gamma astronomy

MAGIC Partial Moon Observations Were Crucial for Discovering the GRB 190114C

- Hence from the beginning of the project MAGIC is regularly observing at the presence of partial Moon, at Dusk and Dawn
- These provide significantly longer observations, albeit at somewhat higher threshold energy
- The Moon observations were inherited already from the HEGRA CT1 operation; back in 1997 we observed Mrk-501 for 244 h dark time and 134 h at partial moonlight. The latter was crucial for finding quasi-periodic γ -ray emission from Mrk-501, see *Kranich et al., 1999*
- MAGIC & Moon observ.: *Albert et al (2007); Ahnen et al (2017)*
- During the GRB 190114C observations the partial Moon increased the anode currents of PMTs by 6-8 times – fully O.k.


GRB 190114C signal footprint

- Very strong signal, almost background-free
- Signal detected up to ~ 40 minutes
- Energy flux emitted @ sub-TeV comparable within factor of two to the one emitted in X-rays (between 60-2000s)




GRB 190114C light curves in the keV, GeV and TeV bands

MAGIC detected the afterglow phase of the GRB



Distribution of TeV-band γ rays in energy versus time for GRB 190114C


MAGIC detected an energetic component not yet seen in GRBs, different from synchrotron




Bottom curves show the expected max. photon energy $\varepsilon_{\text{syn,max}}$ of electron synchrotron radiation in the standard afterglow theory, for two extreme cases; isotropic-equivalent blast wave kinetic energy $E_{k,\text{aft}} = 3 \cdot 10^{55}$ erg
 dotted – $n = \text{const.}$
 dashed – $n \sim 1/R^2$

Multi-wavelength light curves of GRB190114C measured > 20 space-born and ground-based instruments


- MAGIC detected the afterglow phase of the GRB
- A significant share of GRB energy is emitted in the TeV energy range



Development of multi-wavelength spectra of GRB 190114C in time



Luminosity of GRB 190114C as compared to other long GRBs: it is not too special



Take away messages on the detection of GRB 190114C

- MAGIC detected an energetic component not yet seen in GRBs, different from synchrotron
- MAGIC clearly detected the afterglow phase of the GRB 190114C from redshift $z=0.42$
- The new component is energetic, extending $\gg 300$ GeV
- This is the brightest source ever detected by IACTs, the peak rate of gamma-rays ~ 0.1 kCrab
- The flux observed at T_0+80 s corresponds to apparent isotropic-equivalent luminosity $L_{\text{iso}} \sim 3 \times 10^{49}$ erg/s at $E \geq 0.3$ TeV, making this the most luminous source known at these energies

Take away messages on the detection of GRBs

- Most GRBs may possibly possess TeV emission components similar to GRB 190114C, which may be detectable as long as redshift is low and the observing conditions are suitable
- H.E.S.S. reports on γ -ray emission of up to 400 GeV from GRB 180720B at $z=0.653$ 10h after the burst – afterglow
- MAGIC has measured a 3σ hint from the *short* GRB 160821B
- This is just the beginning of the GRB era with IACTs