

The Correlated Spatial Structure of the Proton: Two-body densities as a framework for dynamical imaging

Zaki Panjsheeri,^{a,*} Joshua Bautista^a and Simonetta Liuti^a

^aUniversity of Virginia Physics Department,
382 McCormick Rd, Charlottesville, Virginia

E-mail: zap2nd@virginia.edu, rzp9zf@virginia.edu, sly4@virginia.edu

We present results on two-parton densities in coordinate space which capture a fuller dynamical picture of the proton's internal structure, including information on the relative position between quarks and gluons in the transverse plane. The connection of such two-body densities to observables, proceeds in QCD, via the definition of double generalized parton distributions (DGPDs).

25th International Spin Physics Symposium (SPIN 2023)
24-29 September 2023
Durham, NC, USA

*Speaker

1. Introduction

It is believed that the exclusive process of deeply virtual Compton scattering (DVCS) allows us to access generalized parton distributions (GPDs) [1–3] (we refer the reader to Refs.[4–6] for reviews on the subject). In turn, GPDs, through Fourier transformation, give spatial information on the charge and matter distributions of the quarks and gluons inside the nucleon [7]. The physical properties derived from the Fourier transforms of GPDs include the spatial distributions of each partonic component of the nucleon in the transverse plane with respect to the proton’s motion, as well as the orbital component of angular momentum. Notwithstanding the wealth of information provided by these quantities, to capture a fuller dynamical picture of the proton’s internal structure, information on the *relative* position between partons is crucial. In order to access information on the relative position of quarks and gluons, one needs to define the correlation functions yielding the two-body density distribution in the transverse plane. Connecting the two-body densities to observables, we found that the latter can be defined in QCD with generalized double parton distributions (GDPDs). Using GDPDs, we can address additional observables to describe quark and gluon dynamics, including their overlap probabilities. Such quantities can be extracted from experimental measurements of deeply virtual exclusive processes characterized by multi-particle final states.

2. From one body to two body density

We focus our description on the unpolarized quark inside an unpolarized nucleon described by the GPD, $H^q(X, \zeta = 0, t)$, where x is the quark longitudinal momentum fraction, $t = \Delta^2$ is the four-momentum transfer squared between the initial and final proton, and ζ , the skewness parameter is set to zero. Through Fourier transformation, one obtains the impact parameter-dependent parton distribution function (IPPDF) $\rho(X, \mathbf{b}_T)$, in terms of \mathbf{b}_T , the Fourier conjugate to the transverse momentum transfer, Δ_T [7, 8]. The spatial variables relevant for this problem are described in the left diagram in Figure 1, giving a partonic picture in coordinate space, as first observed using Light Cone (LC) kinematics in Ref.[9].




Figure 1: Spatial coordinates in the transverse plane for one-body distributions (left) and two-body distributions (right).

From the figure, one can see that two relevant variables describe the spatial configuration in a parton-spectator system, namely, $b = b_T$, the relevant transverse distance of the parton from the Center of Momentum (CoM), and R , the position of the nucleon’s CoM. To obtain such a picture

through GPDs, we take the correlation function represented in momentum space by Fig. 2,

$$W_{\Lambda, \Lambda'}^\Gamma = \int \frac{dz_{in}^-}{(2\pi)} \int \frac{dz_{out}^-}{(2\pi)^2} e^{i(k_{in} z_{in})} e^{-i(k_{out} z_{out})} \langle p', \Lambda' | \bar{\psi}(z_{out}) \Gamma \mathcal{U}(z_{in}, z_{out}) \psi(z_{in}) | p, \Lambda \rangle \Big|_{\substack{z_{in}^+(out)=0 \\ \mathbf{z}_{T, in(out)}=0}} \quad (1)$$

We represent the four-vectors in LC notation, $v \equiv (v^+, v^-, \mathbf{v}_T)$, $v^\pm = (v^0 \pm v^3)/\sqrt{2}$, $d^4z =$




Figure 2: Correlation function for the GPD in momentum space

$dz^- dz^+ d^2\mathbf{z}_T$; we evaluate the fields at equal LC time, $z^+ = 0$. $\psi(z_{in})$, $\bar{\psi}_{out}(z)$ are the quark fields; Γ is an operator between quark fields, *i.e.* either a specific gamma matrix, or a combination of gamma matrices; finally, $\mathcal{U}(z_{in}, z_{out})$, is the link ensuring gauge invariance. In what follows we choose the LC gauge for which $\mathcal{U}(z_{in}, z_{out})$ can be taken equal to one. The field operators product in Eq.(1) is evaluated between an incoming proton state, $| p, \Lambda \rangle$, of definite momentum, p , and helicity, Λ , and an outgoing proton state, $| p', \Lambda' \rangle$, with momentum, $p' = p - \Delta$, and helicity, Λ' . Taking $\Gamma = \gamma^+$, and Fourier transforming, one obtains ¹

$$\begin{aligned} W_{\Lambda, \Lambda'}^{\gamma^+} &= \int \frac{dz^-}{(2\pi)} e^{i(X-\zeta/2)p^+z^-} \langle p' = p - \Delta, \Lambda' | \bar{\psi}(0) \gamma^+ \psi(z^-) | p, \Lambda \rangle, \\ &= \frac{1}{2P^+} \left\{ H_q(X, \zeta, t) \bar{u}(p - \Delta, \Lambda') \gamma^+ u(p, \Lambda) + E_q(X, \zeta, t) \bar{u}(p - \Delta, \Lambda') \frac{\sigma^{i+} \Delta_i}{2M} u(p, \Lambda) \right\} \quad (2) \end{aligned}$$

Isolating the GPD H for an unpolarized quark inside an unpolarized proton at skewness ζ equal to 0, in momentum space the $H_q(X, 0, t)$ is off-diagonal in momentum,

$$H_q(X, 0, t) = \int d^2\mathbf{k}_{T, in} \phi^*(X, \mathbf{k}_{T, in} - \Delta) \phi(X, \mathbf{k}_{T, in}), \quad (3)$$

and with Fourier transformations of the vertex functions,

$$\phi(X, \mathbf{k}_{in}) = \frac{1}{(2\pi)^2} \int d^2\mathbf{z}_{in} e^{i\mathbf{k}_{in} \cdot \mathbf{z}_{in}} \tilde{\phi}(X, \mathbf{z}_{in}), \quad (4a)$$

$$\phi^*(X, \mathbf{k}_{out}) = \frac{1}{(2\pi)^2} \int d^2\mathbf{z}_{out} e^{-i\mathbf{k}_{out} \cdot \mathbf{z}_{out}} \tilde{\phi}^*(X, \mathbf{z}_{out}). \quad (4b)$$

the relationship between the GPD and the IPPDF emerges:

$$H_q(X, 0, t) = \int d^2\mathbf{b} e^{i\mathbf{b} \cdot \Delta} \tilde{\phi}^*(X, \mathbf{b}) \tilde{\phi}(X, \mathbf{b}) = \int d^2\mathbf{b} e^{i\mathbf{b} \cdot \Delta} \rho_q(X, \mathbf{b}) \quad (5a)$$

$$\rho_q(X, \mathbf{b}) = \int \frac{d^2\Delta}{(2\pi)^2} e^{-i\mathbf{b} \cdot \Delta} H_q(X, 0, t), \quad (5b)$$

¹The correlation function's parametrization in terms of GPDs also depends on the QCD evolution scale for the process, and Q^2 , that is omitted in these formulae for ease of presentation.

where we have defined the transverse space one-body diagonal density distribution. The index q refers to the quark flavor. A similar distribution is obtained for gluons.

In Figure 3 we show an example of a one-body density distribution, $\rho_g(X, b)$, evaluated for the gluon GPD parametrization from Refs.[10, 11], where H_g was constrained by its first moment in X evaluated in lattice QCD [12, 13]. Notice that in this case, the gluon distribution depends rather strongly on the scale for the process, Q^2 .




Figure 3: The Fourier transform $\rho_g(x, b)$ of the unpolarized gluon GPD H_g calculated as a function of x and Q^2 .




Figure 4: Two-body correlation function

Although the one-body density distributions already give rich spatial information about the proton's internal structure, as we show in Section 3, in order to study the dynamics of the proton's constituents, we must move from a single-body to a two-body picture, studying now double-parton correlations. We thus introduce the two-body correlation function, as in Figure 4, defined through

the following bilinear expression,

$$\begin{aligned}
 W_{\Lambda, \Lambda'}^{\Gamma} &= \int \frac{dz_{1,in}^- d\mathbf{z}_{1,T,in}}{(2\pi)^3} \frac{dz_{2,in}^- d\mathbf{z}_{2,T,in}}{(2\pi)^3} \int \frac{dz_{1,out}^- d\mathbf{z}_{1,T,out}}{(2\pi)^3} \frac{dz_{2,out}^- d\mathbf{z}_{2,T,out}}{(2\pi)^3} \\
 &\times e^{i(k_{1,in}z_{1,in} + k_{2,in}z_{2,in})} e^{-i(k_{1,out}z_{1,out} + k_{2,out}z_{2,out})} \\
 &\times \langle p', \Lambda' | \bar{\psi}(z_{1,out}) \Gamma \psi(z_{1,in}) \bar{\psi}(z_{2,out}) \Gamma \psi(z_{2,in}) | p, \Lambda \rangle \Big|_{z_1^+ = z_2^+ = 0}
 \end{aligned} \tag{6}$$

This definition involving quark fields is similar to the double parton distribution function (DPDF) introduced in [14], but with the initial and final proton states being different, ($p' \neq p$). Proceeding




Figure 5: Visualizing the two-body density as a function of b_1 and b_2 (Eq.9), for different values of x_2 , while keeping x_1 fixed.

similarly to the one-body case, we can now form the following combinations of the two different quark fields locations, $z_{i,in}$ and $z_{i,out}$, ($i = 1, 2$), respectively given by,

$$\begin{cases} z_i = z_{i,in} - z_{i,out} \\ b_i = \frac{1}{2}(z_{i,in} + z_{i,out}) \end{cases} \Rightarrow \begin{cases} z_{i,in} = b_i + \frac{z_i}{2} \\ z_{i,out} = b_i - \frac{z_i}{2} \end{cases}, \tag{7}$$

as well as the conjugate momenta to b_i and z_i ,

$$\begin{cases} \Delta_i = k_{i,in} - k_{i,out} \\ k_i = \frac{1}{2}(k_{i,in} + k_{i,out}) \end{cases} \Leftrightarrow \begin{cases} b_i \\ z_i \end{cases} \tag{8}$$

Following a procedure similar to the one-body case, we derived a formulation for the Fourier transform of the four-point correlation function (6), showing that it can be cast in a two-body density form, $\rho_2^{qq}(x_1, \mathbf{b}_1; x_2, \mathbf{b}_2)$, describing the probability of simultaneously finding quark 1 carrying a momentum fraction x_1 at location \mathbf{b}_1 and quark 2 carrying a momentum fraction x_2 at location \mathbf{b}_2 with respect to the center of momentum of the system. In the absence of correlations in the relative motion of the two quarks, the two-body transverse spatial distribution is described by,

$$\rho_2^{qq}(X_1, \mathbf{b}_1; X_2, \mathbf{b}_2) = \rho_q(X_1, \mathbf{b}_1)\rho_q(X_2, \mathbf{b}_2) \quad (9)$$

where ρ is the diagonal one-body density defined in Eq.(5). Distributions with parton configurations other than qq , *e.g.* qg , or gg , can be written in a similar way, starting from the appropriate two-body correlation function definition.

A visualization of the two-body density distribution, is provided in Fig. 5 where in the upper panel we show a 2D rendition of the gluon-gluon distribution obtained by plotting the quantity, $\rho_2^{gg}(X_1, \mathbf{b}_1; X_2, \mathbf{b}_2)$ at the scale, $Q^2 = 5 \text{ GeV}^2$, for $X_1 = 10^{-3}$, and fixed \mathbf{b}_2 , varying X_2 and \mathbf{b}_1 ; in the lower panels, we show 3D versions obtained at $x_1 = 0.01$ (top) and $X_1 = 10^{-3}$, plotted in the $\mathbf{b}_1, \mathbf{b}_2$ for various values of X_2 .

3. Numerical results/Observables

As described in the previous section, since the Fourier transform of the GPD H for zero skewness gives the IPPDF, we obtain the single-particle spatial density of the parton's charge inside the proton (see *e.g.* Fig. 3 for the gluon distribution). From such single-body densities, we can compute the average impact parameter squared, $\langle \mathbf{b}^2 \rangle$, or the average transverse radius, as,

$$\langle \mathbf{b}_{q,g}^2(X) \rangle = \frac{1}{N_{q,g}} \int d^2\mathbf{b} \mathbf{b}^2 \rho_{q,g}(X, \mathbf{b}), , g \quad (10)$$

where,

$$N_{q,g} = \int d^2\mathbf{b} \rho_{q,g}(X, \mathbf{b}) = H_{q,g}(X, 0, 0). \quad (11)$$

This normalization encapsulates the baryon number, putting the partonic distributions on equal footing to make a faithful comparison. The average radii are shown in Figure 6 where on the *lhs* we plotted the valence u and we d quark geometric radii compared to the gluon one, at $Q^2 = 5 \text{ GeV}^2$; on the *rhs* the behavior of the gluon radius with Q^2 is shown. All calculations were performed using the parametrization from Refs.[10, 11]. One can see that the gluon radius is much smaller than the quark one, consistent with the geometric representation of baryon junctions, as proposed in Ref.[15] and recently investigated in, *e.g.*, Ref.[16]. Furthermore, we find that, while the radii for all q and g components show a small variation with Q^2 at larger values of $X > 0.01$, at small X , this dependence is substantial, and it should, therefore be more easily detectable in experiment.

From the vectors locating the partons 1 and 2 positions in the transverse plane, \mathbf{b}_1 and \mathbf{b}_2 , respectively, we define the relative distance between them, \mathbf{r} , and the distance of the center of mass of the two partons from the origin, \mathbf{R}_{12} ,

$$\mathbf{r} = \mathbf{b}_1 - \mathbf{b}_2 \quad (12)$$

$$\mathbf{R}_{12} = \frac{\mathbf{b}_1 + \mathbf{b}_2}{2} \quad (13)$$




Figure 6: The average radii for u , d (valence) and g (gluon) distributions (left) provide a quantitative method for extracting spatial information through even the one-body density. We can also study the Q^2 -dependence of these average radii through perturbative evolution, as shown for the gluons (right).

Defining the root mean squared of their expectation values, one obtains the square average relative distance and average center of mass of the two partons,

$$\langle \mathbf{r}^2 \rangle (X_1, X_2) = \frac{1}{\mathcal{N}} \int \int d^2 \mathbf{r} d^2 \mathbf{R}_{12} |\mathbf{r}^2| \rho_2 \left(X_1, \mathbf{R}_{12} + \frac{\mathbf{r}}{2}; X_2, \mathbf{R}_{12} - \frac{\mathbf{r}}{2} \right) \quad (14)$$

$$\langle \mathbf{R}_{12}^2 \rangle (X_1, X_2) = \frac{1}{\mathcal{N}} \int \int d^2 \mathbf{r} r^2 \mathbf{R}_{12} |\mathbf{R}_{12}^2| \rho_2 \left(X_1, \mathbf{R}_{12} + \frac{\mathbf{r}}{2}; X_2, \mathbf{R}_{12} - \frac{\mathbf{r}}{2} \right) \quad (15)$$

$$\mathcal{N} = \int \int d^2 \mathbf{r} r^2 \mathbf{R}_{12} \rho_2 \left(X_1, \mathbf{R}_{12} + \frac{\mathbf{r}}{2}; X_2, \mathbf{R}_{12} - \frac{\mathbf{r}}{2} \right) \quad (16)$$

The numerical results obtained using our parametrization for the average relative distance, given in




Figure 7: Average relative distance between the u quark and gluon, the d quark and gluon, and two gluon fields inside the proton plotted vs. $X = X_1 = X_2$, at $Q^2 = 10 \text{ GeV}^2$.

Fig. 7, for the specific case of $X_1 = X_2$, correlate with the results for the single parton distributions, in that the distance between a quark of flavor u , d and a gluon field is always larger than the distance of two gluon fields, revealing that gluonic configurations tend to be more compact.

Through the two-body densities we can now address quantitatively the question of whether the u , quarks and gluons will overlap forming localized “hot spots,” as suggested in an event-by-event

analysis in Refs. [17, 18], or whether the gluon field surrounds uniformly the valence quarks. The geometric average two-parton overlap probability in the transverse plane can be defined as,

$$\langle A(X_1, X_2) \rangle = \frac{1}{\mathcal{N}} \int d^2\mathbf{b}_1 d^2\mathbf{b}_2 \rho_2^{ij}(X_1, \mathbf{b}_1; X_2, \mathbf{b}_2) A(|\mathbf{b}_1 - \mathbf{b}_2|),$$

where $A(r = |\mathbf{b}_1 - \mathbf{b}_2|)$ is taken as the geometric area overlap of the two azimuthally symmetric single parton distributions measured in units of an average radius $a = (\langle b_1 \rangle^2 + \langle b_2 \rangle^2)^{1/2} / \sqrt{2}$,

$$A(r) = a^2 \pi - \frac{r - \delta}{2} (\alpha_1 + \alpha_2) - a_1^2 \arctan \frac{r - \delta}{2\alpha_1} - a_2^2 \arctan \frac{r - \delta}{2\alpha_2} \quad (17)$$

with,

$$\alpha_i = \sqrt{a_i^2 - \frac{(r - \delta)^2}{4}}, \quad i = 1, 2 \quad (18)$$

and $\delta = a_1 - a_2$.

Performing a change of variables, the average two-parton overlap probability is given as,

$$\langle A(X_1, X_2) \rangle = \frac{1}{\mathcal{N}} \int \int d^2\mathbf{r} d^2\mathbf{R}_{12} A(r) \rho_2^{qg} \left(X_1, \mathbf{R}_{12} + \frac{\mathbf{r}}{2}; X_2, \mathbf{R}_{12} - \frac{\mathbf{r}}{2} \right). \quad (19)$$

Numerical results for the overlap probability are given in Figure 8 for the u quark and gluon case. The figure shows $A(X_1, X_2)$, Eq.(19), plotted vs. X , with $X = X_1 = X_2$, and $Q^2 = 10 \text{ GeV}^2$, divided by the maximum overlap surface. Putting together results from Figs.6, 7, and 8, we can see that while all parton types tend to be more broadly distributed at low X (Fig.6), thus also spanning broadly distributed relative distances, their overlap probability also increase (Fig.8). This picture complements the recent ‘‘BNL approach’’, describing ‘‘geometrical fluctuations’’, or event-by-event fluctuations in the proton wave function where the three constituent quarks emit small- x gluons around them, which form ‘‘hot spots’’ at random locations in the transverse plane. [17–20]




Figure 8: The overlap probability of the u quark and gluon distributions in the proton plotted vs $X = X_1 = X_2$ for $Q^2 = 10 \text{ GeV}^2$.

4. Conclusions

We presented results on our study of two-body parton distributions as a means of obtaining information on the relative positions of quarks and gluons inside the proton. While one-body

distributions are limited to capturing average static properties of the proton's internal structure, such as the radius covered by various parton distributions, the proton is a complex, multi-body environment, and it is, therefore, more properly imaged through multi-body distributions. As a first step towards the general problem of imaging this multi-body system, and to provide tools to address multiparton correlations beyond the one-body formalism, we studied the geometric correlations between the u and d quark distributions and gluons.

In Refs.[17, 18] a connection was provided between the two approaches, "GPDs" on one side, and "geometrical fluctuations" on the other. In particular, the dipole-based description of *coherent* diffractive J/ψ production off a proton was connected to the correlation functions for GPDs, within the collinear framework of QCD. However, the more interesting channel of *incoherent* diffraction, described in terms of double dipole diffraction and leading to small- x gluon fluctuations, or event-by-event, color density geometry driven fluctuations has remained a puzzle in that it could not be rendered, so far, within standard QCD approaches. By introducing two-body correlations functions, we provide a framework to investigate these scenarios and to allow us to study relative spatial configurations of quarks and gluons inside the proton and the atomic nuclei.

Acknowledgments

This research is funded by DOE grant DE-SC0016286.

References

- [1] Xiang-Dong Ji. Gauge-Invariant Decomposition of Nucleon Spin. *Phys. Rev. Lett.*, 78:610–613, 1997.
- [2] Xiang-Dong Ji. Deeply virtual Compton scattering. *Phys. Rev.*, D55:7114–7125, 1997.
- [3] A. V. Radyushkin. Nonforward parton distributions. *Phys. Rev.*, D56:5524–5557, 1997.
- [4] M. Diehl. Generalized parton distributions with helicity flip. *Eur.Phys.J.*, C19:485–492, 2001.
- [5] A.V. Belitsky and A.V. Radyushkin. Unraveling hadron structure with generalized parton distributions. *Phys.Rept.*, 418:1–387, 2005.
- [6] Kresimir Kumericki, Simonetta Liuti, and Herve Moutarde. GPD phenomenology and DVCS fitting - Entering the high-precision era. 2016.
- [7] Matthias Burkardt. Impact parameter dependent parton distributions and off forward parton distributions for zeta $\rightarrow 0$. *Phys. Rev. D*, 62:071503, 2000. [Erratum: *Phys.Rev.D* 66, 119903 (2002)].
- [8] M. Diehl. Generalized parton distributions in impact parameter space. *Eur. Phys. J. C*, 25:223–232, 2002. [Erratum: *Eur.Phys.J.C* 31, 277–278 (2003)].
- [9] Davison E. Soper. The Parton Model and the Bethe-Salpeter Wave Function. *Phys. Rev.*, D15:1141, 1977.

- [10] Brandon Kriesten, Philip Velie, Emma Yeats, Fernanda Yopez Lopez, and Simonetta Liuti. Parametrization of quark and gluon generalized parton distributions in a dynamical framework. *Phys. Rev. D*, 105(5):056022, 2022.
- [11] Gary R. Goldstein, J. Osvaldo Gonzalez Hernandez, and Simonetta Liuti. Flexible Parametrization of Generalized Parton Distributions from Deeply Virtual Compton Scattering Observables. *Phys.Rev.*, D84:034007, 2011.
- [12] P. E. Shanahan and W. Detmold. Gluon gravitational form factors of the nucleon and the pion from lattice QCD. *Phys. Rev. D*, 99(1):014511, 2019.
- [13] Dimitra A. Pefkou, Daniel C. Hackett, and Phiala E. Shanahan. Gluon gravitational structure of hadrons of different spin. *Phys. Rev. D*, 105(5):054509, 2022.
- [14] Markus Diehl, Daniel Ostermeier, and Andreas Schafer. Elements of a theory for multiparton interactions in QCD. *JHEP*, 03:089, 2012. [Erratum: JHEP 03, 001 (2016)].
- [15] D. Kharzeev. Can gluons trace baryon number? *Physics Letters B*, 378(1–4):238–246, June 1996.
- [16] R. Terra and F. S. Navarra. Charmonium production in high multiplicity pp collisions and the structure of the proton. *Phys. Rev. D*, 108:054002, Sep 2023.
- [17] Heikki Mäntysaari. Review of proton and nuclear shape fluctuations at high energy. *Rept. Prog. Phys.*, 83(8):082201, 2020.
- [18] Heikki Mäntysaari, Kaushik Roy, Farid Salazar, and Björn Schenke. Gluon imaging using azimuthal correlations in diffractive scattering at the Electron-Ion Collider. *Phys. Rev. D*, 103(9):094026, 2021.
- [19] Heikki Mäntysaari and Björn Schenke. Accessing the gluonic structure of light nuclei at a future electron-ion collider. *Phys. Rev. C*, 101(1):015203, 2020.
- [20] Adrian Dumitru, Heikki Mäntysaari, and Risto Paatelainen. Color charge correlations in the proton at NLO: Beyond geometry based intuition. *Phys. Lett. B*, 820:136560, 2021.