

Extracting the distribution amplitude of pseudoscalar mesons using the HOPE method

Alex Chang,^{*a*} William Detmold,^{*b*} Anthony V. Grebe,^{*c*} Issaku Kanamori,^{*d*} C.-J. David Lin,^{*a,e*} Robert J. Perry^{*b,f,**} and Yong Zhao^{*g*} ^aInstitute of Physics, National Yang Ming Chiao Tung University, 1001 Ta-Hsueh Road, Hsinchu 30010, Taiwan ^bCenter for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA ^c Fermi National Accelerator Laboratory, Batavia, IL 60502, USA ^dRIKEN Center for Computational Science, Kobe 650-0047, Japan ^eCentre for High Energy Physics, Chung-Yuan Christian University, Chung-Li, 32032, Taiwan ^f Departament de Física Quàntica i Astrofísica and Institut de Ciències del Cosmos, Universitat de Barcelona, Martí Franquès 1, E08028, Spain ^gPhysics Division, Argonne National Laboratory, Lemont, IL 60439, USA *E-mail:* s44930e0@gmail.com, wdetmold@mit.edu, agrebe@mit.edu, kanamori-i@riken.jp, dlin@nycu.edu.tw, perryrobertjames@gmail.com, yong.zhao@anl.gov

The pseudoscalar meson light-cone distribution amplitudes (LCDAs) are essential nonperturbative inputs for a range of high-energy exclusive processes in quantum chromodynamics. In this proceedings, progress towards a determination of the low Mellin moments of the pion and kaon LCDAs by the HOPE Collaboration is reported.

The 41st International Symposium on Lattice Field Theory (LATTICE2024) 28 July - 3 August 2024 Liverpool, UK

*Speaker

[©] Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Introduction

The light-cone distribution amplitude (LCDA) is a non-perturbative object of interest for a range of high-energy exclusive processes in quantum chromodynamics (QCD) [1]. It can be defined in terms of the matrix element,

$$\langle 0|\overline{\psi}(z)\gamma^{\mu}\gamma_{5}W[z,-z]\psi(-z)|M^{+}(\vec{p})\rangle = ip^{\mu}f_{M}\int_{-1}^{1}d\xi \ \mathrm{e}^{-i\,\xi\,p\cdot z}\phi_{M}(\xi,\mu),\tag{1}$$

where $\mathcal{W}[-z, z]$ is a light-like $(z^2 = 0)$ Wilson line and $\phi_M(\xi, \mu)$ is the LCDA for the meson M. In the above equation, f_M , \vec{p} , and p_{μ} are the decay constant, three-momentum, and the four-momentum of the meson, respectively.

Despite the clear phenomenological value in computing this quantity from first principles, the presence of a light-like separation in the definition of the relevant operator prohibits a direct calculation using Euclidean formulations of lattice QCD (LQCD). Historically, the solution has been to employ an operator product expansion (OPE) to relate this operator to a sum of local operators, which can be calculated using LQCD [2, 3]. The meson-to-vacuum matrix elements of these local operators can be shown to be related to the Mellin moments of the LCDA,

$$\langle 0|\overline{\psi}\gamma^{\{\mu_1}\gamma_5(i\overset{\leftrightarrow}{D}{}^{\mu_2})\dots(i\overset{\leftrightarrow}{D}{}^{\mu_n\}})\psi|M^+(\vec{p})\rangle = f_M\,\langle\xi^n\rangle_M\,(\mu)[p^{\mu_1}p^{\mu_2}\dots p^{\mu_n} - \mathrm{tr}],\tag{2}$$

where

$$\langle \xi^n \rangle_M(\mu) = \int_{-1}^1 d\xi \, \xi^n \phi_M(\xi,\mu) \tag{3}$$

are the Mellin moments. This approach has been limited to the lowest few moments because the higher spin lattice operators mix with lower dimensional operators and the mixing coefficients contain power divergences. More recently, other approaches have been proposed [4-14].

2. The HOPE method

The heavy-quark operator product expansion (HOPE) is an alternative approach [6, 15] to extracting information about the Mellin moments of the LCDA. It is based on studying the hadronic matrix element

$$R_{M}^{\mu\nu}(t_{-},\vec{p},\vec{q}) = \int d^{3}\vec{z} \, e^{i\vec{q}\cdot\vec{z}} \, \langle 0|T\{J_{A}^{\mu}(t_{-}/2,\vec{z}/2)J_{A}^{\nu}(-t_{-}/2,-\vec{z}/2)\}|M(\vec{p})\rangle \,, \tag{4}$$

where

$$J^{\mu}_{A} = \overline{\Psi}\gamma^{\mu}\gamma^{5}\psi + \overline{\psi}\gamma^{\mu}\gamma^{5}\Psi \tag{5}$$

is an axial-vector current involving a light or strange quark ψ and a fictitious valence heavy quark Ψ with mass m_{Ψ} . The momentum-space hadronic matrix element is given by

$$V_M^{\mu\nu}(p,q) = \int dt_- e^{iq_4t_-} R_M^{\mu\nu}(t_-,\vec{p},\vec{q}).$$
(6)

The matrix element of this two-current operator can be written in terms of local operators via an application of the heavy-quark operator product expansion. To one-loop it can be written as [15]

$$V_M^{\mu\nu}(q,p) = -\frac{2if_\pi \epsilon^{\mu\nu\rho\sigma} q_\rho p_\sigma}{\tilde{Q}^2} \sum_{n=0,\text{even}}^{\infty} \mathcal{F}_n(\tilde{Q}^2,\mu,\tilde{\omega},m_\Psi)\phi_{M,n}(\mu) + \text{higher-twist terms}, \quad (7)$$

where $\phi_{M,n}(\mu)$ are the Gegenbauer moments, μ is the renormalization scale, and \mathcal{F}_n are coefficients that can be computed in QCD perturbation theory and can be expressed as functions of the kinematic variables,

$$\tilde{Q}^2 = Q^2 + m_{\Psi}^2, \quad \tilde{\omega} = \frac{2p \cdot q}{\tilde{Q}^2}, \quad (8)$$

with $Q^2 = -q^2$. To zeroth order in the coupling, the above expression can be equivalently written as

$$V_M^{\mu\nu}(q,p) = -\frac{2if_\pi \epsilon^{\mu\nu\rho\sigma} q_\rho p_\sigma}{\tilde{Q}^2} \sum_{n=0}^{\infty} \left(\frac{\tilde{\omega}}{2}\right)^n \langle \xi^n \rangle_M + \text{higher-twist terms}, \qquad (9)$$

where the Gegenbauer and Mellin moments of the LCDA are related via

$$\phi_{0,M} = \langle \xi^0 \rangle_M = 1, \quad \phi_{1,M} = \frac{5}{3} \langle \xi^1 \rangle_M, \quad \phi_{2,M} = \frac{7}{12} \left(5 \langle \xi^2 \rangle_M - \langle \xi^0 \rangle_M \right), \tag{10}$$

$$\phi_{3,M} = \frac{3}{4} \left(7 \left\langle \xi^3 \right\rangle - 3 \left\langle \xi^1 \right\rangle \right), \quad \phi_{4,M} = \frac{11}{24} \left(21 \left\langle \xi^4 \right\rangle_M - 14 \left\langle \xi^2 \right\rangle_M + \left\langle \xi^0 \right\rangle_M \right), \quad \dots$$

3. Numerical Strategy

In practice, the HOPE strategy involves computing correlation functions using LQCD which provide access to the hadronic matrix element defined by Eq. (4). By fitting the numerical data to the HOPE formula given by Eq. (7), information about the Gegenbauer moments of the meson LCDA can be determined. From the relations given in Eq. (10), this information can be converted into information about the Mellin moments. The hadronic matrix element can be determined from the correlation functions,

$$C_{ij}(t,\vec{p}) = \int d^3x e^{i\vec{p}\cdot\vec{x}} \langle 0|O_i(t,\vec{x})O_j^{\dagger}(0,\vec{0})|0\rangle , \qquad (11)$$

$$C_{j}^{\mu\nu}(t_{e},t_{m},\vec{p}_{e},\vec{p}_{m}) = \int d^{3}x_{e}d^{3}x_{m}e^{i\vec{p}_{e}\cdot\vec{x}_{e}}e^{i\vec{p}_{m}\cdot\vec{x}_{m}} \langle 0|J_{A}^{\mu}(t_{e},\vec{x}_{e})J_{A}^{\nu}(t_{m},\vec{x}_{m})O_{j}^{\dagger}(0,\vec{0})|0\rangle , \quad (12)$$

where $O_i(t, \vec{x})$ is an interpolating operator for the meson. In the studies covered in this proceedings, two interpolating operators, $O_1 = \overline{\psi}\gamma_5\psi$ and $O_2 = \overline{\psi}\gamma_4\gamma_5\psi$, are employed. The optimal linear combination which maximizes the overlap of the operator onto the ground state is constructed by solving the generalized eigenvalue problem [16, 17],

$$\sum_{j} C_{ij}(t_{\text{ref}}, \vec{p}) v_{n,j}(t_{\text{ref}}, t_0) = \lambda_n(t_{\text{ref}}, t_0) \sum_{j} C_{ij}(t_0, \vec{p}) v_{n,j}(t_{\text{ref}}, t_0).$$
(13)

The eigenvector $v_{n,j}(t_{ref}, t_0)$ defines the optimized interpolating operator for the *n*th state. Thus the optimized interpolating operator for the ground state is

$$O_M(t, \vec{x}) = \sum_j v_{0,j}(t_{\text{ref}}, t_0) O_j(t, \vec{x}).$$
(14)

This optimized interpolating operator is used self-consistently in the definitions of the two- and three-point correlation functions. In terms of this optimized interpolating operator for the pion, the two-point correlation function $C_{MM}(t, \vec{p})$ admits the spectral decomposition

$$C_{MM}(t,\vec{p}) = \sum_{n} \frac{|Z_{n,M}(\vec{p})|^2}{2E_n} e^{-E_n(\vec{p})t}$$
(15)



Figure 1: Fitting one-loop HOPE expression to the t_{-} -even and t_{-} -odd components of the hadronic matrix element computed using LQCD.

where $Z_{n,M}(\vec{p}) = \langle 0|O_M(0,\vec{0})|n,\vec{p}\rangle$.

The three-point correlation function C_M can be written as

$$C_{M}^{\mu\nu}(t_{e}, t_{m}, \vec{p}_{e}, \vec{p}_{m}) = R_{M}^{\mu\nu}(t_{-}, \vec{p}, \vec{q}) \frac{Z_{0,M}^{*}(\vec{p})}{2E_{0}(\vec{p})} e^{-E_{0}(\vec{p})t_{+}/2} + \text{excited states},$$
(16)

where $t_+ = t_e + t_m$, $t_- = t_e - t_m$, $\vec{p} = \vec{p}_e + \vec{p}_m$ and $\vec{q} = (\vec{p}_e - \vec{p}_m)/2$. By constructing the ratio

$$\mathcal{R}_{M}^{\mu\nu}(t_{-},t_{+},\vec{p},\vec{q}) = \frac{C_{M}(t_{e},t_{m},\vec{p}_{e},\vec{p}_{m})}{\frac{Z_{0,M}(\vec{p})}{2E_{0}(\vec{p})}e^{-E_{0}(\vec{p})t_{+}/2}},$$
(17)

and studying the large t_+ limit, it is possible to show that

$$\lim_{t_{+} \to \infty} \mathcal{R}_{M}^{\mu\nu}(t_{-}, t_{+}, \vec{p}, \vec{q}) = R_{M}^{\mu\nu}(t_{-}, \vec{p}, \vec{q}).$$
(18)

which may be fit to the HOPE formula to obtain information about the low Mellin moments of the meson LCDA.

4. Results

4.1 Determination of fourth Mellin moment of pion in the quenched approximation

In Ref. [18], the second moment of the pion LCDA was determined using the HOPE method. However, the pion momentum was insufficient to study higher moments of the pion LCDA. The following analysis builds upon that work by studying the hadronic matrix element $R^{\mu\nu}(t, \vec{p}, \vec{q})$ at larger pion three-momenta. In this study, the pion state is taken to have $|\vec{p}|/(2\pi/L) = 2$ units of momentum. The hadronic matrix element is computed on four ensembles generated in the quenched approximation for a range of heavy quark masses. The simulation parameters for these ensembles have previously been reported in Ref. [18].

In this study the ratio $\mathcal{R}^{\mu\nu}_{\pi}(t_-, t_+, \vec{p}, \vec{q})$ is fit to the one-loop HOPE formula evaluated at a renormalization scale of $\mu = 2$ GeV. An example of this fit is shown in Fig. 1. Following this, data is extrapolated to the continuum, twist-two limit using the extrapolation formula,

$$X(a, m_{\Psi}) = X_0 + \frac{A}{m_{\Psi}} + Ba^2 + Ca^2 m_{\Psi} + Da^2 m_{\Psi}^2,$$
(19)





Figure 2: Performing continuum, twist-2 extrapolation of fitted $\langle \xi^2 \rangle$.

Name	$(L/a)^3 \times T/a$	β	<i>a</i> (fm)	<i>K</i> light	K _{strange}	m_{π} (MeV)	m_K (MeV)
B451	$32^{3} \times 64$	3.46	0.075	0.136981	0.136409	422	577
N305	$48^3 \times 128$	3.7	0.049	0.137025	0.136676	428	584
H107	$32^{3} \times 96$	3.4	0.085	0.136946	0.136203	368	550
B452	$32^{3} \times 64$	3.46	0.075	0.137046	0.136378	352	548
N204	$48^3 \times 128$	3.55	0.064	0.137112	0.136575	353	549
N304	$48^3 \times 128$	3.7	0.049	0.137079	0.136665	353	558
N450	$48^{3} \times 128$	3.46	0.075	0.137099	0.136353	287	528

Table 1: Lattice details for dynamical ensembles employed in calculation of second Mellin moment of the pion and low Mellin moments of the kaon. Further details about these ensembles can be found in Ref. [20]

where $X \in \{f_{\pi}, \langle \xi^2 \rangle, \langle \xi^4 \rangle\}$. The form of this extrapolation is inspired by power-counting arguments for the leading higher-twist and lattice artifacts. In order to estimate the systematic uncertainty stemming from the use of this parameterization, the model averaging prescription proposed in Ref. [19] is employed. The resulting continuum, twist-two extrapolations of the second and fourth Mellin moments are presented in Fig. 2.

4.2 Progress in determination of second Mellin moment of pion

Previous work [18] focused on a proof-of-principle determination of the second moment of the pion LCDA. Due to the exploratory nature of that work, the quenched approximation was employed. As a result, the determined second Mellin moment of the LCDA suffers from a uncontrolled systematic error. Therefore, the HOPE Collaboration is currently repeating this calculation using dynamical ensembles generated by the CLS Collaboration [20]. Details of the dynamical ensembles employed in this work are given in Table 1. Using this set of ensembles, it will be possible to perform an extrapolation to the physical point in addition to the continuum, twist-two extrapolation performed in previous HOPE studies.

Except in several small ways, this analysis mirrors the analysis of the fourth moment and the previous quenched analysis of the second moment [18]. A preliminary study of excited state



Figure 3: (Left) Studying excited state contamination in the extracted second Mellin moment of the pion LCDA. (Right) Continuum, twist-two extrapolation of the second moment of the pion LCDA using ensembles with a pion mass of approximately $m_{\pi} \approx 350$ MeV.

contamination in the extracted second Mellin moment is presented in Fig. 3. After extrapolating excited state contamination, the resulting values of the second moment on the ensembles considered here is shown in Fig. 3, along with the preliminary continuum, twist-two extrapolation. This extrapolation employs the same *ansatz* proposed in Eq. (19). It is important to note that this analysis only considers ensembles with approximately equivalent pion masses.

4.3 Progress in determination of low moments of kaon LCDA

Exploratory work has commenced on the determination of the low moments of the kaon LCDA using the HOPE method. Due to the presence of a strange quark, both the odd and even Mellin moments are non-zero. In order to assess the sensitivity of the hadronic matrix element to the presence of the low Mellin moments it is useful to note that in momentum space, the linear combinations

$$V_{K,\text{even}}^{\mu\nu}(p,q) = \frac{1}{2} \left[V_K^{\mu\nu}(p,q) - V_K^{\mu\nu}(p,-q) \right],$$
(20)

$$V_{K,\text{odd}}^{\mu\nu}(p,q) = \frac{1}{2} \bigg[V_K^{\mu\nu}(p,q) + V_K^{\mu\nu}(p,-q) \bigg],$$
(21)

are sensitive to the even and odd Mellin moments, respectively. That is, at tree-level,

$$V_{K,\text{even}}^{\mu\nu}(p,q) = -\frac{2if_{\pi}\epsilon^{\mu\nu\rho\sigma}q_{\rho}p_{\sigma}}{\tilde{Q}^2} \sum_{n=0,\text{even}}^{\infty} \left(\frac{\tilde{\omega}}{2}\right)^n \langle \xi^n \rangle_M + \text{higher-twist terms}, \qquad (22)$$

$$V_{K,\text{odd}}^{\mu\nu}(p,q) = -\frac{2if_{\pi}\epsilon^{\mu\nu\rho\sigma}q_{\rho}p_{\sigma}}{\tilde{Q}^2} \sum_{n=0,\text{odd}}^{\infty} \left(\frac{\tilde{\omega}}{2}\right)^n \langle \xi^n \rangle_M + \text{higher-twist terms}, \qquad (23)$$

In addition to this linear combination, it is possible to further separate the hadronic matrix element into real and imaginary parts, as previously employed in studies of the pion LCDA [18]. The combination of these two approaches make it possible to isolate $\langle \xi^1 \rangle_K$ and $\langle \xi^2 \rangle_K$ in $\text{Im}V_{\text{odd}}^{\mu\nu}(p,q)$ and $\text{Re}V_{\text{even}}^{\mu\nu}(p,q)$, respectively. These amplitudes have been calculated using the ensembles listed in Table 1 and are shown in Fig. 4. The clear non-zero signal for each of these amplitudes is encouraging for a determination of these two Mellin moments using the HOPE method.





Figure 4: Real and imaginary components of the *q*-even and *q*-odd amplitudes which are sensitive to the even and odd Mellin moments of the Kaon LCDA.

5. Conclusions

The LCDA is the non-perturbative object of interest for a range of high-energy exclusive processes in QCD. In this proceedings, recent progress by the HOPE calculation in computing the low Mellin moments of the pion and the kaon LCDA are presented.

Acknowledgements

WD and RJP are supported in part by the U.S. Department of Energy, Office of Science under grant Contract Number DE-SC0011090 and by the SciDAC5 award DE-SC0023116. WD is additionally supported by the National Science Foundation under Cooperative Agreement PHY-2019786 (The NSF AI Institute for Artificial Intelligence and Fundamental Interactions, http://iaifi.org/). RJP has been supported by the projects CEX2019-000918-M (Unidad de Excelencia "Maria de Maeztu"), PID2020-118758GB-I00, financed by MICIU/AEI/10.13039/501100011033/ and FEDER, UE, as well as by the EU STRONG-2020 project, under the program H2020-INFRAIA-2018-1 Grant Agreement No. 824093. AC and CJDL are supported by 112-2112-M-A49-021-MY3 and 113-2123-M-A49-001-SVP. The authors thank ASRock Rack Inc. for their support of the construction of an Intel Knights Landing cluster at National Yang Ming Chiao Tung University, where the numerical calculations were performed. Help from Balint Joo in tuning Chroma is acknowledged. The authors thankfully acknowledge the computer resources at MareNostrum and the technical

support provided by BSC (RES-FI-2023-1-0030). The authors gratefully acknowledge the support of ASGC (Academia Sinica Grid Computing Center, AS-CFII-114-A11, NSTC(NSTC 113-2740-M-001-007) for provision of computational resources. This work utilized Supercomputer Fugaku in RIKEN Center for Computational Science through the HPIC project hp220312 and hp230466. We used Bridge++ code set [21–23]. We thank Enno Scholz, Gunnar Bali, and the CLS Collaboration for generously sharing the ensembles used in the dynamical calculation in this work. This document was prepared using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, Office of High Energy Physics HEP User Facility. Fermilab is managed by Fermi Forward Discovery Group, LLC, acting under Contract No. 89243024CSC000002. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics through Contract No. DE-AC02-06CH11357.

References

- [1] G.P. Lepage and S.J. Brodsky, *Exclusive Processes in Perturbative Quantum Chromodynamics*, *Phys. Rev. D* 22 (1980) 2157.
- [2] A.S. Kronfeld and D.M. Photiadis, *Phenomenology on the Lattice: Composite Operators in Lattice Gauge Theory*, *Phys. Rev. D* **31** (1985) 2939.
- [3] G. Martinelli and C.T. Sachrajda, A Lattice Calculation of the Second Moment of the Pion's Distribution Amplitude, Phys. Lett. B 190 (1987) 151.
- [4] U. Aglietti, M. Ciuchini, G. Corbo, E. Franco, G. Martinelli and L. Silvestrini, *Model independent determination of the light cone wave functions for exclusive processes*, *Phys. Lett. B* 441 (1998) 371 [hep-ph/9806277].
- [5] K.-F. Liu, Parton degrees of freedom from the path integral formalism, Phys. Rev. D 62 (2000) 074501 [hep-ph/9910306].
- [6] W. Detmold and C.J.D. Lin, Deep-inelastic scattering and the operator product expansion in lattice QCD, Phys. Rev. D 73 (2006) 014501 [hep-lat/0507007].
- [7] V. Braun and D. Müller, *Exclusive processes in position space and the pion distribution amplitude*, *Eur. Phys. J. C* **55** (2008) 349 [0709.1348].
- [8] Z. Davoudi and M.J. Savage, Restoration of Rotational Symmetry in the Continuum Limit of Lattice Field Theories, Phys. Rev. D 86 (2012) 054505 [1204.4146].
- [9] X. Ji, Parton Physics on a Euclidean Lattice, Phys. Rev. Lett. 110 (2013) 262002 [1305.1539].
- [10] A.J. Chambers, R. Horsley, Y. Nakamura, H. Perlt, P.E.L. Rakow, G. Schierholz et al., *Nucleon Structure Functions from Operator Product Expansion on the Lattice, Phys. Rev. Lett.* **118** (2017) 242001 [1703.01153].

- [11] A.V. Radyushkin, Quasi-parton distribution functions, momentum distributions, and pseudo-parton distribution functions, Phys. Rev. D 96 (2017) 034025 [1705.01488].
- [12] Y.-Q. Ma and J.-W. Qiu, Exploring Partonic Structure of Hadrons Using ab initio Lattice QCD Calculations, Phys. Rev. Lett. 120 (2018) 022003 [1709.03018].
- [13] A. Shindler, Moments of parton distribution functions of any order from lattice QCD, Phys. Rev. D 110 (2024) L051503 [2311.18704].
- [14] X. Gao, W.-Y. Liu and Y. Zhao, Parton distributions from boosted fields in the Coulomb gauge, Phys. Rev. D 109 (2024) 094506 [2306.14960].
- [15] HOPE collaboration, Parton physics from a heavy-quark operator product expansion: Formalism and Wilson coefficients, Phys. Rev. D 104 (2021) 074511 [2103.09529].
- [16] C. Michael and I. Teasdale, *Extracting Glueball Masses From Lattice QCD*, *Nucl. Phys. B* 215 (1983) 433.
- [17] M. Luscher and U. Wolff, How to Calculate the Elastic Scattering Matrix in Two-dimensional Quantum Field Theories by Numerical Simulation, Nucl. Phys. B 339 (1990) 222.
- [18] HOPE collaboration, Parton physics from a heavy-quark operator product expansion: Lattice QCD calculation of the second moment of the pion distribution amplitude, Phys. Rev. D 105 (2022) 034506 [2109.15241].
- [19] W.I. Jay and E.T. Neil, *Bayesian model averaging for analysis of lattice field theory results*, *Phys. Rev. D* **103** (2021) 114502 [2008.01069].
- [20] RQCD collaboration, Scale setting and the light baryon spectrum in $N_f = 2 + 1$ QCD with Wilson fermions, JHEP **05** (2023) 035 [2211.03744].
- [21] Bridge++ project, "Lattice QCD code Bridge++." https://bridge.kek.jp/Lattice-code/, 2012-.
- [22] S. Ueda, S. Aoki, T. Aoyama, K. Kanaya, H. Matsufuru, S. Motoki et al., Development of an object oriented lattice QCD code 'Bridge++', J. Phys. Conf. Ser. 523 (2014) 012046.
- [23] Y. Akahoshi, S. Aoki, T. Aoyama, I. Kanamori, K. Kanaya, H. Matsufuru et al., General purpose lattice QCD code set Bridge++ 2.0 for high performance computing, J. Phys. Conf. Ser. 2207 (2022) 012053 [2111.04457].