

# Results from the NA48 experiment on rare neutral kaon decays

Andrew Maier\* †

University of Siegen, Fachbereich Physik, Walter-Flex Straße 3, D-57068 Siegen,  
Germany

E-mail: Andrew.Maier@cern.ch

ABSTRACT: New and recent measurements of the branching ratios of rare decays of neutral  $K_S$  and  $K_L$  mesons are presented. The results have been extracted from data collected during the NA48  $\epsilon'/\epsilon$  runs as well as in a special high intensity  $K_S$  and hyperon run. In particular the presentation focuses on the decay of  $K_L \rightarrow \pi^0 \gamma \gamma$ ,  $K_S \rightarrow \pi^0 e^+ e^-$  and  $K_S \rightarrow \gamma \gamma$ . The prospects of future programs aiming to study direct CP violation in charged kaons and to search for rare  $K_S$  decays are given using the example of  $K_S \rightarrow \pi^0 e^+ e^-$ .

hep2001

## 1. Introduction

The NA48 experiment has been designed to measure the direct CP-violation parameter  $\epsilon'/\epsilon$  [1, 2] to a precision of  $2 \times 10^{-4}$  using simultaneous almost collinear  $K_S$  and  $K_L$  beams [3]. The design of the experiment, its good understanding of the systematics necessary for the measurement of  $\epsilon'/\epsilon$  and the high kaon flux make NA48 an excellent tool for the investigation of rare neutral kaon decays.

This paper reviews a small selection of the rare results from data taken during the 1998/1999  $\epsilon'$  run, as well as results from a two day high intensity  $K_S$  run, taken in 1999.


## 2. $K_L \rightarrow \pi^0 \gamma \gamma$

The decay of  $K_L \rightarrow \pi^0 \gamma \gamma$  is interesting in two ways. Firstly, it is related to the CP conserving amplitude of the decay of  $K_L \rightarrow \pi^0 e^+ e^-$ , which has a direct CP violating component. Secondly, it is a good test for Chiral Perturbation Theory ( $\chi$ PT), since at one loop calculations the rate is finite, yet the O(4)  $\chi$ PT calculations only predict about 1/2 to 1/3 of the measured rate. Calculations of O(6) including a Vector Meson Dominance (VMD) contribution which is parameterised by the parameter  $a_v$ , can accommodate the observed rate. The VMD contribution predicts a mass tail at low  $m_{\gamma\gamma}$ .

\*Speaker.

†on behalf of the NA48 collaboration: Cagliari, Cambridge, CERN, Dubna, Edinburgh, Ferrara, Firenze, Mainz, Orsay, Perugia, Pisa, Saclay, Siegen, Torino, Vienna, Warsaw.

‡Address for correspondence: CERN/EP, CH-1211 Genève 23, Switzerland




**Figure 1:** Invariant mass distribution of the photons resulting from the  $\pi^0$  and the  $\gamma\gamma$ . The photons associated to the  $\pi^0$  are labelled 1 and 2, while the  $\gamma\gamma$  pair is labelled 3 and 4. The coloured areas represent three different regions used in Figure 2.

The data sample used for this measurement was taken from the 98/99  $\epsilon'$  data set, with a large number of  $K_L \rightarrow 2\pi^0$  which have a similar topology as  $K_L \rightarrow \pi^0\gamma\gamma$ .  $K_L \rightarrow 2\pi^0$  has also been used as a normalisation channel. Since both data and normalisation channel are taken under the same trigger conditions, trigger efficiencies cancel.

The background for this channel is mainly due to  $2\pi^0$  and  $3\pi^0$  decays. The background from  $2\pi^0$  is rejected using an invariant mass constraint and by building a  $\chi^2$  like variable with a  $\pi^0$  mass hypothesis in order to reject  $2\pi^0$  events. The background from  $3\pi^0$  originates from missing or overlapping photons in the electro-magnetic calorimeter. This background is rejected using combinatorial cuts which make use of the fact that the vertex is correctly calculated for the  $\pi^0$ , but not for the  $K_L$ . Finally, cuts on the shower width of the photon clusters are made.

Figure 1 shows the invariant mass distribution for the two photons associated to the  $\pi^0$  and the two remaining photons, respectively. A low mass tail in the two  $\gamma$  distribution can be seen, which leads to the interpretation of a non vanishing value of  $a_v$ .

To get an understanding of the value of  $a_v$  it is illustrative to select different mass regions of the  $\gamma\gamma$  mass distribution and to compare the expected number of events for



**Figure 2:** Distribution of the low mass tail for the di- $\gamma$  system for different values of  $a_v$  for data and Monte Carlo.

various values of  $a_v$  from Monte Carlo simulation. This can be seen from Figure 2 where the invariant mass of the  $\pi^0$  is compared with simulation for values of  $a_v$  of 0.0,  $-0.46$  and  $-0.72$ , respectively. While the expectation of a vanishing  $a_v$  underestimates the events in the higher mass region, the large value of  $a_v = -0.72$  overestimates the number of events in the low mass region. The final value of  $a_v$  found in the NA48 data is  $a_v = -0.46 \pm 0.03_{(stat)} \pm 0.03_{(syst)} \pm 0.02_{(theo)}$ . The branching ratio of  $K_L \rightarrow \pi^0 \gamma \gamma$  is determined to be  $BR(K_L \rightarrow \pi^0 \gamma \gamma) = (1.36 \pm 0.03_{(stat)} \pm 0.03_{(syst)} \pm 0.03_{(norm)}) \times 10^{-6}$ .

A publication of this result is in preparation.


### 3. $K_S \rightarrow \pi^0 e^+ e^-$

The decays of  $K_L$  into  $\pi^0 l^+ l^-$  are of considerable interest due to their sensitivity to direct CP violation [4]. However, in  $\pi^0 e^+ e^-$  decay, both CP conserving and indirect CP violating amplitudes contribute. The CP conserving component can be measured from the decay  $K_L \rightarrow \pi^0 \gamma \gamma$ , while the indirect CP violating part can be measured from the decay  $K_S \rightarrow \pi^0 e^+ e^-$ . The  $K_S$  decay is expected to be of the order of  $BR(K_S \rightarrow \pi^0 e^+ e^-) = 5.2 a_s^2 \times 10^{-9}$ , where  $a_s$  is expected to be of  $O(1)$ , but is not well bounded theoretically. The decay has so far not been observed and the best limit to date for its branching ratio has been determined to be  $BR(K_S \rightarrow \pi^0 e^+ e^-) < 1.1 \times 10^{-6}$  at the 90% confidence level by NA31 [5].

The data for this analysis was taken in a two day high intensity test run after the end of the 1999  $e'$  data taking period. Events were selected to have at least 4 clusters in the electro-magnetic calorimeter and two tracks identified as electrons with an  $0.9 < E/p < 1.1$  using the calorimeter and the spectrometer. The invariant mass of the event had to be compatible with the  $K^0$  mass and the invariant mass of the two photons resulting from clusters not associated to tracks had to be compatible with the  $\pi^0$ -mass.

Background from the Dalitz decays  $\pi^0 \pi_D^0$  and  $\pi_D^0 \pi_D^0$  is removed by requiring that the mass of the  $e\gamma$  system is at least 30 MeV larger or smaller than the  $\pi^0$  mass. In addition, the mass of the  $ee$  system is required to be larger than 165 MeV/c<sup>2</sup>. While this cut safely removes all background expected from the simulation, it also removes about 50% of the expected signal (see Figure 3).

After this selection, no events remain in the signal region. Using  $K_S \rightarrow \pi^0 \pi_D^0$  as normalisation channel, a new upper limit for the branching ratio can be given to be:



**Figure 3:**  $K_S \rightarrow \pi^0 \pi_D^0$  and  $K_S \rightarrow \pi^0 e^+ e^-$  for data and Monte Carlo.

$BR(K_S \rightarrow \pi^0 e^+ e^-) < 1.4 \times 10^{-7}$  at the 90% confidence level [6]. This result includes a 7% systematic error and it improves the current best measurement by a factor of 8.

#### 4. $K_S \rightarrow \gamma\gamma$

The interest for this decay arises from the fact that it is calculable in the framework of  $\chi$ PT with no counter-terms. It is also sensitive to loops. Therefore the branching ratio of this decay is predicted by theory with a small error to be  $BR(K_S \rightarrow \gamma\gamma) = (2.3 \pm 0.2) \times 10^{-6}$  [7, 8, 9, 10]. Hence a precision measurement of this mode is an important test to Chiral Perturbation Theory.


The data collected originate from the same two day high intensity  $K_S$  run, which has been used to extract  $K_S \rightarrow \pi^0 e^+ e^-$ . Since the flux of  $K_S$  and  $K_L$  produced at the target are the same, it is important to subtract the number of the events originating from  $K_L \rightarrow \gamma\gamma$  decays from the number of all  $K \rightarrow \gamma\gamma$  events by measuring the flux of the  $K_L$  and using the measured branching ratio of  $K_L \rightarrow \gamma\gamma$ .

Events were selected to have at least two clusters in the electro-magnetic calorimeter. The highest invariant mass  $M_{\gamma\gamma}$  that can be formed by odd pairing photons from  $K_S \rightarrow 2\pi^0$  decays, amounts to 458 MeV, which translates into a vertex shift of 9 m, if such an event is analysed under the  $K_S \rightarrow \gamma\gamma$  hypothesis. The choice of a short decay region, therefore allows to distinguish the signal from the background of  $K_S \rightarrow 2\pi^0$  with 2 missed photons. Monte Carlo studies show, that taking into account also overlapping showers, a region of 5 m leads to an almost background free sample.

The vertex distribution for signal and background, both for data and Monte Carlo, is shown in Figure 4. The contribution from  $K_L \rightarrow \gamma\gamma$  is removed and using  $K_S \rightarrow 2\pi^0$  as a normalisation channel, a branching ratio of  $BR(K_S \rightarrow \gamma\gamma) = (2.58 \pm 0.36_{(stat)} \pm 0.22_{(sys)}) \times 10^{-6}$  is measured [11], in good agreement with theoretical expectations.

#### 5. Outlook

NA48 is involved in a rich program for the search of rare neutral kaon decays. A future run with a high intensity  $K_S$  beam, planned for 2002, will allow competitive physics in the area of rare  $K_S$  and hyperon decays. Already the two day high intensity  $K_S$  run in 1999 was able to improve the limit for the branching ratio for  $K_S \rightarrow \pi^0 e^+ e^-$  by almost an order



**Figure 4:** The vertex distribution for  $K_S \rightarrow \gamma\gamma$ ,  $K_L \rightarrow \gamma\gamma$  and the  $\gamma\gamma$  background originating from  $K_S \rightarrow 2\pi^0$ .

of magnitude. In addition to the  $K_S$  run, a special run of charged kaons is envisaged. This run is scheduled for 2003.

## 6. Acknowledgements

The efforts of all the NA48 members and technical staff are gratefully acknowledged. It is a pleasure to thank the organisers of the EPS meeting in Budapest for an excellent program.

## References

- [1] V. Fanti *et. al.*, *A new measurement of direct cp violation in two pion decays of the neutral kaon*, *Physics Letters B* **465** (1999) 335–348.
- [2] NA48 Collaboration, *A precise measurement of the direct CP violation parameter  $Re(\epsilon'/\epsilon)$* , *CERN-EP-2001-065*, *hep-ex/0110019* (2001). Submitted to European Physical Journal.
- [3] C. Biino *et. al.*, *The simultaneous and nearly-colinear  $K^0$  beams for experiment NA48*, in *Proceedings of the 6th European Particle Accelerator Conference* (S. Myers, L. Lijeby, C. Petit-Jean-Genaz, *et. al.*, eds.), 1998.
- [4] G. D'Ambrosio and G. Isidori, *CP violation in kaon decays*, *International Journal of Modern Physics A* **A13** (1998) 1.
- [5] G. Barr *et. al.* *Physics Letters B* **304** (193) 381.
- [6] A. Lai *et. al.*, *Search for the decay  $K_S \rightarrow \pi^0 e^+ e^-$* , *Physics Letters B* **514** (2001) 253–262.
- [7] G. D'Ambrosio and D. Espriu *Physics Letters B* **175** (1986).
- [8] J. L. Goity *Zeitschrift für Physik C* **34** (1987) 341.
- [9] Z. E. S. Uy *Physical Review D* **3** (1971) 234.
- [10] J. F. Donoghue, “Private communication.”
- [11] A. Lai *et. al.*, *A new measurement of the branching ratio of  $K_S \rightarrow \gamma\gamma$* , *Physics Letters B* **493** (2000) 29–35.