

## Light baryon spectroscopy and diquark correlations in quenched QCD with overlap fermions.

---

**Ronald Babich<sup>\*a</sup>, Federico Berruto<sup>b</sup>, Nicolas Garron<sup>c</sup>, Christian Hoelbling<sup>d</sup>, Joseph Howard<sup>a</sup>, Laurent Lellouch<sup>e</sup>, Claudio Rebbi<sup>a</sup>, and Noam Shoresh<sup>f</sup>**

<sup>a</sup>*Department of Physics, Boston University, Boston, MA*

<sup>b</sup>*Brookhaven National Laboratory, Upton, NY*

<sup>c</sup>*DESY, Platanenallee 6, 15738 Zeuthen, Germany*

<sup>d</sup>*Department of Physics, Bergische Universität Wuppertal, Germany*

<sup>e</sup>*Centre de Physique Théorique<sup>†</sup>, Marseille, France*

<sup>f</sup>*Harvard University, Cambridge, MA*

*E-mail:* rbabich@physics.bu.edu

A simulation of quenched QCD with the overlap Dirac operator has been carried out using 100 Wilson gauge configurations at  $\beta = 6$  on an  $18^3 \times 64$  lattice and at  $\beta = 5.85$  on a  $14^3 \times 48$  lattice. Here we present results for baryon masses and quark and diquark propagators in the Landau gauge.

For the content of this contribution, see PoS(LAT2005)043 in these proceedings.

*XXIIIrd International Symposium on Lattice Field Theory  
25-30 July 2005  
Trinity College, Dublin, Ireland*

---

<sup>\*</sup>Speaker.

<sup>†</sup>UMR 6207 du CNRS et des universités d'Aix-Marseille I, II et du Sud Toulon-Var, affiliée à la FRUMAM.