

Wilson fermions quark bilinears to three loops

Francesco Di Renzo*

University of Parma and I.N.F.N.

E-mail: direnzo@fis.unipr.it

Andrea Mantovi

University of Parma and I.N.F.N.

E-mail: mantovi@fis.unipr.it

Vincenzo Miccio[†]

University of Parma and I.N.F.N.

E-mail: vincenzo.miccio@mib.infn.it

Christian Torrero

University of Parma and I.N.F.N.

E-mail: torrero@fis.unipr.it

Luigi Scorzato

Humboldt University, Berlin

E-mail: Luigi.Scorzato@physik.hu-berlin.de

Quark currents renormalization constants can in principle be safely computed in lattice perturbation theory. In practice, traditional lattice perturbative computations are quite cumbersome, so that so far only the first loop results were available. By making use of Numerical Stochastic Perturbation Theory we reached three (and with less statistical precision even four) loops, both in quenched and in unquenched theory. Convergence properties of the series can be assessed and comparison with non perturbative results (where available) can be made: high loops computations of renormalization constants can be a valuable tool for lattice QCD.

XXIIIrd International Symposium on Lattice Field Theory

25-30 July 2005

Trinity College, Dublin, Ireland

*Speaker.

[†]Current address: I.N.F.N. Milano.

1. Introduction

Despite the fact that logarithmic divergent renormalization constants like those for quark bilinears could be in principle safely computed in Lattice Perturbation Theory (LPT), one usually tries to compute them non-perturbatively, relying on some intermediate scheme. Popular choices are *RI'-MOM* [1] and *SF* [2]. Drawbacks of traditional perturbative computations are well known. LPT is hard and as a result computations of renormalization constants are usually at one loop, with second order computations just now making their entrance¹. Due to the bad convergence properties of LPT, large use is made of so-called Boosted Perturbation Theory (BPT), often in the Tadpole-Improved (TI) variant. By making use of the Numerical Stochastic Perturbation Theory (NSPT) method², we are in a position to perform quite high orders computations: this means three (and even four) loops for quark bilinears. By making use of BPT, we are in a position to assess convergence properties and truncation errors of the series and to make contact with non perturbative results, where available³. This is relevant, since both perturbative and non perturbative computations rely on the same assumptions, which strictly speaking are only proved in the perturbative framework.

2. Computational setup

We performed our computations for Wilson gauge action and (unimproved, at the moment) Wilson fermion action ($r = 1$). The quenched computations have been performed on lattice sizes of both 32^4 and 16^4 . Unquenched computations are at the moment⁴ for $n_f = 2, 3$ on 32^4 . The bigger lattice size fits well on our *APEmille* crate, while standard PC's are enough for the smaller size. In the following we will focus on the $n_f = 2$ results. Configurations have been stored in order to perform many other computations.

We work in the *RI'-MOM* scheme [1]. Quark bilinears operators are computed between (off shell) quark states of momentum p and then amputated to get Γ -functions

$$\langle p | \bar{\psi} \Gamma \psi | p \rangle = G(p) \rightarrow \Gamma(p) = S^{-1}(p) G(p) S^{-1}(p). \quad (2.1)$$

By making use of a convenient projector P_O one then projects on the tree level structure

$$O(p) = \text{Tr}(\hat{P}_O \Gamma(p)) \quad (2.2)$$

to obtain the operators in terms of which the renormalization conditions are given

$$Z_O Z_q^{-1} O(p) \Big|_{p^2=\mu^2} = 1. \quad (2.3)$$

Z_q is the quark field renormalization constants defined via

$$Z_q = -i \frac{1}{12} \frac{\text{Tr}(\not{p} S^{-1})}{p^2} \Big|_{p^2=\mu^2}. \quad (2.4)$$

¹See for example [3] at this conference.

²The method is reviewed in [4] and new technical details were also presented at this conference [5].

³This is the case for the quantities at hand: see [6] at this conference.

⁴ $n_f = 4$ computations are also on their way.

Renormalization conditions are given in the zero quark mass limit in order to get a mass independent scheme. We note with this respect that we are in a position to stay at zero mass by plugging in the Dirac operator the convenient (critical mass) counterterms. These are analytically known at one and two loops [7], while third loop has been computed in [4]. Going to four loop, we obtained as a byproduct the corresponding new order for the critical mass⁵.

The gauge is fixed to Landau, which determines the one loop quark field anomalous dimension to vanish.

One should keep in mind that the *RI'-MOM* scheme is defined in infinite volume, while we are of course forced to finite volume approximations: as we will see, care is to be taken of this point.

In a traditional perturbative computation fixing the divergences (*i.e.* the log's) is the *easy* part of the job, while obtaining finite parts is usually much harder. In NSPT it is just the other way around. We actually take the anomalous dimensions γ 's for granted. In *RI'-MOM* they are known to three loop order [8]: notice that is the reason why we stop at third order. Only for RG-invariant quantities (like the ratio Z_p/Z_s on which we will focus in the following) we have almost no conceptually severe limitations.

In the end, at a given loop L this is what one expects for the coefficient of a renormalization constant:

$$z_L = c_L + \sum_i^L d_i(\gamma) \log(\hat{p})^i + F(\hat{p}) \quad (\hat{p} = pa). \quad (2.5)$$

The first (finite) term is the one we are interested in (in the continuum limit). log's coefficients are known function of anomalous dimensions while the remainder is a scalar function of hypercubic invariants. The way to extract the continuum limit from *hypercubic-symmetric* Taylor expansions is explained in [5], which also contains a practical example, namely for the quark field renormalization constant.

It is useful to see how our master formula Eq. (2.3) reads in a specific case, *i.e.* for the scalar current (first order⁶)

$$\left(1 - \frac{z_q^{(1)}}{\beta} + \dots\right) \left(1 - \frac{z_s^{(1)} - \gamma_s^{(1)} \log(\hat{p}^2)}{\beta} + \dots\right) \left(1 - \frac{o_s^{(1)}}{\beta} + \dots\right) = 1 \quad (2.6)$$

which determines

$$z_s^{(1)} = z_q^{(1)} - \left(o_s^{(1)} - \gamma_s^{(1)} \log(\hat{p}^2)\right) \quad (2.7)$$

Notice that $o_s^{(1)}$ is what is actually coming from measurements, while $z_q^{(1)}$ comes from an independent computation (there is no log because we are in Landau gauge). Higher orders simply requires some more algebra in order to solve for the unknown quantities (modulo some remarks we will make later).

3. Z_p/Z_s to four loops

The *perfect* quantity to compute is the ratio Z_p/Z_s . Quark field renormalization constants drops

⁵A quite surprising result is that resumming the critical mass at four loop in a convenient BPT scheme one obtains a result which is not so far from non perturbative determinations.

⁶We slightly change notation with respect to Eq. (2.3): now the order is understood as the superscript, while subscript denotes the observable.




Figure 1: The resummation of Z_p/Z_s and of Z_s/Z_p ($n_f = 2$) at $\beta = 5.8$. On the x -axis the values of the different couplings x_0, x_1, x_2, x_3 (see text). Color code for y -axis as follows: blue is one loop result, green two loops, red three loops and pale blue four loops.

out in the ratio. There is no anomalous dimension around and so there is no conceptual limitation to our computations (apart from practical ones, at moment mainly dictated by computer memory limitations). As an extra *bonus*, in our perturbative approach Z_p/Z_s and Z_s/Z_p come from *different* signals, so that one has the handle of verifying that the series are actually inverse of each other.

For $n_f = 2$ we obtain

$$\begin{aligned} Z_p/Z_s &= 1 - 0.487(2)\beta^{-1} - 1.46(2)\beta^{-2} - 5.36(7)\beta^{-3} + \dots \\ Z_s/Z_p &= 1 + 0.487(2)\beta^{-1} + 1.70(2)\beta^{-2} + 6.9(1)\beta^{-3} + \dots \end{aligned} \quad (3.1)$$

One can verify that series are actually inverse of each other to a very good precision and we point out that finite size effects have been proved to be well under control by comparing results on different lattice sizes. These results come from the (quite huge) collection of three loops configurations that we stored. We actually pushed the computation even to four loops, even though with less statistics. At four loop we only quote the resummed result. For this ratio the computation is anyway safe. For quantities involving anomalous dimensions things are more cumbersome: one has also to fit an unknown γ , which comes from \log 's. Next step is now to resum the series. We now make use of BPT and resum (for $ap = 1$) using different expansion parameters according to the definitions

$$x_0 \equiv \beta^{-1} \quad x_1 \equiv \frac{\beta^{-1}}{\sqrt{P}} \quad x_2 \equiv -\frac{1}{2} \log(P) \quad x_3 \equiv \frac{\beta^{-1}}{P} \quad (3.2)$$

where P is the plaquette⁷. We resum at $\beta = 5.8$ in order to make contact with the results of [6]. Resummation is plotted in Figure 1: different colors are for different orders of resummation. We

⁷While x_2 and x_3 are widely used, x_1 was defined just in order to better point out the general picture that emerges from the choice of different couplings: see later.




Figure 2: Computation of one loop renormalization constant for the scalar current. Upper points are the unsubtracted $o_s^{(1)}$, while lower (ticker symbols) stand for the subtracted $o_s^{(1)} - \gamma_s^{(1)} \log(\hat{p}^2)$. Analytic result is marked with a darker symbol. On the left: no correction for finite volume. On the right: finite volume *tamed-log* taken into account.

regret that there was an error in the four loop resummation shown at the conference. We here display the correct four loop result, which does not change the overall picture.

From the figure one can better understand the effect of one loop BPT: one is actually sitting on a straight line, whose slope is dictated by the first loop coefficient. Only if higher loops are taken into account a reliable understanding of what is going on can be achieved. In particular, one can easily force the series to oscillate widely by changing the coupling. We stress that convergence properties should be assessed on a case by case basis, *i.e.* depending on the observable. The same holds for TI-BPT and in general if also the scale is changed while switching to a new coupling.

As expected, at a fixed coupling we get milder and milder variations by switching on higher and higher orders, while at a fixed order we get milder and milder variations by changing the coupling. Convergence properties of the series seem reasonable for x_2 and x_3 . For these couplings it is interesting to point out that if one adds to the result at a given order the deviation from the immediately lower order, one always ends up at the same result, which for example in the case of Z_p/Z_s is $0.77(1)$ ⁸. Besides that, resummed series are almost inverse of each other. The comparison with the non perturbative result in [6] yields encouraging result: things are working pretty well in the *RI'-MOM* scheme.

4. Treating anomalous dimensions

Let us go back to Eq. (2.7) (the scalar current at one loop). We make use of this one loop example since the analytic result is known [9]. A naive application of Eq. (2.7) yields what is plotted in Figure 2 on the left: the result is missed, *as if one were subtracting too much by subtracting the log*. The effect is systematic: a similar picture is obtained for example for Z_p . By computing on different lattice sizes, one realizes that is a finite size effect, getting worse and worse as the volume

⁸This is a popular way to pin down a truncation error.

decreases. We stress that this is a $pL \equiv \hat{p}Na = \hat{p}N$ effect, L being expressed in terms of the lattice spacing as $L = Na$. One could say that the log is *tamed* by finite volume. Since we want to regard our finite volume computations as infinite volume approximations, we need to correct for this effect. A solution comes from computing the expected log in the continuum at the same values of $pL = \hat{p}N$ one is interested in. The result is shown in Figure 2 on the right. Notice anyway that if one stays away from the lower momenta results are anyway quite safe.

5. Conclusions and prospects

NSPT is a valuable tool to compute logarithmic-divergent renormalization constants for lattice QCD. Now that configurations are stored, a lot of computations are possible and, in particular, a full account of quark bilinears to three loops will be ready soon. The main caveat when making use of our method is that care is needed when dealing with log's, because of finite volume effects.

We are in a position to assess convergence properties of the series and to shed some light on BPT, a blind application of which is never advisable. Even when a non perturbative result is available, our method can provide an important check to better assess systematic errors.

References

- [1] G. Martinelli, C. Pittori, C.T. Sachrajda, M. Testa and A. Vladikas, *A general method for non perturbative renormalization of lattice operators*, Nucl. Phys. B 445 (1995) 81 [hep-lat/9411010].
- [2] M. Luescher, R. Narayanan, P. Weisz and U. Wolff, *The Schroedinger functional: a renormalizable probe for non abelian gauge theories*, Nucl. Phys. B 384 (1992) 168 [hep-lat/9207009].
- [3] Q.Mason, *High precision fundamental constants using lattice Perturbation Theory*, these proceedings, PoS(Lat2005).
- [4] F. Di Renzo and L. Scorzato, *Numerical Stochastic Perturbation Theory for full QCD*, JHEP 04 (2004) 073 [hep-lat/0410010].
- [5] V. Miccio, F. Di Renzo, A. Mantovi, C. Torrero and L. Scorzato, *Fermionic observables in Numerical Stochastic Perturbation Theory*, these proceedings, PoS(Lat2005)108.
- [6] D. Becirevic, B. Blossier, Ph. Boucaud, V. Gimenez, V. Lubicz, F. Mescia, S. Simula and C. Tarantino, *Non perturbatively renormalized light quark masses with two dynamical fermions*, these proceedings, PoS(Lat2005)079 [hep-lat/0509091].
- [7] E. Follana and H. Panagopoulos, *The critical mass of Wilson fermions: a comparison of perturbative and Monte Carlo results*, Phys. Rev. D 63 (2001) 017501 [hep-lat/0006001];
S. Caracciolo, A. Pelissetto and A. Rago, *Two-loop critical mass for Wilson fermions*, Phys. Rev. D 64 (2001) 094506 [hep-lat/0106013].
- [8] J.A. Gracey, *Three loop anomalous dimension of non-singlet quark currents in the RI' scheme*, Nucl. Phys. B 662 (2003) 247 [hep-ph/0304113].
- [9] G. Martinelli and Yi-Cheng Zhang, *The connection between local operators on the lattice and in the continuum and its relation to meson decay constants*, Phys. Lett. B 123 (1983) 433.