

B Decays with J/psi or Baryons from Belle

Ruslan Chistov^{*†}

Institute for Theoretical and Experimental Physics (ITEP), Russia

E-mail: chistov@itep.ru

We report the results of a study of B -meson decays with J/ψ or baryons in the final state. It includes observation of $B^- \rightarrow J/\psi \Lambda \bar{p}$, $B^{+0} \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^{+0}$ and $B^+ \rightarrow \bar{\Xi}_c^0 \Lambda_c^+$ decays and search for $B^- \rightarrow J/\psi \Sigma^0 \bar{p}$, $B^- \rightarrow J/\psi p \bar{p}$, $B^0 \rightarrow J/\psi \bar{D}^0$, $B^0 \rightarrow J/\psi \bar{D}^0 \pi^+$ and $B^0 \rightarrow \bar{\Xi}_c^- \Lambda_c^+$ decays. These results are based on the analysis of data collected at the $\Upsilon(4S)$ resonance with the Belle detector at the KEKB asymmetric-energy e^+e^- collider.

*International Europhysics Conference on High Energy Physics
July 21st - 27th 2005
Lisboa, Portugal*

^{*}Speaker.

[†]On behalf of the Belle Collaboration

1. Introduction

Recently, the Belle Collaboration observed a number of baryonic B decays, charmless [1] and charmful [2]. Belle continues a study of such decays and reports the results on the new modes.

To isolate the B -meson signal For all reported decays, to isolate the B -meson signal we use $\Delta E = \Sigma_i E_i - E_{\text{beam}}$ (or $\Delta M = M(B)_{\text{inv.}} - M(B)_{\text{table}}$) and beam-energy constrained mass $M_{\text{bc}} = \sqrt{E_{\text{beam}}^2 - (\Sigma_i \vec{p}_i)^2}$. Here $E_{\text{beam}} = \sqrt{s}/2$ is the beam energy in the center of mass, \vec{p}_i and E_i are the three-momenta and energies of the B candidate's decay products, $M(B)_{\text{inv.}}$ and $M(B)_{\text{table}}$ are the reconstructed B -meson mass and the world average B -meson mass.

2. Observation of $B^- \rightarrow J/\psi \Lambda \bar{p}$ and Searches for $B^- \rightarrow J/\psi \Sigma^0 \bar{p}$ and $B^0 \rightarrow J/\psi p \bar{p}$

We present the observation of the decay mode $B^- \rightarrow J/\psi \Lambda \bar{p}$, which is a new type of baryonic B decay, $B \rightarrow \text{charmonium} + \text{baryons}$. Modes of this type were proposed as a potential explanation for the excess in the low momentum region of the inclusive J/ψ momentum spectrum in B decays [3]. The measured branching fraction and the results of a search for the related modes are shown in Table 1.

3. Search for $B^0 \rightarrow J/\psi \bar{D}^0$ and $B^+ \rightarrow J/\psi \bar{D}^0 \pi^+$

A search for this mode is motivated by the proposed intrinsic charm ($q\bar{b}c\bar{c}$) in the B meson as another explanation for the excess in the soft part of inclusive J/ψ momentum spectrum mentioned in the previous section [4]. The results of this search are presented in Table 1.

4. Observation of $B^{+0} \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^{+0}$

Recently, Belle observed two-body baryonic B decay $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{p}$ [10] proceeding via $b \rightarrow c\bar{u}d$ transition. In this and next section we present the first observations of exclusive B decays into two charmed baryons in the final state proceeding via $b \rightarrow c\bar{c}s$ transition. Figure 1 shows the ΔM and M_{bc} for the B^+ -meson decaying into $\Lambda_c^+ \bar{\Lambda}_c^- K^+$. The measured branching fraction for this decay together with that for the isospin related mode $B^0 \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^0$ are presented in Table 1.

5. Observation of $B^+ \rightarrow \bar{\Xi}_c^0 \Lambda_c^+$ and Evidence for $B^0 \rightarrow \bar{\Xi}_c^- \Lambda_c^+$

The $\Delta E(M_{bc})$ distributions for the $B^+ \rightarrow \bar{\Xi}_c^0 \Lambda_c^+$ and $B^0 \rightarrow \bar{\Xi}_c^- \Lambda_c^+$ decays are shown in Fig. 2 a(b) and Fig. 2 c(d), respectively, where the superimposed curves are the fit results. The measured products of branching fractions are presented in Table 1.

Taking into account the theoretical predictions for $\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$ of $\sim (0.9 - 2)\%$ [9] and the Belle measurement of $\mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+ \bar{p})$ [10] we obtain $\mathcal{B}(B \rightarrow \bar{\Xi}_c^0 \Lambda_c^+) / \mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+ \bar{p}) \sim 100$. This disagrees with the naive expectation that the branching fractions for two-body baryonic B decays proceeding via $b \rightarrow c\bar{c}s$ and $b \rightarrow c\bar{u}d$ transitions should be of the same order [11].

6. Summary

In conclusion, the Belle Collaboration observes for the first time new types of B decays with J/ψ or baryons in the final state: $B^- \rightarrow J/\psi \Lambda \bar{p}$, $B^{+0} \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^{+0}$ and $B^+ \rightarrow \bar{\Xi}_c^0 \Lambda_c^+$. Despite small energy release in all of these modes, the measured branching fractions are in the range of $(10^{-5} - 10^{-3})$. Two latter modes are the first examples of exclusive B decays into two charmed baryons. All these new results provide additional information on the mechanism in B decays of baryon formation.

Table 1: Summary of reported results

| Decay Branching Fraction | Result | Significance, σ 's |
|--|--|---------------------------|
| $\mathcal{B}(B^- \rightarrow J/\psi \Lambda \bar{p})$ | $(11.6 \pm 2.8_{-2.3}^{+1.8}) \times 10^{-6}$ [5] | 11.1 |
| $\mathcal{B}(B^- \rightarrow J/\psi \Sigma^0 \bar{p})$ | $< 1.1 \times 10^{-5}$ @ 90% CL [5] | - |
| $\mathcal{B}(B^0 \rightarrow J/\psi p \bar{p})$ | $< 8.3 \times 10^{-7}$ @ 90% CL [5] | - |
| $\mathcal{B}(B^0 \rightarrow J/\psi \bar{D}^0)$ | $< 2.0 \times 10^{-5}$ @ 90% CL [6] | - |
| $\mathcal{B}(B^+ \rightarrow J/\psi \bar{D}^0 \pi^+)$ | $< 2.5 \times 10^{-5}$ @ 90% CL [6] | - |
| $\mathcal{B}(B^+ \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^+)$ | $(6.5_{-0.9}^{+1.0} \pm 1.1 \pm 3.4) \times 10^{-4}$ [7] | 15.4 |
| $\mathcal{B}(B^0 \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^0)$ | $(7.9_{-2.3}^{+2.9} \pm 1.2 \pm 4.1) \times 10^{-4}$ [7] | 6.6 |
| $\mathcal{B}(B^+ \rightarrow \bar{\Xi}_c^0 \Lambda_c^+) \times \mathcal{B}(\bar{\Xi}_c^0 \rightarrow \bar{\Xi}^+ \pi^-)$ | $(4.8_{-0.9}^{+1.0} \pm 1.1 \pm 1.2) \times 10^{-5}$ [8] | 8.7 |
| $\mathcal{B}(B^0 \rightarrow \bar{\Xi}_c^- \Lambda_c^+) \times \mathcal{B}(\bar{\Xi}_c^- \rightarrow \bar{\Xi}^+ \pi^- \pi^-)$ | $(9.3_{-2.8}^{+3.7} \pm 1.9 \pm 2.4) \times 10^{-5}$ [8] | 3.8 |

References

- [1] Belle Collaboration, Y.-J. Lee, M.-Z. Wang *et al.*, Phys. Rev. Lett. **93**, 211801 (2004); Belle Collaboration, M.-Z. Wang, Y.-J. Lee, *et al.*, Phys. Rev. Lett. **90**, 201802 (2003); Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **88**, 181803 (2002).
- [2] Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **89**, 151802 (2002); H. Kichimi, Nucl. Phys. B Proc. Suppl. **142**, 197 (2005).
- [3] S. J. Brodsky and F. S. Navarra, Phys. Lett. B **411**, 152 (1997).
- [4] C. H. Chang and W. S. Hou, Phys. Rev. D **64**, 071501(R) (2001).
- [5] Belle Collaboration, L. M. Zhang, Z. P. Zhang *et al.*, Phys. Rev. D **71**, 091107 (2005).
- [6] Belle Collaboration, Q. L. Xie *et al.*, Phys. Rev. D **72**, 051105(R) (2005).
- [7] Belle Collaboration, K. Abe *et al.*, hep-ex/0508015.
- [8] Belle Collaboration, R.Chistov *et al.*, hep-ex/0510074.
- [9] B. Desplanques, J. F. Donoghue and B. R. Holstein, Annals Phys. **124**, 449 (1980); P. Zenczykowski, Phys. Rev. D **40**, 2290 (1989); P. Zenczykowski, Phys. Rev. D **50**, 402 (1994).
- [10] Belle Collaboration, N. Gabyshev, H. Kichimi *et al.*, Phys. Rev. Lett. **90**, 121802 (2003).
- [11] V. L. Chernyak and I. R. Zhitnitsky, Nucl. Phys B **345**, 137 (1990).




Figure 1: The ΔM (upper) and M_{bc} (lower) distributions for the B^+ -meson decaying into $\Lambda_c^+ \bar{\Lambda}_c^- K^+$.




Figure 2: The ΔE and M_{bc} distributions for the $B^+ \rightarrow \bar{\Lambda}_c^0 \Lambda_c^+ K^+$ (a, b) and $B^0 \rightarrow \bar{\Lambda}_c^- \Lambda_c^+ K^+$ (c, d) candidates.