

The complete lowest order chiral Lagrangian from a little box

Thomas DeGrand*

Physics Department, University of Colorado, Boulder CO 80309 USA

E-mail: thomas.degrand@colorado.edu

Stefan Schaefer

NIC, DESY, Platanenallee 6, D-15738 Zeuthen, Germany

E-mail: stefan.schaefer@desy.de

We recently performed a pilot study determining the parameters of the leading order chiral Lagrangian from distributions of the eigenvalues of a quenched Dirac operator coupled to an imaginary isospin chemical potential. We complement a quick survey of our recent preprint [1] by addressing some points raised during discussions at the conference.

*The XXV International Symposium on Lattice Field Theory
July 30 - August 4 2007
Regensburg, Germany*

*Speaker.

1. The project, in a nutshell

Two coupling constants, F and Σ , parameterize the leading-order chiral effective Lagrangian

$$\mathcal{L}_{\text{eff}} = \frac{F^2}{4} \text{Tr}(\partial_\mu U \partial^\mu U^\dagger) - \frac{\Sigma}{2} \text{Tr}[\mathcal{M}(U + U^\dagger)]. \quad (1.1)$$

A long-standing problem for lattice simulations is to determine them directly from QCD. Typically, this is done by fitting a correlation function to some theoretical formula parameterized by F , Σ and m_π . Usually one has to measure a correlation function of operators at long distance to be sensitive to these parameters. This makes the measurement expensive. In addition, the simulations do not give F and Σ directly; instead, they give the mass-dependent decay constant and condensate, which depend, through the standard formulas of chiral perturbation theory, on F and Σ and on the coefficients of higher-order terms in \mathcal{L} . So the determination of F and Σ from the lattice is rather indirect.

A series of papers by Akemann, Damgaard, Heller, Osborn, Splittorf, Svetitsky, and Toublan[2, 3, 4, 5] provide an alternative path toward a direct measurement of F and Σ . This is done via the properties of eigenvalues of the Dirac operator, from simulations in the epsilon regime ($m_\pi L \ll 1$ but $\Lambda L \gg 1$ for box length L and Λ any nonperturbative parameter of QCD except m_π). The particular implementation we used is that of Ref. [5]. An imaginary isospin chemical potential μ is coupled to a doublet of quenched quarks. The correlator of the eigenvalues from an ordinary $\mu = 0$ simulation, λ_i , with the quenched eigenvalues, λ_j ,

$$\rho_{(1,1)}^{(2)\text{conn}}(x,y) = \langle \sum_i \delta(x - \lambda_i) \sum_j \delta(y - \lambda_j) \rangle - \langle \sum_i \delta(x - \lambda_i) \rangle \langle \sum_j \delta(y - \lambda_j) \rangle \quad (1.2)$$

is a function of Σ and F which is given by Random Matrix Theory (RMT). The particular formula is given by Eq. (3.49) of Ref. [5], in terms of the usual rescaled variables $\lambda_i \Sigma V$, $\lambda_j \Sigma V$, the rescaled mass $m_q \Sigma V$, and the rescaled isospin chemical potential $\delta = \mu F \sqrt{V}$. V is the volume.

To combine data sets, we integrate the data. Refs. [2] and [3] suggest using

$$C(x, \zeta_{\text{max}}) = \int_0^{\zeta_{\text{max}}} dy \rho(x+y, y). \quad (1.3)$$

We did one more integral:

$$I(X, \zeta_{\text{max}}) = \int_{-\zeta_{\text{max}}}^X C(x, \zeta_{\text{max}}) dx. \quad (1.4)$$

The first integral $C(x)$ shows a spike at $x = 0$ whose width goes roughly as δ^2 . Then $I(X)$ will show a sharp step at $X = 0$.

In our numerical simulations we used overlap fermions[6, 7]. They preserve the full $SU(N_f) \otimes SU(N_f)$ chiral symmetry (including the anomalous singlet current and the index theorem). Our data set uses a lattice volume of 12^4 points. The lattice spacing a , as determined from the Sommer parameter r_0 [8], is $r_0/a = 3.71(5)$. The pseudoscalar mass is $am_\pi = 0.329(3)$. This is not in the epsilon regime, but experience shows RMT is robust enough to work well outside the epsilon regime, and we are making a pilot study. In our conversions to physical units we use $r_0 = 0.5$ fm. Details of our action are to be found in Refs. [9, 10, 11]. The one new ingredient is the gauge




Figure 1: Data (squares) and fit from $|\nu| = 1$. A cut $\zeta_{max} = 0.07$ is enforced. The best fit values for this bootstrap sample are $\Sigma_L V = 147$ and $aF_L = 0.071$.




Figure 2: Best-fit values of aF_L , varying ζ_{max} .

connection: we used the differentiable hypercubic smeared link of Ref. [12]. We collected about 400 thermalized HMC trajectories of unit length and analyzed 30 lattices in topological charge sector $\nu = 0$ and 75 $|\nu| = 1$ ones. To do the fitting, we took bootstrap averages of the integrated data.

We reproduce two figures from the paper: Fig. 1 shows an example of a fit from one of our bootstraps. Fig. 2 shows the dependence of fits on ζ_{max} . These fits all include only the three lowest (but nonzero) eigenvalues in a topological sector. Smaller ζ_{max} means that less data is included in the fit; larger ζ_{max} means that more eigenmodes are needed to saturate the correlation function. It appears that results for this data set do not depend on the choice of ζ_{max} .

In finite volume, the eigenvalue correlator is not a direct function of Σ and F , but of the quantities Σ_L and F_L which reflect the finite volume. For the condensate, this dependence is known: $\Sigma_L = \rho_\Sigma \Sigma$ where

$$\rho_\Sigma = 1 + C_\Sigma \frac{1}{F^2} \Delta(0) + \dots \quad (1.5)$$

with $\Delta(0)$ the contribution to the tadpole graph (propagator at zero separation) from finite-volume image terms. In the epsilon regime, $\Delta(0) = -D/\sqrt{V}$ and D depends on the geometry[13]. (It is 0.1405 for hypercubes.) For N_f flavors, $C_\Sigma = -(N_f^2 - 1)/N_f$.

A second complication for the condensate is that Σ is scheme-dependent. We perform the conversion from lattice regularization the \overline{MS} using a measurement of the matching factor from the RI-MOM scheme as an intermediary. Correcting for finite volume and converting schemes, we find

$$r_0 \Sigma(\overline{MS}, \mu = 2 \text{ GeV})^{1/3} = 0.594(13) \quad (1.6)$$

or

$$(\Sigma(\overline{MS}, \mu = 2 \text{ GeV}))^{1/3} = 234(4) \text{ MeV} \quad (1.7)$$

This is consistent with an analysis of cumulants of eigenvalues, and with other determinations of the $N_f = 2$ condensate[14, 15, 16, 17].

F is also expected to be volume dependent, and the correction is also likely to be of the form $F_L = \rho_F F$ with

$$\rho_F = 1 + C_F \frac{1}{F^2} \Delta(0) + \dots \quad (1.8)$$

Unfortunately, there is no calculation (yet) of C_F . We found

$$r_0 F_L = 0.255(13) \quad (1.9)$$

or with $r_0 = 0.5 \text{ fm}$,

$$F_L = 101(6) \text{ MeV}. \quad (1.10)$$

We expect that C_F is a number on the order 1-2, probably N_f dependent; that would be sufficient to lower F from F_L by 20 per cent or so and bring it in line with phenomenological estimates of about 86 MeV [18, 19].

So at this point the calculation is incomplete: one needs the finite volume correction. However, the size of the error is very interesting: while typical large scale simulations can get f_π to a fraction of a per cent, F is harder: for example, MILC[20] quotes (their Table IV, Fit A, our conversion) $r_1 F = 0.131(10)$ or $F = 82(6) \text{ MeV}$, compared to their number for f_π , which has a 0.3 per cent error [21]). Our quite competitive error on F , with the expenditure of a tiny amount of computer resources, is what makes this calculation worth repeating.

2. Questions from the conference

- You used fat links. Aren't you worried they will mess something up?

On the contrary. Everyone who has tried using fat links for light fermions has only good things to say about them, from the point of view of perturbation theory, exceptional configurations (for Wilson fermions), taste symmetry restoration (for staggered fermions) and efficiency in computing the overlap action. The one exception was MILC, who long ago[22] tried to use APE smearing with 0.45×10 (!!!) smearing steps in a calculation of the properties of heavy-light mesons. This gave big scaling violations in f_D : too much smearing decoupled the quarks from the short distance part of the one gluon exchange potential. We are using hypercubic smearing. This is even more local than AsqTad – in fact, you cannot be more local. Readers might remember that the Alpha collaboration studied static-light properties with hypercubic links, with great success [23].

- Why did you measure F ? Isn't f_π more interesting?

Actually, what is interesting is the ratio f_K/f_π , which can be used to determine the Cabibbo angle[24]. But then you have to ask, how do you measure f_π ? This is not done with simulations at the physical quark masses; instead, one performs simulations at any light quark masses (heavy enough to be cheaply simulated, light enough to be in the region of validity of chiral perturbation theory). Then the mass-dependence of parameters is fit to a chiral perturbation theory formula. The uncertainties between F , Σ and the higher order terms cancel at the input quark mass values, for the particular observables which are being fit. However, for other observables (predictions) the larger error on F can be problematic. That's why this particular measurement scheme – provided the finite volume correction can be controlled – is so interesting.

- Isn't there a problem using individual eigenvalues to determine F and Σ , rather than using the complete spectral density $\rho(\zeta)$?

We don't think so (and anyway, we used formulas which involved integrated eigenvalue distributions). Ref. [25] showed that the spectral density and its moments, and even eigenvalues, renormalize pretty much as one would naively expect. Notice that at low values of ζ , $\rho(\zeta)$ coincides with the distribution of the lowest eigenvalue $p_1(\zeta)$ (because that is all that there is)! Akemann and Damgaard [26] have derived exact expression relating the individual eigenvalue distributions to integrals over the eigenvalue density and all the spectral correlators. If these are true, it is hard to imagine that something different would happen to the individual eigenvalue distributions and to the integrated ones. (They are also working on a single-eigenvalue version of the correlator we studied here[27].)

- I don't have a chiral action. How much of this project could I do?

Doing exactly what we did with fermions which do not recognize the index theorem could be tricky, since you have to work in sectors of fixed topology, and you have to project the nonzero eigenvalues, which are complex for a generic non-chiral Wilson-type fermion, onto the imaginary axis. This might be more trouble than it is worth. You might try looking at quantities in the epsilon regime which average over topology; the original papers by Hansen and Leutwyler[28] give formulas for the pseudoscalar and axial correlators, which can also be analyzed to get Σ and F .

- Tell me more about the “Primme” package.

The “Primme” package of McCombs and Stathopoulos[29] is a library for computing eigenvalues of Hermitian matrices. (For overlap fermions, that is all you need.) It implements a variety of algorithms, preconditioning and optimization options which can be used via a very convenient interface. It is about a factor of three faster than our version of the Bunk-Kalkreuter-Simma [30] Conjugate Gradient algorithm.

- Can you suggest a good book about lattice QCD?

Yes [31].

3. Acknowledgments

This work was supported by the US Department of Energy. T. D. would like to thank Poul Damgaard for his encouragement and for much valuable advice. He would also like to thank Kim Splittorff for clarifying correspondence. We would like to thank Hidenori Fukaya and Silvia Necco for discussions about finite volume corrections.

References

- [1] T. DeGrand and S. Schaefer, arXiv:0708.1731 [hep-lat].
- [2] P. H. Damgaard, U. M. Heller, K. Splittorff and B. Svetitsky, Phys. Rev. D **72**, 091501 (2005) [arXiv:hep-lat/0508029].
- [3] P. H. Damgaard, U. M. Heller, K. Splittorff, B. Svetitsky and D. Toublan, Phys. Rev. D **73**, 074023 (2006) [arXiv:hep-lat/0602030].
- [4] P. H. Damgaard, U. M. Heller, K. Splittorff, B. Svetitsky and D. Toublan, Phys. Rev. D **73**, 105016 (2006) [arXiv:hep-th/0604054].
- [5] G. Akemann, P. H. Damgaard, J. C. Osborn and K. Splittorff, Nucl. Phys. B **766**, 34 (2007) [arXiv:hep-th/0609059].
- [6] H. Neuberger, Phys. Lett. B **417**, 141 (1998) [arXiv:hep-lat/9707022].
- [7] H. Neuberger, Phys. Rev. Lett. **81**, 4060 (1998) [arXiv:hep-lat/9806025].
- [8] R. Sommer, Nucl. Phys. B **411**, 839 (1994) [arXiv:hep-lat/9310022].
- [9] T. A. DeGrand [MILC collaboration], Phys. Rev. D **63**, 034503 (2001) [arXiv:hep-lat/0007046].
- [10] T. A. DeGrand and S. Schaefer, Phys. Rev. D **71**, 034507 (2005) [arXiv:hep-lat/0412005].
- [11] T. DeGrand and S. Schaefer, JHEP **0607**, 020 (2006) [arXiv:hep-lat/0604015].
- [12] A. Hasenfratz, R. Hoffmann and S. Schaefer, JHEP **0705**, 029 (2007) [arXiv:hep-lat/0702028].
- [13] J. Gasser and H. Leutwyler, Phys. Lett. B **184**, 83 (1987); Nucl. Phys. B **307**, 763 (1988). See also P. Hasenfratz and H. Leutwyler, Nucl. Phys. B **343**, 241 (1990).
- [14] Compare the summaries in C. McNeile, Phys. Lett. B **619**, 124 (2005) [arXiv:hep-lat/0504006].
- [15] T. DeGrand, Z. Liu and S. Schaefer, Phys. Rev. D **74**, 094504 (2006) [Erratum-ibid. D **74**, 099904 (2006)] [arXiv:hep-lat/0608019].
- [16] C. B. Lang, P. Majumdar and W. Ortner, Phys. Lett. B **649**, 225 (2007) [arXiv:hep-lat/0611010].
- [17] H. Fukaya *et al.* [JLQCD Collaboration], Phys. Rev. Lett. **98**, 172001 (2007) [arXiv:hep-lat/0702003].
- [18] J. Gasser and H. Leutwyler, Annals Phys. **158**, 142 (1984).
- [19] G. Colangelo and S. Durr, Eur. Phys. J. C **33**, 543 (2004) [arXiv:hep-lat/0311023].
- [20] C. Aubin *et al.* [MILC Collaboration], Phys. Rev. D **70**, 114501 (2004) [arXiv:hep-lat/0407028].
- [21] C. Bernard *et al.* [MILC Collaboration], arXiv:hep-lat/0609053.
- [22] C. Bernard *et al.* [MILC Collaboration], Phys. Rev. D **66**, 094501 (2002) [arXiv:hep-lat/0206016].

- [23] M. Della Morte, S. Durr, J. Heitger, H. Molke, J. Rolf, A. Shindler and R. Sommer [ALPHA Collaboration], Phys. Lett. B **581**, 93 (2004) [Erratum-ibid. B **612**, 313 (2005)] [arXiv:hep-lat/0307021].
- [24] W. J. Marciano, Phys. Rev. Lett. **93**, 231803 (2004) [arXiv:hep-ph/0402299].
- [25] L. Del Debbio, L. Giusti, M. Luscher, R. Petronzio and N. Tantalo, JHEP **0602**, 011 (2006) [arXiv:hep-lat/0512021].
- [26] G. Akemann and P. H. Damgaard, Phys. Lett. B **583**, 199 (2004) [arXiv:hep-th/0311171].
- [27] G. Akemann and P. H. Damgaard, arXiv:0709.0484 [hep-lat].
- [28] F. C. Hansen, Nucl. Phys. B **345**, 685 (1990); F. C. Hansen and H. Leutwyler, Nucl. Phys. B **350**, 201 (1991).
- [29] A. Stathopoulos, Nearly optimal preconditioned methods for hermitian eigenproblems under limited memory. Part I: Seeking one eigenvalue, SIAM J. Sci. Comput., Vol. 29, No. 2, (2007), 481–514. A. Stathopoulos and J. R. McCombs, Nearly optimal preconditioned methods for hermitian eigenproblems under limited memory. Part II: Seeking many eigenvalues, Tech Report: WM-CS-2006-02, June, 2006.
- [30] See B. Bunk, K. Jansen, M. Lüscher, and H. Simma, unpublished DESY report (1994); T. Kalkreuter and H. Simma, Comput. Phys. Commun. **93**, 33 (1996) [hep-lat/9507023].
- [31] T. DeGrand and C. DeTar, *Lattice Methods for Quantum Chromodynamics*, (World Scientific, Singapore, 2006).