

Perturbative determination of c_{SW} with Symanzik improved gauge action and stout smearing

Roger Horsley

School of Physics, University of Edinburgh, Edinburgh EH9 3JZ, UK

E-mail: rhorsley@ph.ed.ac.uk

Holger Perlt*, Arwed Schiller

Institut für Theoretische Physik, Universität Leipzig, PF 100 920, 04009 Leipzig, Germany

E-mail: Holger.Perlt@itp.uni-leipzig.de

E-mail: Arwed.Schiller@itp.uni-leipzig.de

Paul E.L. Rakow

Theoretical Physics Division, Department of Mathematical Sciences, University of Liverpool,

Liverpool L69 3BX, UK

E-mail: rakow@amtp.liv.ac.uk

Gerrit Schierholz

Deutsches Elektronen-Synchrotron DESY, 22603 Hamburg, Germany

E-mail: Gerrit.Schierholz@desy.de

We determine the improvement factor c_{SW} in one-loop lattice perturbation theory for the plaquette and Symanzik improved gauge actions. The fermionic action is $\mathcal{O}(a)$ clover improved with one-time stout smearing. c_{SW} is derived from the one-loop correction to the quark-quark-gluon vertex in the off-shell regime. We give a first numerical value for the one-loop contribution to the non gauge-invariant improvement coefficient c_{NGI} for the quark field using the plaquette action. A discussion of mean field improvement is included.

The XXV International Symposium on Lattice Field Theory

July 30 - August 4, 2007

Regensburg, Germany

*Speaker.

1. Introduction

Current simulations with 2 + 1 flavors require highly improved gauge and quark actions. Renormalization group improved gauge actions are order $\mathcal{O}(a^2)$ improved and should be preferred to the $\mathcal{O}(a)$ improved plaquette gauge action. In accordance with our numerical simulations we take the Symanzik improved gauge action [1, 2]

$$S_G^{\text{Symanzik}} = \frac{6}{g^2} \left[c_0 \sum_{\text{plaq}} \frac{1}{3} \text{Re Tr}(1 - U_{\text{plaq}}) + c_1 \sum_{\text{rect}} \frac{1}{3} \text{Re Tr}(1 - U_{\text{rectangle}}) \right] \quad (1.1)$$

with

$$c_1 = -\frac{1}{12}, \quad c_0 = 1 - 8c_1.$$

As the fermionic action, we use the clover improved action as proposed by Sheikholeslami and Wohlert [3] which means that one has to add the so-called clover term to the standard Wilson fermion action

$$S_F^{\text{clover}} = S_F^{\text{Wilson}} - c_{\text{SW}} \sum_n \sum_{\mu, \nu} i g \frac{r}{4} \bar{\psi}_n \sigma_{\mu\nu} F_{\mu\nu}(n) \psi_n, \quad (1.2)$$

where $F_{\mu\nu}(n)$ is the field strength in clover form and $\sigma_{\mu\nu} = i/2(\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu)$. An additional improvement can be achieved with ultraviolet filtering or smearing the gauge links U_μ in the fermionic Wilson action S_F^{Wilson} : it reduces the chiral symmetry breaking of Wilson quarks among light flavors. There have been proposed several smearing techniques (for a detailed discussion see [4]). We use the stout smearing of Morningstar and Peardon [5]. It is given by a sequence of transformations

$$U_\mu \rightarrow U_\mu^{(1)} \rightarrow U_\mu^{(2)} \dots \rightarrow U_\mu^{(n)} = \tilde{U}_\mu, \quad (1.3)$$

with

$$U_\mu^{(n+1)}(x) = e^{iQ_\mu^{(n)}(U, \omega_{\mu\nu})} U_\mu^{(n)}(x).$$

The function $Q_\mu^{(n)}(U, \omega_{\mu\nu})$ depends on the staples of the gauge link under consideration and on the stout parameters $\omega_{\mu\nu}$ which determine the strength of smearing. We chose an isotropic parameter $\omega_{\mu\nu} = \omega$ and one step smearing which is recommended by various investigations.

It is of importance to determine the improvement factor c_{SW} appearing in (1.2) as precisely as possible. Non-perturbative determinations are to be preferred but for the combination described above there are no results obtained so far. In perturbation theory c_{SW} has the form

$$c_{\text{SW}} = 1 + g^2 c_{\text{SW}}^{(1)} + \mathcal{O}(g^4). \quad (1.4)$$

There have been published results for $c_{\text{SW}}^{(1)}$ for plaquette action with twisted antiperiodic boundary conditions [6] and Schrödinger functional method [7]. For some popular improved gauge actions Aoki and Kuramashi [8] calculated the one-loop correction using conventional perturbation theory. All results are obtained for unsmearred gauge links in the on-shell regime.

In this paper we calculate $c_{\text{SW}}^{(1)}$ for Symanzik improved gauge action with stout smearing in conventional perturbation theory. We do the calculation off-shell. This enables us to determine the one-loop contribution to the non gauge-invariant improvement coefficient c_{NGI} for the quark

fields ψ as proposed in [9]. Using BRST symmetry arguments the authors proposed the off-shell improvement for the quark fields ψ_* to be

$$\psi_* = (1 + a c_D \overrightarrow{D} + a i g c_{NGI} \not{A}) \psi, \quad (1.5)$$

where the coefficient c_{NGI} does not contribute on-shell. Its perturbative expansion is known to be [9]

$$c_{NGI} = g^2 c_{NGI}^{(1)} + \mathcal{O}(g^4). \quad (1.6)$$

In order to determine $c_{NGI}^{(1)}$ either a two-loop calculation of the quark propagator or a one-loop calculation of the quark-quark-gluon vertex is required.. The improvement coefficient c_D has been calculated to one-loop order in [14].

2. Improvement procedure

In the approach of conventional perturbation theory we use the quark-quark-gluon vertex $\Lambda_\mu(p_1, p_2)$ as discussed in [8] already. Looking at the $\mathcal{O}(a)$ expansion of tree-level $\Lambda_\mu^{(0)}(p_1, p_2)$ as derived from action (1.2)

$$\Lambda_\mu^{(0)}(p_2, p_1) = -i g \gamma_\mu - g \frac{1}{2} a r \mathbf{1}(p_1 + p_2)_\mu - c_{SW} i g \frac{1}{2} a r \sigma_{\mu\alpha} (p_2 - p_1)_\alpha + \mathcal{O}(a^2), \quad (2.1)$$

one can see by inserting (1.4) that a one-loop calculation for $\Lambda_\mu(p_2, p_1)$ provides necessary conditions to determine $c_{SW}^{(1)}$. We omit in all three-point functions the common overall color factor t^{ac} . In (2.1) p_1 (p_2) are the incoming (outgoing) momenta. The off-shell improvement condition states that the non-amputated improved three-point function $G_{*,\mu}(p_2, p_1)$ has to be free of $\mathcal{O}(a)$ terms in one-loop. The unimproved and improved three-point functions are defined by

$$G_\mu(p_2, p_1) = S(p_2) \Lambda_\nu(p_2, p_1) S(p_1) D_{\nu\mu}(q), \quad (2.2)$$

$$G_{*,\mu}(p_2, p_1) = S_*(p_2) \Lambda_{*,\nu}(p_2, p_1) S_*(p_1) D_{\nu\mu}(q), \quad (2.3)$$

with $q = p_2 - p_1$. $D_{\nu\mu}(q)$ is the full gluon propagator which is $\mathcal{O}(a)$ -improved already. $\Lambda_\mu(p_2, p_1)$ and $\Lambda_{*,\mu}(p_2, p_1)$ are the unimproved and improved amputated three-point functions. The corresponding quark propagators are given by

$$S^{-1}(p) = i \not{p} \Sigma_p(p) + \frac{ap^2}{2} \Sigma_W(p) = i \not{p} \Sigma_p(p) \left(1 - \frac{1}{2} a i \not{p} \frac{\Sigma_W(p)}{\Sigma_p(p)} \right), \quad (2.4)$$

$$S_*^{-1}(p) = i \not{p} \Sigma_p(p). \quad (2.5)$$

In terms of the improved quark fields (1.5) $G_\mu(p_2, p_1)$ can be related to its improved version

$$G_\mu(p_2, p_1) = G_{*,\mu}(p_2, p_1) - a i g c_{NGI} \mathcal{F} \left[\langle (\not{A} \overrightarrow{D}^{-1} + \overrightarrow{D}^{-1} \not{A}) A_\mu \rangle \right]. \quad (2.6)$$

In deriving (2.6) we have assumed $\langle A \rangle = 0$, \mathcal{F} denotes the Fourier transform. Taking into account (1.6) we insert in our one-loop calculation the corresponding tree-level expressions

$$a i g c_{NGI} \mathcal{F} \left[\langle (\not{A} \overrightarrow{D}^{-1} + \overrightarrow{D}^{-1} \not{A}) A_\mu \rangle^{tree} \right] = a i g^3 c_{NGI}^{(1)} \left(\gamma_\nu \frac{1}{i \not{p}_1} + \frac{1}{i \not{p}_2} \gamma_\nu \right) D_{\nu\mu}^{tree}(q), \quad (2.7)$$

or its amputated version

$$aig c_{NGI} \mathcal{F} [\langle (A \mathcal{D}^{-1} + \mathcal{D}^{-1} A) A_\mu \rangle_{amp}^{tree}] = -a g^3 c_{NGI}^{(1)} (\not{p}_2 \gamma_\mu + \gamma_\mu \not{p}_1). \quad (2.8)$$

If we amputate (2.2) and use (2.4), (2.6) and (2.8) we get the off-shell improvement condition

$$\begin{aligned} \Lambda_\mu(p_2, p_1) &= \Lambda_{*,\mu}(p_2, p_1) + a g^3 c_{NGI}^{(1)} (\not{p}_2 \gamma_\mu + \gamma_\mu \not{p}_1) \\ &\quad - \frac{1}{2} a i \not{p}_2 \frac{\Sigma_W(p_2)}{\Sigma_p(p_2)} \Lambda_{*,\mu}(p_2, p_1) - \frac{1}{2} a i \Lambda_{*,\mu}(p_2, p_1) \not{p}_1 \frac{\Sigma_W(p_1)}{\Sigma_p(p_1)}, \end{aligned} \quad (2.9)$$

which should hold to order $\mathcal{O}(g^3)$ by determining $c_{NGI}^{(1)}$ and $c_{SW}^{(1)}$ correctly.

3. Calculation




Figure 1: One-loop diagrams contributing to the amputated quark-quark-gluon vertex

The diagrams contributing to the amputated one-loop three-point function are shown in Fig. 1. The calculation is performed combining symbolic and numerical methods. For the symbolic computation we use a *Mathematica* package that we developed for one-loop calculations in lattice perturbation theory (for a more detailed description see [11]). It is based on the infinite volume algorithm of Kawai et al. [10]. The analytic treatment has several advantages: one can extract the infrared singularities exactly and the results are given as functions of lattice integrals which can be determined with high precision. The disadvantage consists in very large expressions especially for the problem under consideration. In the analytic method the divergencies are isolated by differentiation with respect to external momenta. As can be seen in Fig. 1 diagrams (b) and (c) have two gluon propagators. So no parametrization can be chosen with only internal momentum flowing

through the gluon lines. Therefore at least one gluon propagator has to be differentiated. Looking at the full analytic form of the gluon propagator for improved gauge actions [13] one easily sees that huge analytic expression would arise. As discussed in [13] one can split the full gluon propagator $D_{\mu\nu}^{\text{improved}}(k)$

$$D_{\mu\nu}^{\text{improved}}(k) = D_{\mu\nu}^{\text{plaquette}}(k) + \Delta D_{\mu\nu}(k). \quad (3.1)$$

The diagrams with $D_{\mu\nu}^{\text{plaquette}}(k)$ only contain the logarithmic parts and are treated with the analytic *Mathematica* package. The diagrams with at least one $\Delta D_{\mu\nu}(k)$ are infrared finite and can be determined safely with pure numeric methods. We have written a C program with a Gauss-Legendre integration algorithm in four dimensions (for a description of the method see [11, 12]). We choose a sequence of small external momenta (p_1, p_2) and perform an extrapolation to vanishing momenta in order to extract the corresponding values. Additionally, we have written an independent FORTRAN code which computes the one-loop contributions for each diagram including the infrared logarithms. Results for both methods agree within accuracy.

The Feynman rules for non-smearred Symanzik gauge action have been summarized in [8]. For the stout smeared gauge links in the clover action the rules are given for the forward case by [4]. The corresponding Feynman rules needed for the quark-quark-gluon vertex are much more complicated and have been derived by the authors. They are too long as to be given in this proceedings [15].

The calculation has been done in Feynman gauge with Wilson parameter $r = 1$. All the one-loop coefficients are calculated at $c_{sw} = 1$ because $g^3 c_{sw} = g^3 + O(g^5)$.

4. Results

The anticipated general structure for the amputated three-point function in one-loop is

$$\begin{aligned} \Lambda_\mu(p_2, p_1) = & \Lambda_\mu^{\overline{MS}}(p_2, p_1) + A_{lat} i g^3 \gamma_\mu \\ & + B_{lat} \frac{a}{2} g^3 (\not{p}_2 \gamma_\mu + \gamma_\mu \not{p}_1) + C_{lat} \frac{ia}{2} g^3 \sigma_{\mu\alpha} q_\alpha \end{aligned} \quad (4.1)$$

$\Lambda_\mu^{\overline{MS}}(p_2, p_1)$ is the universal part of the three-point function independent of the chosen gauge action computed in the \overline{MS} -scheme

$$\begin{aligned} \Lambda_\mu^{\overline{MS}}(p_2, p_1) = & -i g \gamma_\mu - g \frac{a}{2} \mathbf{1} (p_{1,\mu} + p_{2,\mu}) - c_{sw} i g \frac{a}{2} \sigma_{\mu\alpha} q_\alpha \\ & + i g^3 F_{1,\mu}(p_1, p_2, q) + a g^3 F_{2,\mu}(p_1, p_2, q). \end{aligned} \quad (4.2)$$

$F_{1,\mu}(p_1, p_2, q)$ and $F_{2,\mu}(p_1, p_2, q)$ are complicated functions involving polylogarithms and logarithms. They will be given in [15]. The quantities A_{lat} , B_{lat} and C_{lat} are obtained as

$$\begin{aligned} A_{lat} = & C_F (0.03783 - 0.93653 \omega + 3.42833 \omega^2 + 0.01266 \log(a\mu)) \\ & + N_c (-0.02200 + 0.01266 \log(a\mu)), \\ B_{lat} = & C_F (0.03804 - 1.03749 \omega + 3.43791 \omega^2 + 0.02533 \log(a\mu)) \\ & + N_c (-0.02432 + 0.01925 \omega + 0.01266 \log(a\mu)), \\ C_{lat} = & C_F (0.11618 + 0.82813 \omega - 2.45508 \omega^2) \\ & + N_c (0.01215 + 0.01109 \omega - 0.30228 \omega^2), \end{aligned} \quad (4.3)$$

with $C_F = (N_c^2 - 1)/(2N_c)$ for $SU(N_c)$. As shown in (2.9) we need the self energy parts $\Sigma_p(p)$ and $\Sigma_W(p)$ as defined in (2.4) to solve the off-shell improvement condition

$$\begin{aligned}\Sigma_p(p) &= 1 - \frac{g^2 C_F}{16\pi^2} [\log(ap)^2 + \Sigma_1] , \\ \Sigma_W(p) &= 1 - \frac{g^2 C_F}{16\pi^2} [2 \log(ap)^2 + \Sigma_2] .\end{aligned}\quad (4.4)$$

It turns out that the self energy parts Σ_1 and Σ_2 contribute only to $c_{NGI}^{(1)}$. For the Symanzik gauge action we will present them in [15]. For the plaquette action we get

$$\begin{aligned}\Sigma_1^{plaq} &= 8.20627 - 196.44600 \omega + 739.68364 \omega^2 , \\ \Sigma_2^{plaq} &= 7.35794 - 208.58321 \omega + 711.56526 \omega^2 .\end{aligned}\quad (4.5)$$

We use (4.2) and (4.4) to construct the left hand side of (2.9) whereas (4.4) with (4.5) are inserted into the right hand side. In order to fulfill (2.9) we get the following improvement coefficients for the plaquette action

$$c_{NGI}^{(1,plaq)} = N_c (0.00143 - 0.01166 \omega) , \quad (4.6)$$

$$\begin{aligned}c_{SW}^{(1,plaq)} &= C_F (0.16764 + 1.07915 \omega - 3.68668 \omega^2) \\ &\quad + N_c (0.01502 + 0.00962 \omega - 0.28479 \omega^2) .\end{aligned}\quad (4.7)$$

For the Symanzik improved gauge action we find the improvement coefficient $c_{SW}^{(1)}$

$$\begin{aligned}c_{SW}^{(1)} &= C_F (0.11618 + 0.82813 \omega - 2.45508 \omega^2) \\ &\quad + N_c (0.01215 + 0.01109 \omega - 0.30228 \omega^2) .\end{aligned}\quad (4.8)$$

5. Mean field improvement

It is known that lattice artefacts make the perturbative expansion worse. One possible improvement procedure is to replace the naive coupling constant g by its mean field improved value $g_{MF} = g/u_0^2$ where u_0^4 is the average plaquette value for the corresponding gauge field action. By scaling all gauge links in the clover field strength $F_{\mu\nu}(n)$ in (1.2) by $1/u_0$ one obtains the mean field improved c_{SW} as

$$c_{SW}^{MF} = u_0^3 c_{SW} . \quad (5.1)$$

The perturbative expansion of u_0 is known to be

$$u_0 = 1 - \frac{g_{MF}^2 C_F}{16\pi^2} k_u , \quad (5.2)$$

where k_u for popular gauge actions are given in [13]. Therefore, the perturbative expression for the mean field improved c_{SW} is given by

$$c_{SW} = c_{SW}^{MF} u_0^{-3} = \frac{1}{u_0^3} \left(1 + g_{MF}^2 \left(c_{SW}^{(1)} - \frac{3C_F}{16\pi^2} k_u \right) + \mathcal{O}(g_{MF}^4) \right) = c_{SW}^{MF,p} + \mathcal{O}(g_{MF}^4) . \quad (5.3)$$

For the future simulations of the QCDSF collaboration we have the following numbers for the Symanzik action and 2+1 flavors

$$C_F = 4/3, \quad N_c = 3, \quad u_0^4 = 0.6065, \quad g_{MF}^2 = 1.71335, \quad k_u = 0.732524 \pi^2.$$

This gives the one-loop expression for c_{SW} parameter as

$$c_{SW} = 1 + g^2 (0.19136 + 1.13745 \omega - 4.18029 \omega^2) + \mathcal{O}(g^4), \quad (5.4)$$

$$\begin{aligned} c_{SW}^{MF,p} &= \frac{1}{u_0^3} (1 + g_{MF}^2 (0.19136 + 1.13745 \omega - 4.18029 \omega^2) - g_{MF}^2 0.18313) \\ &= 1.47557 + 2.83568 \omega - 10.42148 \omega^2 \end{aligned} \quad (5.5)$$

For no stout-smearing ($\omega = 0$) the result (5.4) has to be compared with the number given in [8]: $c_{SW}^{(1,AK)} = 0.19624449(1)$. The minor difference to our value $c_{SW}^{(1)} = 0.19136$ can possibly be related to an inaccuracy in our numerical integrations. In the simulation the stout parameter ω is chosen to be $\omega = 0.1$ leading to a mean field improved value $c_{SW}^{MF,p} = 1.65492$.

This investigation has been supported by the DFG under contract FOR 465 (Forschergruppe Gitter-Hadronen-Phänomenologie).

References

- [1] P. Weisz, *Nucl. Phys. B* **212** (1983) 1.
- [2] P. Weisz and R. Wohlert, *Nucl. Phys. B* **236** (1984) 397 [*Erratum-ibid. B* **247** (1984) 544].
- [3] B. Sheikholeslami and R. Wohlert, *Nucl. Phys. B* **259** (1985) 572.
- [4] S. Capitani, S. Dürr and C. Hoelbling, *JHEP* **0611** (2006) 028 [[arXiv:hep-lat/0607006](https://arxiv.org/abs/hep-lat/0607006)].
- [5] C. Morningstar and M. J. Peardon, *Phys. Rev. D* **69** (2004) 054501 [[arXiv:hep-lat/0311018](https://arxiv.org/abs/hep-lat/0311018)].
- [6] R. Wohlert, DESY preprint 87/069 (unpublished).
- [7] M. Lüscher and P. Weisz, *Nucl. Phys. B* **479** (1996) 429 [[arXiv:hep-lat/9606016](https://arxiv.org/abs/hep-lat/9606016)].
- [8] S. Aoki and Y. Kuramashi, *Phys. Rev. D* **68** (2003) 094019 [[arXiv:hep-lat/0306015](https://arxiv.org/abs/hep-lat/0306015)].
- [9] G. Martinelli, G. C. Rossi, C. T. Sachrajda, S. R. Sharpe, M. Talevi and M. Testa, *Nucl. Phys. B* **611** (2001) 311 [[arXiv:hep-lat/0106003](https://arxiv.org/abs/hep-lat/0106003)].
- [10] H. Kawai, R. Nakayama and K. Seo, *Nucl. Phys. B* **189** (1981) 40.
- [11] M. Göckeler, R. Horsley, E.-M. Ilgenfritz, H. Perlt, P. Rakow, G. Schierholz and A. Schiller, *Nucl. Phys. B* **472** (1996) 309 [[arXiv:hep-lat/9603006](https://arxiv.org/abs/hep-lat/9603006)].
- [12] M. Göckeler, R. Horsley, H. Perlt, P. E. L. Rakow, A. Schäfer, G. Schierholz and A. Schiller, *Nucl. Phys. B* **717** (2005) 304 [[arXiv:hep-lat/0410009](https://arxiv.org/abs/hep-lat/0410009)].
- [13] R. Horsley, H. Perlt, P. E. L. Rakow, G. Schierholz and A. Schiller [QCDSF Collaboration], *Nucl. Phys. B* **693** (2004) 3 [*Erratum-ibid. B* **713** (2005) 601] [[arXiv:hep-lat/0404007](https://arxiv.org/abs/hep-lat/0404007)].
- [14] S. Capitani, M. Göckeler, R. Horsley, H. Perlt, P. E. L. Rakow, G. Schierholz and A. Schiller, *Nucl. Phys. B* **593** (2001) 183 [[arXiv:hep-lat/0007004](https://arxiv.org/abs/hep-lat/0007004)].
- [15] R. Horsley, H. Perlt, P. E. L. Rakow, G. Schierholz and A. Schiller, in preparation.