

Higgs Searches at the Fermilab Tevatron $p\bar{p}$ Collider

Jianming Qian^{*†}

Department of Physics, University of Michigan, Ann Arbor, Michigan 48109, USA

E-mail: qianj@umich.edu

Searching for and potential discovery of Higgs boson(s) both within and beyond the standard model is perhaps the most visible physics goal of the current Fermilab Tevatron program. In this proceeding, recent results from both the CDF and DØ experiments based on analyses of datasets with integrated luminosities between 1.7 and 3.0 fb⁻¹ are summarized. The combined Tevatron cross section upper limits on the production of a standard model Higgs boson are fast approaching the expected standard model values for a wide mass range. Particularly, the Tevatron has now excluded a standard model Higgs boson with a mass of 170 GeV at 95% C.L.

2008 Physics at LHC

September 29 - 4 October 2008

Split, Croatia

^{*}Speaker.

[†]On behalf of the CDF and DØ Collaborations

1. Introduction

The standard model of particle physics has been remarkably successful in interpreting existing experimental data. However, the electroweak symmetry breaking of the model remains to be understood. In the model, the symmetry is postulated to be spontaneously broken through the so-called Higgs mechanism [1]. As a result, a neutral scalar fundamental particle called the Higgs boson is predicted to exist. Searching for this elusive particle has been a major undertaking of the experimental particle physics community over the last two decades.

With the exception of its mass, the couplings of the Higgs boson to fermions and gauge bosons are theoretically well known. Its mass, however, can be inferred from precision measurements of electroweak data, as the Higgs boson often appears in the loops of high-order electroweak corrections. A recent global fit to precision electroweak data yields a central value for the Higgs boson mass of 84_{-26}^{+34} GeV and a 95% C.L. upper bound of 154 GeV [2]. Direct searches at LEP have set a lower bound of 114.4 GeV on the mass, again at 95% C.L. [3].

The relatively low mass preferred by the fit is encouraging for the search for and potential discovery of the Higgs boson at the Fermilab Tevatron $p\bar{p}$ collider. At the Tevatron, the Higgs boson can be produced through the following four processes (ordered in decreasing cross section): (a) $gg \rightarrow H$, (b) $q\bar{q}' \rightarrow WH$, (c) $q\bar{q} \rightarrow ZH$ and (d) $qq \rightarrow qqH$. Since it couples to other particles in proportion to the mass of the particle, the Higgs boson decays predominantly to the heaviest particle kinematically allowed. Thus in the low mass region of $114 < m_H \lesssim 135$ GeV, Higgs decays mostly to $b\bar{b}$, while for masses above 135 GeV, $H \rightarrow WW^*$ decay dominates. For example, $Br(H \rightarrow b\bar{b}) = 73\%$ at 115 GeV and $Br(H \rightarrow WW^*) = 97\%$ at 170 GeV.

The minimal extension of the standard model, the minimal supersymmetric standard model (MSSM), predicts five Higgs bosons: two CP even (h , H) and one CP odd (A) neutral bosons as well as two charged bosons (H^+ and H^-) [4]. Two of the three neutral bosons are generally degenerate in mass and are often collectively denoted as ϕ . Two main production processes for ϕ at the Tevatron are $b\bar{b} \rightarrow \phi$ and $gg \rightarrow \phi$. Their cross sections are proportional to $\tan^2 \beta$, here $\tan \beta$ is the ratio of the vacuum expectation values of the two Higgs doublet fields in the model. Thus at high $\tan \beta$, the cross section could be large. Moreover, ϕ decays mostly to $b\bar{b}$ and $\tau\tau$. At $\tan \beta \sim 40$ ($\approx m_t/m_b$), for example, $Br(\phi \rightarrow b\bar{b}) \approx 90\%$ and $Br(\phi \rightarrow \tau\tau) \approx 10\%$.

The results summarized in this proceeding are from the CDF and DØ experiments at the Fermilab Tevatron $p\bar{p}$ collider at $\sqrt{s} = 1.96$ TeV. The Tevatron has been running well with over 5 fb^{-1} integrated luminosity delivered to each experiment. Analyzing these huge datasets takes time. Consequently the results described below are from data samples with luminosities between $1.7 - 3.0 \text{ fb}^{-1}$. The CDF and DØ detectors are designed to explore high p_T phenomena and are described in detail elsewhere [5].

The expected low rate of a potential Higgs signal poses a number of experimental challenges including, but not limited to, event selection efficiency, b -tagging performance and dijet mass resolution. Both the CDF and DØ experiments have invested considerable effort in all these areas. For example, CDF utilizes tracking and calorimetry information to significantly extend their muon identification capability, and uses their central tracking to supplement the calorimetry to improve the dijet mass resolution. DØ employs a neural network to combine several b -tagging algorithms to increase the b -tagging efficiency by over 30% for a fixed fake rate. The maturing

understanding of the datasets enables both collaborations to utilize advanced analysis techniques such as a neural network (NN), matrix-element (ME), and boosted decision trees (BDT) to exploit subtle differences between the signal and background processes, and hence greatly improve search sensitivities. NN is a pattern recognition paradigm that consists of a large number of interconnected processing elements trained to identify a specific pattern. The ME method on the other hand constructs likelihood discriminant using the matrix-elements (generally leading-order) of the underlying physical processes. And BDT is a classification algorithm that combines a set of weak classifiers into a powerful discriminant.

It is not possible to do justice to all the efforts that have gone into the Higgs searches in this short proceeding. Readers are encouraged to consult collaborations' web pages [6] for a full slate of the results as well as the latest updates.

2. Searches for a standard model Higgs boson

Though the $gg \rightarrow H$ production dominates over the entire mass region of interest at the Tevatron, the large $H \rightarrow b\bar{b}$ decay branching ratio at low masses renders it impractical for the search in this mass region, due to an overwhelming QCD $b\bar{b}$ background. Consequently searches at low masses are generally focused on Higgs productions in association with vector bosons such as WH and ZH . The decays of $W \rightarrow \ell\nu$ ($\ell = e, \mu$) and $Z \rightarrow \ell\ell, \nu\nu$ lead to distinct signatures for triggering and identification while $H \rightarrow b\bar{b}$ decay provides the benefit of the Higgs mass reconstruction. For a Higgs mass above 135 GeV, the $gg \rightarrow H$ production followed by the $H \rightarrow WW^* \rightarrow \ell\ell'\nu\nu$ decay offers the most promise for a potential Higgs discovery, as backgrounds are strongly suppressed by the presence of two high p_T leptons and a large missing transverse energy (E_T).

2.1 $WH \rightarrow \ell\nu b\bar{b}$

The $q\bar{q}' \rightarrow WH \rightarrow \ell\nu b\bar{b}$ final state is characterized by one high p_T lepton, a large E_T and a pair of b -jets. At 115 GeV, the expected cross section is about 30 fb. Main background processes include W +jets (Wbb, Wcc, Wjj), $t\bar{t}$, diboson (WW, WZ, ZZ) and multijet productions. The analyses generally begin with a pre-tagged sample of events with a W and two or more jets. The large statistics of this sample allows for the normalizations of different contributions. Since the W +jets cross section is not well known, its overall normalization is fixed to the data, after subtracting contributions from other processes such as $t\bar{t}$, diboson, multijets etc. The relative contributions of Wbb, Wcc and Wjj of the W +jets component are determined from Monte Carlo (MC). The multijet background is determined from the data. b -tagging algorithms are then applied to the data to yield samples with 1-tagged and 2-tagged events. MC background processes are subjected to tag-rate-functions of b -tagging algorithms to determine their contributions in these tagged samples. Advanced techniques are then employed to further separate the signal from backgrounds. After b -tagging, most information about Higgs is encoded in the dijet mass distribution. Thus good dijet mass resolution is critical for this analysis. Figure 1 (left) shows the dijet mass distribution of the 2-tagged sample from CDF. There is no apparent excess in the distribution. The corresponding BDT output including ME as an input is shown in Fig. 1 (right). Higgs candidates are expected to populate to the right of the distribution. An upper limit on the signal cross section can then be extracted by fitting the observed BDT distribution in the data to the sum of distributions expected

from background processes. CDF has analyzed a dataset of 2.7 fb^{-1} with both NN and BDT as the final discriminants and observed an upper cross section limit of 5.0 times the expected standard model value for $m_H = 115 \text{ GeV}$. DØ, on the other hand, has only analyzed 1.7 fb^{-1} of the data using a NN discriminant. The observed upper limit is therefore considerably weaker at 9.3 times the standard model value.




Figure 1: The invariant mass of the dijet (left) and the corresponding BDT outputs (right) of events with two b -tagged jets from CDF. For $m_H = 115 \text{ GeV}$, 2.0 ± 0.2 Higgs events are expected in this sample.

2.2 $ZH \rightarrow \nu\nu b\bar{b}$

The signature of this final state is two b -jets and a large E_T . The process has a cross section of $\sigma(ZH \rightarrow \nu\nu b\bar{b}) \approx 15 \text{ fb}$ for $m_H = 115 \text{ GeV}$. Before b -tagging, backgrounds are dominated by multijets with mis-measured E_T and W +light-jets as shown in Fig. 2 (left) from DØ. To reduce these backgrounds, two b -tagged jets are required. The remaining backgrounds are W + b/c -jets, dibosons and $t\bar{t}$ after the b -tagging. Though intended initially for $ZH \rightarrow \nu\nu b\bar{b}$, the analysis is found to be also sensitive to $WH \rightarrow \ell\nu b\bar{b}$ with an identified lepton (including τ). In fact, the WH and ZH contribute almost equally to the final signal yield. Without a high p_T lepton in the final state, the E_T becomes the main handle for the trigger and thus a relatively high E_T is required. The DØ analysis for example requires a E_T to be greater than 50 GeV. In addition, a set of topological cuts need to be applied to keep the multijet background under control. As in the case of $WH \rightarrow \ell\nu b\bar{b}$, the $ZH \rightarrow \nu\nu b\bar{b}$ production would manifest itself as a mass bump in the 2-tagged b -jets mass distribution. Figure 2 (right) compares the observed dijet mass distribution with those from background processes, along with the expected distribution from a potential Higgs signal. Both CDF and DØ have analyzed 2.1 fb^{-1} of the data. CDF uses a NN while DØ applies a BDT for the final signal-background separation. The observed upper limits for $m_H = 115 \text{ GeV}$ are 7.9 (CDF) and 7.5 (DØ) times the expected standard model cross section.

2.3 $ZH \rightarrow \ell\ell b\bar{b}$

The $q\bar{q} \rightarrow ZH \rightarrow \ell\ell b\bar{b}$ process in principle provides the cleanest Higgs signature since every final-state object can be reconstructed. Unfortunately even for a Higgs process, the rate is low ($\sim 5 \text{ fb}$ at $m_H = 115 \text{ GeV}$). Z boson production together with jets or another vector boson (W or Z) constitutes the majority of the background. $t\bar{t}$ background is important as well. Candidates are selected by first identifying two leptons from Z decays, followed by b -tagging, and finally by




Figure 2: The E_T distribution before b -tagging (left) and the dijet mass distribution (right) for events with two b -tagged jets. For $m_H = 115$ GeV, ~ 3.7 signal events are expected in the double-tagged sample.

applying advanced discriminants. CDF has analyzed 2.4 fb^{-1} of the data and used a 2-dimensional NN output to distinguish Z +jets and $t\bar{t}$ backgrounds from the signal. $D\emptyset$ has analyzed a dataset of similar size (2.3 fb^{-1}), but used a NN for the ee and a BDT for the $\mu\mu$ channel as the final signal-background discriminant. At $m_H = 115$ GeV, the observed upper limit in CDF is 11.6 times of the standard model value and the limit observed in $D\emptyset$ is 11.0.

2.4 $H \rightarrow WW^* \rightarrow \ell\ell' \nu\nu$

The $gg \rightarrow H \rightarrow WW^* \rightarrow \ell\ell' \nu\nu$ production and decay offers the best discovery potential at the Tevatron for a Higgs mass above 135 GeV. Compared with searches for a low mass Higgs, the direct reconstruction of Higgs mass in this final state is unfortunately infeasible due to the two undetected neutrinos. However, the final result below shows that this drawback is well compensated by the cleanliness of the event topology. The electroweak WW production, which has the identical final state as the signal, is the largest source of background. However, due to the scalar nature of the Higgs boson, WW from Higgs are expected to have different kinematics. Particularly the two leptons from the Higgs decay are expected to have a smaller average opening angle in the transverse plane than those from the electroweak WW process, a feature keenly exploited by both CDF and $D\emptyset$. Other backgrounds include $t\bar{t} \rightarrow WWbb$ and W +jets with one lepton from a mis-identified jet.




Figure 3: The E_T distribution of the $e\mu$ channel from $D\emptyset$ (left) and the combined NN output distribution from CDF (right).

Both CDF and $D\emptyset$ have analyzed 3.0 fb^{-1} of the data for this final state. CDF divides their analyses into 0-, 1- and 2-jet bins, but combines ee , $e\mu$ and $\mu\mu$ final states. This division is motivated by the fact that signal and background kinematics are expected to be different for different

POS(2008LHC)005

jet multiplicity. Moreover, it enables the 2-jet analysis to pick up contributions from other signal processes such as $ZH \rightarrow \ell\ell b\bar{b}$ and $qq \rightarrow qqH \rightarrow qqWW^* \rightarrow qq\ell\ell'v\nu$. Three different NNs, one for each jet bin, are trained. In contrast, the $D\emptyset$ analyses combine jet multiplicities but separate into ee , $e\mu$ and $\mu\mu$ channels. This approach follows naturally from the trigger and takes advantage of different signal-background ratios in the three channels. A NN is designed and trained for each channel. Efficient lepton identification and good E_T resolution are essential for this analysis. Figure 3 (left) compares the observed E_T distribution in the $e\mu$ channel in $D\emptyset$ with expected contributions from backgrounds while Fig. 3 (right) is the NN output distribution from CDF, combining three jet bins. For a mass of 165 GeV, both experiments expect to see about 16 signal events. The observed upper cross section limit for the $gg \rightarrow H$ production normalized to that of the standard model value is 1.6 for CDF and 2.0 for $D\emptyset$ at $m_H = 165$ GeV, by far the most stringent limits in all searches of a standard model Higgs boson.

2.5 Other searches

There are several other searches from both experiments that was not cover in the presentation, nor in this proceeding. The list includes $WH \rightarrow WWW^* \rightarrow \ell\ell\ell$ (CDF, $D\emptyset$), $WH/ZH \rightarrow qq\bar{b}\bar{b}$ (CDF), $t\bar{t}H \rightarrow t\bar{t}b\bar{b}$ ($D\emptyset$), $(H \rightarrow \tau\tau) + 2\text{-jets}$ (CDF) and $H \rightarrow \gamma\gamma$ ($D\emptyset$). Though less powerful than the searches presented above, these additional ones contribute significantly to the overall combination. Interested readers should see Ref. [6] for details.

2.6 Combined limits

The limits of individual search final states presented above are all above their respective standard model cross sections. To arrive at a Tevatron limit on the Higgs production for a given mass, results from CDF and $D\emptyset$ experiments are combined, taking into account correlations in systematic uncertainties. At the time of this presentation, the two collaborations have not updated the combination for the low mass searches. However, when they are done, the combined limit is expected to be significantly stringent than the individual limits presented above. For the high mass searches, the




Figure 4: Observed and expected 95% CL Tevatron combined upper limit on the Higgs cross section normalized to the standard model value as a function of the Higgs mass.

collaborations have recently combined the results of the $H \rightarrow WW^* \rightarrow \ell\ell'v\nu$ analyses. The combined Tevatron limit is shown in Fig. 4 as a function of the Higgs mass. It is encouraging to note

that the Tevatron upper cross section limit has reached the standard model value at $m_H = 170$ GeV, an indication of exciting results to come in the near future.

3. Searches for supersymmetric Higgs bosons

While searches for a standard model Higgs boson have the spotlight, both experiments have significant efforts in searches for the supersymmetric Higgs bosons as well. Ongoing analyses include $gb \rightarrow \phi b \rightarrow b\bar{b}b$ (CDF, DØ), $bb \rightarrow \phi \rightarrow \tau\tau$ (CDF, DØ) and $gb \rightarrow \phi b \rightarrow \tau\tau b$ (DØ).

For the same reason as $H \rightarrow b\bar{b}$ in the standard model case, inclusive search of $\phi \rightarrow b\bar{b}$ is impractical due to the large QCD $b\bar{b}$ background. The extra b -jet makes $b\phi \rightarrow b\bar{b}b$ final state manageable. Since there is neither a high p_T lepton nor a significant E_T , signal events have to be triggered by multijet triggers. QCD multijet production with b quarks constitutes the majority of the background and its contribution is determined from data. The analysis selects events with three b -tagged jets and searches for mass bumps of two b -tagged jets. Therefore good modeling of the dijet mass distribution is critical. As an example, Fig. 5 (left) shows the mass distribution of two b -tagged jets from CDF demonstrating a good agreement between the distribution of the signal sample and the sum of distributions from various background sources. Neither CDF nor DØ has seen any significant excess beyond the expected backgrounds in datasets of 1.8 and 2.6 fb^{-1} respectively. A significant region of the supersymmetry parameter space with a large $\tan\beta$ value is excluded.




Figure 5: Left: the mass distribution of two b -tagged jets in events with three b -tagged jets from CDF. Right: 95% C.L. limit on the $\tan\beta$ vs m_ϕ (M_A in the figure) plane from the DØ $\phi \rightarrow \tau\tau$ analysis.

$\phi \rightarrow \tau\tau$ is another important search final state at the Tevatron. With at least one τ decaying to an electron or a muon, the final state is sufficiently clean that the signal can be identified over backgrounds, comprised mostly of $Z \rightarrow \tau\tau$, W +jets and multijets. Efficient and clean identification of hadronically decaying τ 's is the key for the analysis. Both CDF and DØ have developed sophisticated τ algorithms based on large samples of $W \rightarrow \tau\nu$ and $Z \rightarrow \tau\tau$ events. The $\phi \rightarrow \tau\tau$ analysis searches for excesses in the visible mass distribution of the two τ 's. DØ has analyzed 2.2 fb^{-1} of the data and sees no evidence for $\phi \rightarrow \tau\tau$ production. The limit in one of the MSSM scenarios from this analysis is shown in Fig. 5 (right). Neither has any signal been seen by CDF

from a dataset of 2.0 fb^{-1} . DØ has also searched for the ϕ production in $b\phi \rightarrow b\tau\tau$ final state. The extra b -jet provides an additional tool for background rejection. This analysis yields a similar sensitivity as the inclusive $\phi \rightarrow \tau\tau$ analysis.

The combination of the CDF and DØ results is ongoing. We expect significant improvement in the limit once it is done.

4. Summary and Prospect

The quest for Higgs bosons in both standard and non-standard models is at the core of the Fermilab Tevatron Run II physics program. Both CDF and DØ have a comprehensive Higgs search effort. The sensitivities continue to improve at a rate faster than the rate of data accumulation. The search has reached a tipping point, particularly for a Higgs mass around 170 GeV. For the first time, direct searches at the Tevatron are contributing to our knowledge of the Higgs boson mass beyond those from LEP.

The Fermilab Tevatron collider continues to shatter luminosity records. It is scheduled to run in 2009. A decision, likely a positive one, on 2010 running is expected next spring. At the current pace, it is very likely that both experiments will have over 8 fb^{-1} of data recorded by the end of 2010. These huge datasets along with ever-improving analyses will undoubtedly lead to further improvements in sensitivities for the Higgs search. It appears that the Tevatron will be able to probe the standard model Higgs in a mass region between 145 – 185 GeV. The low mass region is as expected more challenging, just as it will be at the LHC. Nevertheless, with potential improvements the Tevatron could well have sensitivities in this region as well. For supersymmetric Higgs bosons, the Tevatron will continue to explore the parameter space which may lead to either evidence/discovery or large exclusion regions.

I conclude by saying that the best Tevatron Run II results are yet to come. I thank the CDF and DØ Collaborations for the opportunity to present these exciting results and the Fermilab Tevatron staff for delivering outstanding integrated luminosity! I wish to express my gratitude to G. Davies, M. Herndon, and D. Lincoln for commenting on this proceeding.

References

- [1] F. Englert and R. Brout, Phys. Rev. Lett. 13, 321 (1964); P.W. Higgs, Phys. Rev. Lett. 13, 508 (1964).
- [2] LEP Electroweak Working Group, <http://lepewwg.web.cern.ch/LEPEWWG>.
- [3] LEP Collaborations (ALEPH, DELPHI, L3 and OPAL), Phys. Lett. B 565, 61 (2003).
- [4] For a review, please see H.E. Haber and G.L Kane, Phys. Rep. 117, 75 (1985).
- [5] CDF Collaboration, Phys. Rev. D 71, 032001 (2005);
DØ Collaboration, Nucl. Instrum. Methods A 565, 463 (2006).
- [6] CDF Collaboration: <http://www-cdf.fnal.gov/physics/physics.html>;
DØ Collaboration: <http://www-d0.fnal.gov/Run2Physics/WWW/results.htm>