

## Transverse Momentum Distributions of Quarks in the Nucleon from Lattice QCD

---

**Bernhard U. Musch<sup>\*a</sup>, Philipp Hägler<sup>a</sup>, Andreas Schäfer<sup>b</sup>, Dru B. Renner<sup>c</sup>,  
John W. Negele<sup>d</sup>, LHPC (Lattice Hadron Physics Collaboration)**

<sup>a</sup> *Institut für Theoretische Physik T39, Physik-Department, Technische Universität München,  
James-Franck-Straße, D-85747 Garching, Germany*

<sup>b</sup> *Institut für Theoretische Physik, Universität Regensburg,  
D-93040 Regensburg, Germany*

<sup>c</sup> *Theory Group, Deutsches Elektronen-Synchrotron DESY,  
Platanenallee 6, D-15738 Zeuthen, Germany*

<sup>d</sup> *Center for Theoretical Physics, Massachusetts Institute of Technology,  
Cambridge, MA02139, USA*

*E-mail: [bmusch@ph.tum.de](mailto:bmusch@ph.tum.de)*

Transverse momentum dependent parton distribution functions (TMDPDFs) encode information about the intrinsic motion of quarks inside the nucleon. They are important non-perturbative ingredients in our understanding of, e.g., azimuthal asymmetries and other qualitative features in semi-inclusive deep inelastic scattering experiments. Here we present a first lattice approach to polarized and unpolarized transverse momentum dependent quark densities, based on MILC gauge configurations and propagators from LHPC. The operators we employ are non-local and consist of spatially separated quark creation and annihilation operators connected by a straight Wilson line. We briefly discuss their renormalization properties and our present strategies to remove linear divergences in  $a^{-1}$ .

*The XXVI International Symposium on Lattice Field Theory  
July 14 - 19, 2008  
Williamsburg, Virginia, USA*

---

<sup>\*</sup>Speaker.

## 1. Introduction

Semi-inclusive deeply inelastic scattering (SIDIS) experiments are sensitive to many correlations between the direction of parton and hadron spins and intrinsic transverse momenta. These have been parameterized in a systematic manner using transverse momentum dependent parton distribution functions (TMDPDFs), see [1]. TMDPDFs describe the distribution of partons carrying a longitudinal momentum fraction  $x$  and an intrinsic transverse momentum  $k_\perp$  in a hadron as illustrated in Fig. 1. Here we give an update of our effort [2] to develop techniques suitable for the calculation of moments of TMDPDFs on the lattice. Note that TMDPDFs are not to be confused with generalized parton distribution functions (GPDs), which provide probability distributions with respect to the impact parameter  $b_\perp$  rather than  $k_\perp$ . For an overview of recent hadron structure studies in lattice QCD, we refer to [3].

Fig. 2 illustrates the factorization of SIDIS into perturbative and non-perturbative parts. The lower blob represents the non-perturbative contribution of the nucleon and is described by

$$\Phi^{[\Gamma]}(x, k_\perp; P, S) \equiv \frac{1}{2} \int dk^- \int \frac{d^4 \ell}{(2\pi)^4} e^{-ik \cdot \ell} \langle P, S | \bar{q}(\ell) \Gamma \mathcal{U}_{\mathcal{C}(\ell, 0)} q(0) | P, S \rangle, \quad (1.1)$$

where  $|P, S\rangle$  represents a nucleon state of momentum  $P$  and spin  $S$ ,  $\Gamma$  is a Dirac matrix and  $k$  is the quark momentum, with  $k^+ = xP^+$ . The Wilson line  $\mathcal{U}_{\mathcal{C}(\ell, 0)}$  connecting the quark operators ensures gauge invariance. In SIDIS,  $\mathcal{U}_{\mathcal{C}(\ell, 0)} = \mathcal{U}_{[\ell, \ell + \infty \hat{n}_-]} \mathcal{U}_{[\ell + \infty \hat{n}_-, \infty \hat{n}_-]} \mathcal{U}_{[\infty \hat{n}_-, 0]}$  is a concatenation of three straight Wilson lines running to light cone infinity and back [4, 5].




Figure 1




Figure 2: Factorized tree level diagram of SIDIS.

Some examples of TMDPDFs are  $f_1$ ,  $g_{1L}$  and  $g_{1T}$ , defined via [1]

$$\Phi^{[\gamma^+]}(x, k_\perp; P, S) = f_1(x, k_\perp^2) \quad (1.2)$$

$$\Phi^{[\gamma^+ \gamma^5]}(x, k_\perp; P, S) = \frac{m_N}{P^+} S^+ g_{1L}(x, k_\perp^2) + \frac{k_\perp \cdot S_\perp}{m_N} g_{1T}(x, k_\perp^2). \quad (1.3)$$

## 2. TMDPDFs from the Euclidean lattice

In this exploratory study, we simplify the calculation and employ a single straight Wilson line  $\mathcal{U}_{\mathcal{C}(\ell, 0)} = \mathcal{U}_{[\ell, 0]}$  running from 0 to  $\ell$ . In this case, the matrix element appearing in eq. (1.1) can be

evaluated directly on the Euclidean lattice, as long as we set  $\ell^0 = 0$ . To facilitate the translation of the correlators evaluated on the lattice into TMDPDFs, we first parameterize the matrix element in terms of Lorentz-invariant amplitudes  $\tilde{A}_i(\ell^2, \ell \cdot P)$ , for example

$$\langle P, S | \bar{q}(\ell) \gamma_\mu \mathcal{U} q(0) | P, S \rangle = 4 \tilde{A}_2 P_\mu + 4i m_N^2 \tilde{A}_3 \ell_\mu, \quad (2.1)$$

$$\langle P, S | \bar{q}(\ell) \gamma_\mu \gamma^5 \mathcal{U} q(0) | P, S \rangle = -4 m_N \tilde{A}_6 S_\mu - 4i m_N \tilde{A}_7 P_\mu \ell \cdot S + 4 m_N^3 \tilde{A}_8 \ell_\mu \ell \cdot S. \quad (2.2)$$

The amplitudes  $\tilde{A}_i(\ell^2, \ell \cdot P)$  are extracted on the lattice and then Fourier transformed into TMDPDFs. Setting  $\ell^0 = 0$  translates into the constraints

$$\ell^2 \leq 0, \quad |\ell \cdot P| \leq |\vec{\ell}| |\vec{P}|, \quad (2.3)$$

which restrict the analysis of the full  $x$ - and  $k_\perp$ -dependence but are harmless if we are only interested in the first Mellin moment, i.e. if we integrate over  $x$ . For example, we obtain

$$f_1^{(1)}(k_\perp^2) \equiv \int_{-1}^1 dx f_1(x, k_\perp^2) = \int_0^\infty \frac{d(-\ell^2)}{2(2\pi)} J_0(\sqrt{-\ell^2} |k_\perp|) 2\tilde{A}_2(\ell^2, \ell \cdot P = 0), \quad (2.4)$$

$$g_{1T}^{(1)}(k_\perp^2) \equiv \int_{-1}^1 dx g_{1T}(x, k_\perp^2) = \int_0^\infty \frac{d(-\ell^2)}{2(2\pi)} \frac{J_1(\sqrt{-\ell^2} |k_\perp|)}{\sqrt{-\ell^2} |k_\perp|} (\ell^2 m_N^2) 2\tilde{A}_7(\ell^2, \ell \cdot P = 0), \quad (2.5)$$

where  $J_0$  and  $J_1$  are Bessel-functions.

### 3. Simulation technique and parameters

The matrix element  $\langle P, S | \bar{q}(\ell) \Gamma \mathcal{U}_{[\ell,0]} q(0) | P, S \rangle$  is evaluated using ratios of three- and two-point functions as described in Ref. [2]. The non-local operator  $\mathcal{O}^\Gamma(\ell) \equiv \bar{q}(\ell) \Gamma \mathcal{U}_{[\ell,0]} q(0)$  inserted in the three-point function contains the Wilson-line  $\mathcal{U}_{[\ell,0]}$ , which is implemented as a product of link variables. For oblique angles, we approximate a straight line by a step-like path as illustrated in Fig. 3.




Figure 3

For our studies we work with MILC gauge configurations [6] based on an AsqTad improved staggered quark action with 2+1 flavors on a  $20^3 \times 64$  lattice with a lattice spacing  $a \approx 0.12$  fm and a strange quark mass  $am_s = 0.050$ . We have used three different light quark masses,  $am_{u,d} = 0.020$  ( $m_\pi \approx 500$  MeV, 239 configurations),  $am_{u,d} = 0.030$  ( $m_\pi \approx 600$  MeV, 281 configurations) and the three-flavor degenerate case  $am_{u,d} = 0.050$  ( $m_\pi \approx 760$  MeV, 213 configurations).

The gauge configurations have been HYP smeared and bisected in the temporal direction to double statistics. We are using domain wall propagators and sequential propagators previously calculated by the LHPC collaboration on these configurations, with the valence quark mass tuned to match the staggered sea (see, e.g., [7]). The sequential propagators feature a source-sink separation of  $t_{\text{sink}} - t_{\text{source}} = 10$ , and are available for two lattice nucleon momenta  $\vec{P} = (0, 0, 0)$  and  $\vec{P} = (-1, 0, 0)$ , the latter corresponding to 500 MeV in physical units. We neglect contributions from disconnected diagrams.




Figure 4: (a) Amplitude  $2 \operatorname{Re} \tilde{A}_2(\ell^2, \ell \cdot P = 0)$  for up quarks. The continuous curves are fits of the form  $C_1 \exp(-\ell^2/\sigma_1^2) + C_2 \exp(-\ell^2/\sigma_2^2)$ . We have used three different sets of renormalization constants: For the upper curve, we have determined  $a\delta m$  from 1-loop perturbation theory, for the two lower curves we have employed two variants of the taxi driver method. (b) The same for the amplitude  $-2 \operatorname{Re} \tilde{A}_7(\ell^2, \ell \cdot P = 0)$ .

#### 4. Renormalization of the Wilson Lines

The Wilson line  $\mathcal{U}_{[\ell,0]}$  in our non-local operator gives rise to a linear divergence, which has to be removed by a renormalization constant  $\delta m$  proportional to the cutoff, given by  $a^{-1}$  on the lattice. Refs. [8, 9] show within continuum theory that the renormalized operator is of the form

$$\mathcal{O}_{\text{ren}}^\Gamma(\ell) = Z^{-1} \exp(-\delta m L) \mathcal{O}^\Gamma(\ell). \quad (4.1)$$

Here  $Z^{-1}$  subsumes renormalization factors associated with divergences at the end points, and  $L$  is the total length of the smooth Wilson line.

In lattice QCD, the linear divergence has been a long standing issue in the context of heavy quark propagators [10]. We have calculated  $a\delta m$  for link paths on the axes in leading order perturbation theory, adapting the procedure in Refs. [11, 12] to our action according to Refs. [13, 14].

However, perturbation theory is not expected to give accurate results. Therefore we have also sought to determine  $a\delta m$  non-perturbatively with our “taxi driver method”, which is based on the assumption that on the lattice,  $L$  in eq. (4.1) is given by the total number of link variables, and thus allows us to deduce the renormalization constants from the comparison of straight and step-like link paths. There are two variants of this method, one based on data from Wilson lines in a Landau gauge fixed ensemble, and one based on Wilson loops. For the moment, we assume that the quark mass dependence is weak, and use the constants determined on the three-flavor degenerate lattice also for the lighter quark masses. We are currently testing the validity of the taxi driver approach and alternative methods on several different lattice spacings.

#### 5. Results with Preliminary Renormalization

Here we present some results obtained by applying the techniques sketched above. Note that at the present stage we regard our renormalization procedure still as preliminary. In the following, we

label our distributions “sW” to indicate that they are based on straight Wilson lines and are therefore not strictly identical to the TMDPDFs defined and used in the literature and for the description of, e.g., SIDIS.

In Fig. 4a we display results for  $2\tilde{A}_2(\ell^2, \ell \cdot P = 0)$ . They are obtained from a three-point function with the operator  $\mathcal{O}^A(\ell)$ , where the Wilson line is renormalized using different approaches as described above. The overall normalization can be obtained requiring charge conservation,  $2\tilde{A}_2^{u-d}(0,0) = 1$ . Applying eq. (2.4) to the fit results, we get  $f_1^{(1)\text{sW}}(k_\perp^2)$  as plotted in Fig. 5a, which is interpreted as the unpolarized distribution of quarks in the unpolarized nucleon in Fig. 6a. In the unpolarized channels, the nucleon looks axially symmetric. For the large pion masses currently analyzed, the quark mass dependence of the width of this distribution appears to be rather weak, see Fig. 5c.

The axial symmetry is broken in the polarized case. As an example, we can consider the distribution of quarks with positive helicity  $\lambda = +1$ , corresponding to an operator  $\mathcal{O}^\Gamma(\ell)$  with  $\Gamma = \gamma^+ \frac{1}{2}(1 + \gamma^5)$  in a nucleon transversely polarized with transverse spin  $S_\perp$ . Using eq. (1.2,1.3)

$$\rho(k_\perp, S_\perp) \equiv \int_{-1}^1 dx \Phi^{[\gamma^+(1+\gamma^5)/2]}(x, k_\perp; P, S) = \frac{1}{2} \left( f_1^{(1)}(k_\perp^2) + \frac{k_\perp \cdot S_\perp}{m_N} g_{1T}^{(1)}(k_\perp^2) \right). \quad (5.1)$$

This density is axially asymmetric due to the contribution from the TMDPDF  $g_{1T}$ , which contains information about the correlation of spins and momenta of the form  $\vec{s} \cdot \vec{P} \cdot k \cdot S \sim \lambda k_\perp \cdot S_\perp$ . According to eq. (2.5), we obtain  $g_{1T}^{(1)\text{sW}}(k_\perp^2)$  from the amplitude  $\tilde{A}_7$ , as shown in the case of up quarks in Fig. 4b and 5b. For down quarks, the results are of opposite sign and smaller by a factor of about 1/6. In terms of the density  $\rho(k_\perp, S_\perp)$ , we find that quarks of a specific polarization can have a non-vanishing average transverse momentum

$$\langle k_\perp \rangle = \frac{\int d^2 k_\perp k_\perp \rho(k_\perp, S_\perp)}{\int d^2 k_\perp \rho(k_\perp, S_\perp)} \quad (5.2)$$




Figure 5: (a)  $f_1^{(1)\text{sW}}(k_\perp^2)$  for up quarks as obtained from the Fourier transform of the fits in Fig. 4a. (b)  $\frac{|k_\perp|}{m_N} g_{1T}^{(1)\text{sW}}(k_\perp^2)$  for up quarks as obtained from the fits in Fig. 4b. (c) Linear chiral extrapolation of the root mean squared transverse momentum  $\langle k_\perp^2 \rangle^{1/2}$  from  $f_1^{(1)\text{sW}}(k_\perp^2)$  for up quarks minus down quarks.

and that it is opposite in sign for up- and down quarks, see Figs. 6b and 6c. That such deformed quark densities are to be expected has been mentioned, e.g., in Ref. [15]. Similar deformations have also been observed in the framework of GPDs in [16, 17].




Figure 6: Quark density plots. Here the renormalization constants have been chosen according to the taxi driver method based on Wilson lines.

(a)  $f_1^{(1)SW}(k_\perp^2) = \int dx \Phi^{[\gamma^+]}(x, k_\perp)$  for up quarks at  $m_\pi = 500 \text{ MeV}$ . We interpret this as the charge density of up quarks in the nucleon in the transverse momentum plane.

(b) Density  $\rho(k_\perp, S_\perp)$  of up quarks with positive helicity  $\lambda = +1$  (i.e. with spin pointing in  $z$ -direction) in a nucleon polarized in transverse  $x$ -direction  $S_\perp = (1, 0)$ , evaluated at a pion mass  $m_\pi = 500 \text{ MeV}$ . The distribution features an average transverse momentum shift  $\langle k_x \rangle = (67 \pm 5_{\text{stat}} \pm 3_{\text{renorm.}}) \text{ MeV}$ , where the uncertainty from renormalization has been estimated from the comparison of the three different sets of renormalization constants employed in Figs. 4 and 5.

(c) Same as in (b) but for down quarks. The average transverse momentum shift  $\langle k_x \rangle = (-24 \pm 5_{\text{stat}} \pm 3_{\text{renorm.}}) \text{ MeV}$  has the opposite sign as for up quarks.

(d) Same as in (b) for up quarks minus down quarks. The deformation appears amplified. Note that this is not a density and not necessarily positive.

## 6. Conclusions and Outlook

Presently, we employ a straight Wilson line between the quark fields in the definition of TMD-PDFs. We have shown first preliminary results from lattice QCD for the TMDPDFs  $f_1^{sW}$  and  $g_{1T}^{sW}$  as a function of transverse momentum. We find that densities of longitudinally polarized quarks in a transversely polarized proton are deformed.

Concerning our renormalization procedure, further investigations are in progress. Furthermore, we would like to extend our work towards non-straight Wilson lines, similar to those appearing in the definition of TMDPDFs for experimental processes such as SIDIS.

## Acknowledgments

Thanks are due to Vladimir Braun, Gunnar Bali and Meinulf Gökeler for very helpful discussions and to the members of the LHPC collaboration for providing propagators and technical expertise. B. M. and Ph. H. acknowledge support by the DFG Emmy Noether-program and the Excellence Cluster Universe at the TU München, A. S. acknowledges support by BMBF. This work was supported in part by funds provided by the U.S. Department of Energy under grant DE-FG02-94ER40818.

## References

- [1] P. J. Mulders and R. D. Tangerman, *Nucl. Phys.* **B461** (1996) 197–237.
- [2] B. U. Musch, P. Hägler, A. Schäfer, M. Gökeler, D. B. Renner, J. W. Negele, and LHPC collaboration, *PoS LAT2007* (2007) 155.
- [3] J. Zanotti, *PoS LAT2008* (2008) 007.
- [4] A. V. Belitsky, X. Ji, and F. Yuan, *Nucl. Phys.* **B656** (2003) 165–198.
- [5] D. Boer, P. J. Mulders, and F. Pijlman, *Nucl. Phys.* **B667** (2003) 201–241.
- [6] C. W. Bernard *et al.*, *Phys. Rev.* **D64** (2001) 054506.
- [7] P. Hägler *et al.*, LHPC collaboration, *Phys. Rev.* **D77** (2008) 094502.
- [8] N. S. Craigie and H. Dorn, *Nucl. Phys.* **B185** (1981) 204.
- [9] H. Dorn, *Fortsch. Phys.* **34** (1986) 11–56.
- [10] L. Maiani, G. Martinelli, and C. T. Sachrajda, *Nucl. Phys.* **B368** (1992) 281–292.
- [11] E. Eichten and B. R. Hill, *Phys. Lett.* **B240** (1990) 193.
- [12] P. Boucaud, L. C. Lung, and O. Pene, *Phys. Rev.* **D40** (1989) 1529.
- [13] B. Bistrovic. PhD thesis, Massachusetts Institute of Technology, Dept. of Physics, 2005. <http://hdl.handle.net/1721.1/32304>.
- [14] T. A. DeGrand, *Phys. Rev.* **D67** (2003) 014507.
- [15] G. A. Miller, *Phys. Rev.* **C76** (2007) 065209.
- [16] M. Diehl and P. Hägler, *Eur. Phys. J.* **C44** (2005) 87–101.
- [17] M. Gökeler *et al.*, QCDSF collaboration, *Phys. Rev. Lett.* **98** (2007) 222001.