

The rapidity dependence of the proton-to-pion ratio in Au+Au and p+p collisions at $\sqrt{s_{NN}} = 62.4$ and 200 GeV

P. Staszal^{*†}

Smoluchowski Inst. of Physics, Jagiellonian University, ul. Reymonta 4, 30-059 Kraków, Poland

E-mail: ufstasze@if.uj.edu.pl

The proton-to-pion yield ratios measured in the BRAHMS experiment for Au+Au and p+p collisions at $\sqrt{s_{NN}} = 62.4$ GeV and $\sqrt{s_{NN}} = 200$ GeV are presented as a function of transverse momentum and collision centrality within the pseudorapidity range $0 \leq \eta \leq 3.8$. A strong pseudorapidity dependence of these ratios is observed. In Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, $\eta \approx 2.2$, and at $\sqrt{s_{NN}} = 62.4$ GeV, $\eta = 0$, the bulk medium can be characterized by the common value of $\mu_B \approx 65$ MeV. The $p/\pi^+(p_T)$ ratios measured for these two selections display a striking agreement in the p_T range covered (up to 2.2 GeV/c). At collision energy of 62.4 GeV and forward pseudorapidity we found a crossing point of p/π^+ ratios measured in central and semi-peripheral Au+Au and in p+p reactions. The crossing occurs in the narrow η bin around value of 3.2, simultaneously in the whole covered p_T range ($0.3 \text{ GeV}/c < p_T < 1.8 \text{ GeV}/c$).

*European Physical Society Europhysics Conference on High Energy Physics, EPS-HEP 2009,
July 16 - 22 2009
Krakow, Poland*

^{*}Speaker.

[†]for the BRAHMS collaboration.

The measured p_T dependence of the baryon-to-meson ratio appears to be related to modifications in the hadronization mechanisms as it happens in a partonic medium. It was pointed out that the baryon-to-meson ratio p_T dependence should be sensitive to the hadronization scenario due to the different quark content of baryons and mesons and/or to radial flow of the bulk medium because of significant differences in baryon and meson masses. Both flow and medium quark coalescence are expected to enhance protons over pions at intermediate p_T .

The PHENIX \bar{p}/π^- data at mid-rapidity is well described by the Greco, Ko, and Levai quark coalescence model where the introduced coalescence involves partons from the medium (thermal) and partons from mini-jets [1]. The Hwa and Yang quark recombination model is also successful in describing BRAHMS and PHENIX mid-rapidity data for p/π^+ [2]. On the other hand, the comparison with the hydrodynamical model shows that hydro-flow cannot itself account for the large observed ratio above ≈ 3 GeV/c and that the model overpredicts the data at low p_T [3].

These results support the view of a hadronization process driven by parton recombination with negligible final state interactions between produced hadrons. However, at large μ_B thus large rapidities, the picture, suggested by mid-rapidity measurements, might be contaminated by final state hadron interactions leading to a transition from the parton recombination scheme to a hydrodynamical description that has a common velocity field for baryons and mesons [4, 5].

The setup of the BRAHMS experiment is described in details in [6]. Here we just point out that the arrangement of BRAHMS spectrometers, namely of the Mid-Rapidity Spectrometer (MRS) and the Forward Spectrometer (FS), makes it possible to measure identified particle spectra over a pseudorapidity interval from $\eta = 0$ to $\eta = 3.8$. Particle identification in the FS is provided by TOF measurements for low and medium particle momenta. High momentum particles are identified using a Ring Imaging Cherenkov detector (RICH) [7].

The data analysis reported in this proceedings utilizes the feature of the same pion and proton acceptance in the η versus p_T space in the same real time measurement. For a given η - p_T bin the p/π ratios are calculated on a setting by setting basis. In order to avoid mixing different PID techniques, which usually lead to different systematic uncertainties, the ratios are calculated separately for the TOF PID and the RICH PID. In this way all factors such as acceptance corrections, tracking efficiencies, trigger normalization and bias related to the centrality cut cancel out in the ratio. The remaining species dependent corrections are: (i) decay in flight, interaction with the beam pipe and other detector material, and (ii) the PID efficiency correction. The corrections for (ii) are determined from the single particle response of pions and protons with energy in a realistic GEANT [9] model description of the BRAHMS experimental setup. We estimate that the overall systematic uncertainty related to this correction is at the level of 2%. The PID procedure used in the analysis leads to a relatively clean sample of pions with some contamination by kaons having spurious rings associated in the RICH counter. Together with the kaon - proton overlap at larger momenta, this contamination effect is a source of systematic errors that are represented by shadow boxes in Figure 1.

Figure 1 (left) shows p/π^+ ratios obtained for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV for two centrality selections of events, namely, for centrality 0 – 10% (solid dots) and 40 – 80% (open squares) The shaded boxes plotted for the most central events only, represent the systematic uncertainties discussed in the previous section. The ratios extracted from p+p data at the same energy are plotted for comparison (solid stars). The p_T coverage depends on the pseudorapidity bins and




Figure 1: Centrality dependent p/π^+ (left) and \bar{p}/π^- (right) ratio for Au+Au system colliding at $\sqrt{s_{NN}} = 200$ GeV for central (0 – 10%) and semi-peripheral (40 – 80%) reactions in comparison with p+p collision data at the same energy. The vertical bars represent the statistical errors and the shaded boxes (plotted only for central Au+Au) show the systematic uncertainties.

extends up to $p_T = 4$ GeV/c for $\eta = 2.6$ and 3.1. At low p_T (< 1.5 GeV/c) the p/π^+ ratios exhibit a rising trend with a weak dependence on centrality. The dependency on centrality begins above 1.5 GeV/c. The ratios appear to reach a maximum value at p_T around 2.5 GeV/c (whenever there is enough p_T coverage). The maxima of the ratios increase with the level of centrality and at $\eta = 3.1$, are equal to about 2.5 and 1.5 for the 0 – 10% and 40 – 80% centrality bins, respectively. The p+p ratios are consistent with Au+Au data at low p_T and begin to deviate significantly above $p_T = 1$ GeV/c. At $\eta = 3.1$ a maximum value of the ratio of 0.55 is reached in p+p collisions which is a factor of 4.5 smaller than that observed for central Au+Au reactions.

The values of the \bar{p}/π^- ratios plotted in Fig. 1 (right) are significantly lower than the p/π^+ ratios (note the difference in the vertical scale), however, the centrality dependence shows the same features as those observed in the p/π^+ ratios, namely, that the ratios for different centrality classes are consistent with each other up to $p_T \approx 1.2$ GeV/c and a strong dependence on centrality appears at larger transverse momenta reaching a maximum at similar p_T as the positive particles. Looking at the p+p data alone, one can note the difference in shape between the p/π^+ and \bar{p}/π^- ratios: a clear shift of the \bar{p}/π^- peaks towards lower p_T , as well as a much broader p/π^+ peaks. These large difference between the Au+Au and p+p both in shape and overall magnitude may reflect significant medium effects in Au+Au at $\sqrt{s_{NN}} = 200$ GeV in the pseudorapidity intervals covered.

Figure 2 presents p/π^+ measured in Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV and $\eta = 0.0$ shown with open red (on-line) triangles and the same ratio measured in Au+Au reactions at $\sqrt{s_{NN}} = 200$ GeV and $\eta = 2.2$ shown with the black triangles. The pseudorapidity intervals selected for this comparison corresponds to the ranges where the observed \bar{p}/p ratios are similar, of approximately 0.45. The similarity of proton-to-pion ratios for these selected heavy ion collisions suggests that the baryon and meson production at the p_T interval studied (up to 2 GeV/c) is dominated by medium effects and is determined by the bulk medium properties. Figure 3 compares the p/π^+ ratio from




Figure 2: p/π^+ ratio in Au+Au collisions for $\eta = 0.0$ at $\sqrt{s_{NN}} = 62.4$ GeV and in Au+Au reactions for $\eta = 2.2$ at $\sqrt{s_{NN}} = 200$ GeV.




Figure 3: p/π^+ ratio from p+p and Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV and $\eta \approx 3.2$.

p+p and Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV and $\eta \approx 3.2$ ($\mu_B \approx 250$ MeV, [10]). There is remarkably little difference in the p/π^+ ratios over a very wide range of the colliding system volume, namely, from p+p reactions up to central Au+Au collisions. It should be noted, that (what is not shown) at $\eta = 2.67$ the p/π^+ ratio from central Au+Au reactions is enhanced in respect to p+p collisions by a factor of 1.6, whereas at $\eta = 3.5$ the situation is reversed, namely, p/π^+ ratio in p+p exceeds that measured in central Au+Au by a factor of about 1.4. This indicates that the consistency observed at $\eta \approx 3.2$ is a result of simultaneous crossing of the ratios for different systems (from p+p up to central Au+Au) at this particular pseudorapidity bin.

We presented the (p_T) dependence of the p/π ratios measured in Au+Au and p+p collisions at energies 62.4 and 200 GeV as a function of pseudorapidity and collision centrality (Au+Au). For Au+Au and p+p reactions at $\sqrt{s_{NN}} = 200$ GeV the p/π^+ and \bar{p}/π^- ratios show noticeable dependency on centrality at intermediate p_T with a rising trend from p+p to central Au+Au collisions. We have shown that p/π^+ ratios are remarkably similar for central Au+Au at $\sqrt{s_{NN}} = 200$ GeV, $\eta \approx 2.2$ and central Au+Au at $\sqrt{s_{NN}} = 62.4$ GeV, $\eta \approx 0$, where the bulk medium is characterized by the same value of \bar{p}/p . This observation, together with the observed centrality dependence suggests, that at these energies and pseudorapidity intervals, particle production at intermediate p_T is governed by the size and chemical properties of the created medium. Finally, the Au+Au and p+p measurements at $\sqrt{s_{NN}} = 62.4$ GeV shows that the p/π^+ ratios for p+p and for all analysed Au+Au centralities cross simultaneously at the same η value (≈ 3.2) and are consistent with each other in the covered p_T range i.e. from 0.3 GeV/c up to 1.8 GeV/c.

References

- [1] V. Greco, C.M. Ko, and P. Levai, Phys. Rev. Lett. **90**, 022302 (2003).
- [2] R. C. Hwa, and C. B. Yang, Phys. Rev. C **67**, 034902 (2003).
- [3] E. J. Kim (BRAHMS Collaboration), Nucl. Phys. A **774**, 493 (2006).
- [4] T. Hirano, and Y. Nara, Nucl. Phys. A **743**, 305 (2004).

- [5] W. Broniowski, and W. Florkowski, Phys. Rev. Lett. **87**, 272302 (2001).
- [6] M. Adamczyk et. al., BRAHMS Collaboration, Nucl. Instr. Meth. **A 499**, 437 (2003).
- [7] R. Debbel et. al., Nucl. Instr. Meth. **A 570** 216 (2007).
- [8] M. Gyulassy and X. N. Wang, Comput. Phys. Commun. **83** 307 (1994).
- [9] "GEANT: Detector Description and Simulation Tool", wwwinfo.cern.ch/asdoc/geant_html3/
- [10] I.C. Arsene et. al., for BRAHMS Collaboration, Int. Jour. of Mod. Phys. **E 16**, 2035 (2007).