

Unitarity-Cuts, Stokes' Theorem and Berry's Phase

Pierpaolo Mastrolia^{*†}

Centro Studi e Ricerche "E. Fermi", Piazza del Viminale 1, I-00184, Roma, Italy

Dipartimento di Fisica, Università di Salerno, Via Ponte don Melillo, I 84084 Fisciano, Italy

Theory Department, CERN, CH-1211 Geneva 23, Switzerland

E-mail: pierpaolo.mastrolia@cern.ch

Two-particle unitarity-cuts of scattering amplitudes can be efficiently computed by applying Stokes' Theorem, in the fashion of the Generalised Cauchy Theorem. Consequently, the Optical Theorem can be related to the Berry Phase, showing how the imaginary part of arbitrary one-loop Feynman amplitudes can be interpreted as the flux of a complex 2-form.

*RADCOR 2009 - 9th International Symposium on Radiative Corrections (Applications of Quantum Field Theory to Phenomenology),
October 25 - 30 2009
Ascona, Switzerland*

^{*}Speaker.

[†]In memory of my friend Thomas (Binoth), who reacted with his characteristic enthusiasm when I first shew him the results now collected in this manuscripts. I acknowledge Tania Robens for clarifying discussions. My participation to the conference was supported by HepTools.

1. Introduction

Unitarity and geometric phases are two ubiquitous properties of physical systems. The Berry phase is the phase acquired by a system when it is subjected to a cyclic evolution, resulting only from the geometrical properties of the path traversed in the parameter space because of anholonomy [1, 2]. Unitarity represents the probability conservation in particle scattering processes described by the unitary *scattering operator*, S . The relation, $S = 1 + iT$, between the S -operator and the *transition operator*, T , leads to the Optical Theorem, $-i(T - T^\dagger) = T^\dagger T$. The matrix elements of this equation between initial and final states are expressed, in perturbation theory, in terms of Feynman diagrams. The evaluation of the right hand side requires the insertion of a complete set of intermediate states. Therefore, since $-i(T - T^\dagger) = 2 \text{Im}T$, the Optical Theorem yields the computation of the imaginary part of Feynman integrals from a sum of contributions from all possible intermediate states.

The Cutkosky-Veltman rules, implementing the unitarity conditions, allow the calculation of the discontinuity across a branch cut of an arbitrary Feynman amplitude, which corresponds to its imaginary part [3]. Accordingly, the imaginary part of a given Feynman integral can be computed by evaluating the phase-space integral obtained by cutting two internal particles, which amounts to applying the on-shell conditions and replacing their propagators by the corresponding δ -function, $(p^2 - m^2 + i0)^{-1} \rightarrow (2\pi i) \delta^{(+)}(p^2 - m^2)$. In later studies the problem of finding the discontinuity of a Feynman integral associated to a singularity was addressed in the language of homology theory and differential forms [4].

More recently multi-particle cuts have been combined with the use of complex momenta for on-shell internal particles into very efficient techniques, by-now known as *unitarity-based methods*, to compute scattering amplitudes for arbitrary processes. These methods exploit two general properties of scattering amplitudes like analyticity, granting that amplitudes are determined by their own singularity-structure, and unitarity, granting that the residues at the singular points factorize into products of simpler amplitudes. Unitarity and analyticity become tools for the quantitative determination of one-loop amplitudes [5] when merged with the existence of an underlying representation of amplitudes as a combination of basic scalar one-loop functions [6]. These functions, known as Master Integrals (MI's), are n -point one-loop integrals, I_n ($1 \leq n \leq 4$), with trivial numerator, equal to 1, characterised by external momenta and internal masses present in the denominator. In general, the fulfillment of multiple-cut conditions requires loop momenta with complex components. Since the loop momentum has four components, the effect of the cut-conditions is to fix some of them according to the number of the cuts. Any *quadruple-cut* [11] fixes the loop-momentum completely, yielding the algebraic determination of the coefficients of I_n , ($n \geq 4$); the coefficient of 3-point functions, I_3 , are extracted from *triple-cut* [12, 13, 14, 15, 16, 17]; the evaluation of *double-cut* [18, 19, 20, 14, 21, 16, 22, 23, 17] is necessary for extracting the coefficient of 2-point functions, I_2 ; and finally, in processes involving massive particles, the coefficients of 1-point functions, I_1 , are detected by *single-cut* [16, 24, 25]. In cases where fewer than four denominators are cut, the loop momentum is not frozen: the free-components are left over as phase-space integration variables.

In [26, 27], I introduced a novel efficient method for the analytic evaluation of the coefficients of one-loop 2-point functions *via* double-cuts. Spun-off from the spinor-integration technique [18, 19, 20], that method is an application of Stokes' Theorem. Due to a special decomposition of the

loop-momentum, the double-cut phase-space integral is written as parametric integration of rational function in two complex-conjugated variables. By applying Stokes' Theorem, the integration is carried on in two simple steps: an indefinite integration in one variable, followed by Cauchy's Residue Theorem in the conjugated one.

The coefficients of the 2-point scalar functions, being proportional to the rational term of the double-cut, can be directly extracted from the indefinite integration by Hermite Polynomial Reduction.

2. Double-Cut

The two-particle Lorentz invariant phase-space (LIPS) in the K^2 -channel is defined as,

$$\int d^4\Phi = \int d^4\ell_1 \delta^{(+)}(\ell_1^2 - m_1^2) \delta^{(+)}((\ell_1 - K)^2 - m_2^2), \quad (2.1)$$

where K^μ is the total momentum across the cut. We introduce a suitable parametrization for ℓ_1^μ [26, 20], in terms of four massless momenta, which is a solution of the two on-shell conditions, $\ell_1^2 = m_1^2$ and $(\ell_1 - K)^2 = m_2^2$,

$$\ell_1^\mu = \frac{1 - 2\rho}{1 + z\bar{z}} \left(p^\mu + z\bar{z} q^\mu + z \varepsilon_+^\mu + \bar{z} \varepsilon_-^\mu \right) + \rho K^\mu, \quad (2.2)$$

where p_μ and q_μ are two massless momenta with the requirements,

$$p_\mu + q_\mu = K_\mu, \quad p^2 = q^2 = 0, \quad 2 p \cdot q = 2 p \cdot K = 2 q \cdot K \equiv K^2; \quad (2.3)$$

the vectors ε_+^μ and ε_-^μ are orthogonal to both p^μ and q^μ , with the following properties¹,

$$\varepsilon_+^2 = \varepsilon_-^2 = 0 = \varepsilon_\pm \cdot p = \varepsilon_\pm \cdot q, \quad 2 \varepsilon_+ \cdot \varepsilon_- = -K^2. \quad (2.4)$$

The pseudo-threshold $\rho = (K^2 + m_1^2 - m_2^2 - \sqrt{\lambda}) / (2K^2)$, with $\lambda = (K^2)^2 + (m_1^2)^2 + (m_2^2)^2 - 2K^2 m_1^2 - 2K^2 m_2^2 - 2m_1^2 m_2^2$, depends only on the kinematics. The complex conjugated variables z and \bar{z} parametrize the degrees of freedom left over by the cut-conditions. Because of (2.2), the LIPS in (2.1) reduces to the remarkable expression,

$$\int d^4\Phi = (1 - 2\rho) \iint \frac{dz \wedge d\bar{z}}{(1 + z\bar{z})^2}. \quad (2.5)$$

The double-cut of a generic n -point amplitude in the K^2 -channel is defined as

$$\Delta \equiv \int d^4\Phi A_L^{\text{tree}}(\ell_1) A_R^{\text{tree}}(\ell_1), \quad (2.6)$$

where $A_{L,R}^{\text{tree}}$ are the tree-level amplitudes sitting at the two sides of the cut, see Fig.1. By using (2.5) for the LIPS, and (2.2) for the loop-momentum ℓ_1^μ , one has,

$$\Delta = (1 - 2\rho) \iint dz \wedge d\bar{z} \frac{A_L^{\text{tree}}(z, \bar{z}) A_R^{\text{tree}}(z, \bar{z})}{(1 + z\bar{z})^2}, \quad (2.7)$$

¹In terms of spinor variables that are associated to massless momenta, we can define $p^\mu = (1/2)\langle p | \gamma^\mu | p \rangle$ and $q^\mu = (1/2)\langle q | \gamma^\mu | q \rangle$, hence $\varepsilon_+^\mu = (1/2)\langle q | \gamma^\mu | p \rangle$ and $\varepsilon_-^\mu = (1/2)\langle p | \gamma^\mu | q \rangle$.




Figure 1: Double-cut of one-loop amplitude in the K^2 -channel.

where the tree-amplitudes A_L^{tree} and A_R^{tree} are rational in z and \bar{z} . Since ρ is independent of z and \bar{z} , its presence in the integrand is understood. By applying a special version of the so called *Generalised Cauchy Formula* also known as the *Cauchy-Pompeiu Formula* [28], one can write the two-fold integration in z - and \bar{z} -variables appearing in Eq.(2.7) simply as a convolution of an unbounded \bar{z} -integral and a contour z -integral² [26],

$$\Delta = (1 - 2\rho) \oint dz \int d\bar{z} \frac{A_L^{\text{tree}}(z, \bar{z}) A_R^{\text{tree}}(z, \bar{z})}{(1 + z\bar{z})^2}, \quad (2.8)$$

where the product $A_L^{\text{tree}} A_R^{\text{tree}}$ is a rational function of z and \bar{z} , and the integration contour has to be chosen as enclosing all the complex z -poles. The equivalence of Eq.(2.7) and Eq.(2.8) is due to Stokes' Theorem [26]. Accordingly, the double-cut Δ in (2.7) is the flux of a 2-form, corresponding to an integral over the complex tangent bundle of the Riemann sphere³, where the curvature 2-form is defined as $\Omega = 1/(1 + |z|^2)^2 dz \wedge d\bar{z}$.

2.1 Coefficient of the 2-point function

The formula in Eq.(2.8) can be integrated straightforwardly in two steps. To begin with the integration, we find a primitive with respect to \bar{z} , say F , by keeping z as independent variable,

$$F(z, \bar{z}) = \int d\bar{z} \frac{A_L^{\text{tree}}(z, \bar{z}) A_R^{\text{tree}}(z, \bar{z})}{(1 + z\bar{z})^2}, \quad (2.9)$$

so that Δ becomes,

$$\Delta = (1 - 2\rho) \oint dz F(z, \bar{z}). \quad (2.10)$$

Since F is the primitive of a rational function, its general form can only contain two types of terms: a rational term and a logarithmic one,

$$F(z, \bar{z}) = F^{\text{rat}}(z, \bar{z}) + F^{\text{log}}(z, \bar{z}). \quad (2.11)$$

The coefficient of a 2-point function in the K^2 -channel will appear in Δ^{rat} , namely the result of the Residue Theorem in z applied only to F^{rat} . The choice of p and q specified in Eqs.(2.3) grants that there exists a pole at $z = 0$ associated to the 2-point function in the K^2 -channel, $I_2(K^2)$; while

²The roles of z and \bar{z} can be equivalently exchanged.

³In [26] it has been shown that the double-cut of the scalar 2-point function, $\Delta I_2 = \int d^4\Phi$ amounts to the integral $\iint \Omega = -2\pi i$. This result corresponds to the integration of the first Chern class, $(i/\pi) \iint \Omega = 2$.

the reduction of higher-point functions that have $I_2(K^2)$ as subdiagram can generate poles at finite z -values. The Residue Theorem has to be applied by reading all the residues in z , and substituting the corresponding complex-conjugate values where \bar{z} appears. Since it can be shown [26] that the double-cut of the 2-point scalar function in the K^2 -channel amounts to $\Delta I_2 = -2\pi i (1 - 2\rho)$, the coefficient of the 2-point function can be finally extracted from the ratio,

$$c_2 = \frac{\Delta^{\text{rat}}}{\Delta I_2} = - \left(\text{Res}_{z=0} F^{\text{rat}}(z, \bar{z}) + \text{Res}_{z \neq 0} F^{\text{rat}}(z, \bar{z}) \right), \quad (2.12)$$

which will depend on ρ .

Recently, the method just described has been successfully applied to the completion of the analytic calculation of the one-loop virtual corrections to $H + 2\text{jets}$ via gluon fusion [29, 30].⁴

3. Optical Theorem and Berry's Phase

In [27] the following observation was made. In the double-cut integral (2.7), we did not make any assumptions on the tree-level amplitudes sewn along the cut, thus providing a general framework to the integration method developed in [26]. If we now choose $A_L^{\text{tree}} = A_{m \rightarrow 2}^{*, \text{tree}}$, that is the conjugate scattering amplitude of a process $m \rightarrow 2$, and $A_R^{\text{tree}} = A_{n \rightarrow 2}^{\text{tree}}$, that is the amplitude of a process $n \rightarrow 2$, then Δ reads,

$$\Delta = \int d^4\Phi A_{m \rightarrow 2}^{*, \text{tree}} A_{n \rightarrow 2}^{\text{tree}} = -i \left[A_{n \rightarrow m}^{\text{one-loop}} - A_{m \rightarrow n}^{*, \text{one-loop}} \right] = 2 \text{Im} \left\{ A_{n \rightarrow m}^{\text{one-loop}} \right\}, \quad (3.1)$$

which is the definition of the two-particle discontinuity of the one-loop amplitude $A_{n \rightarrow m}^{\text{one-loop}}$ across the branch cut in the K^2 -channel, corresponding to the field-theoretic version of the Optical Theorem for one-loop Feynman amplitudes. On the other side, because of Stokes' Theorem in (2.7, 2.8), one has,

$$\Delta = (1 - 2\rho) \iint dz \wedge d\bar{z} \frac{A_{m \rightarrow 2}^{*, \text{tree}} A_{n \rightarrow 2}^{\text{tree}}}{(1 + z\bar{z})^2} = (1 - 2\rho) \oint dz \int d\bar{z} \frac{A_{m \rightarrow 2}^{*, \text{tree}} A_{n \rightarrow 2}^{\text{tree}}}{(1 + z\bar{z})^2}, \quad (3.2)$$

which provides a geometrical interpretation of the imaginary part of one-loop scattering amplitudes, as a flux of a complex 2-form through a surface bounded by the contour of the z -integral (the contour should enclose all the poles in z exposed in the integrand after the integration in \bar{z} [26]).

Given the equivalence of (3.1) and (3.2), a correspondence between the imaginary part of scattering amplitudes and the anholonomy of Berry's phase does emerge, since the latter is indeed defined as the flux of a 2-form in presence of curved space [1, 2]. In this context, one could establish a parallel description between the Aharonov-Böhm (AB) effect [31] and the double-cut of one-loop Feynman integrals. Accordingly, let us follow the evolution in Fig.1 from the left to the right. The two particles produced in the A_L -scattering, going around the loop and initiating the A_R -process, at the A_R -interaction point would experience a phase-shift due to the non-trivial geometry in effective momentum space induced by the on-shell conditions. As in the AB-effect, the anholonomy phase-shift is a consequence of Stokes' Theorem, and here it corresponds to the imaginary part of the one-loop Feynman amplitude.

⁴See also S. Badger, and C. Williams in these proceedings.

References

- [1] M. V. Berry, Proc. Roy. Soc. Lond. A **392** (1984) 45.
- [2] A. D. Shapere and F. Wilczek (ed.), Adv. Ser. Math. Phys. **5** (1989) 1.
- [3] L. D. Landau, Nucl. Phys. **13**, 181 (1959). R. E. Cutkosky, J. Math. Phys. **1**, 429 (1960). R. J. Eden, P. V. Landshoff, D. I. Olive, J. C. Polkinghorne, *The Analytic S Matrix*, Cambridge University Press, 1966. M. J. G. Veltman, Physica **29** (1963) 186. E. Remiddi, Helv. Phys. Acta **54** (1982) 364.
- [4] R. C. Hwa, and V. L. Teplitz, Mathematical Physics Monographs, W. A. Benjamin Inc., 1966.
- [5] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, Nucl. Phys. B **425** (1994) 217; Nucl. Phys. B **435** (1995) 59.
- [6] G. Passarino and M. J. G. Veltman, Nucl. Phys. B **160** (1979) 151.
- [7] G. Ossola, C. G. Papadopoulos and R. Pittau, Nucl. Phys. B **763**, 147 (2007); JHEP **0707**, 085 (2007).
- [8] R. K. Ellis, W. T. Giele and Z. Kunszt, JHEP **0803** (2008) 003.
- [9] W. T. Giele, Z. Kunszt and K. Melnikov, JHEP **0804** (2008) 049.
- [10] C. F. Berger *et al.*, Phys. Rev. D **78** (2008) 036003.
- [11] R. Britto, F. Cachazo and B. Feng, Nucl. Phys. B **725**, 275 (2005).
- [12] P. Mastrolia, Phys. Lett. B **644**, 272 (2007).
- [13] T. Binoth, G. Heinrich, T. Gehrmann and P. Mastrolia, Phys. Lett. B **649**, 422 (2007).
- [14] D. Forde, Phys. Rev. D **75**, 125019 (2007).
- [15] N. E. J. Bjerrum-Bohr, D. C. Dunbar and W. B. Perkins, JHEP **0804**, 038 (2008).
- [16] W. B. Kilgore, arXiv:0711.5015 [hep-ph].
- [17] S. D. Badger, JHEP **0901** (2009) 049.
- [18] R. Britto, E. Buchbinder, F. Cachazo and B. Feng, Phys. Rev. D **72**, 065012 (2005).
- [19] R. Britto, B. Feng and P. Mastrolia, Phys. Rev. D **73**, 105004 (2006).
- [20] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia, Phys. Lett. B **645**, 213 (2007); JHEP **0703**, 111 (2007).
- [21] R. Britto and B. Feng, Phys. Rev. D **75**, 105006 (2007); JHEP **0802**, 095 (2008).
- [22] R. Britto, B. Feng and P. Mastrolia, Phys. Rev. D **78** (2008) 025031.
- [23] R. Britto, B. Feng and G. Yang, JHEP **0809** (2008) 089.
- [24] E.W.N. Glover and C. Williams, JHEP **0812** (2008) 067.
- [25] R. Britto and B. Feng, Phys. Lett. B **681**, 376 (2009).
- [26] P. Mastrolia, Phys. Lett. B **678** (2009) 246.
- [27] P. Mastrolia, to appear in Lett. Math. Phys., arXiv:0906.3789.
- [28] M. J. Ablowitz, and A. S. Fokas, *Complex Variables*, Cambridge Texts in Applied Mathematics, 2003, 2nd Edition.
- [29] S. Badger, E. W. Nigel Glover, P. Mastrolia and C. Williams, to appear in JHEP, arXiv:0909.4475.
- [30] S. Badger, J. M. Campbell, R. K. Ellis and C. Williams, JHEP **0912** (2009) 035.
- [31] Y. Aharonov and D. Bohm, Phys. Rev. **115** (1959) 485.