

NNLO QCD corrections to lepton charge asymmetry in W decays at hadron colliders

G. Ferrera*

Dipartimento di Fisica e Astronomia, Università di Firenze and INFN, Sezione di Firenze

E-mail: ferrera@fi.infn.it

We discuss the effect of next-to-next-to-leading order (NNLO) QCD corrections to the charged leptons asymmetry from W^\pm boson decays in hadron collisions. The calculation includes the lepton kinematical cuts which are necessary applied in experimental analyses. We show some numerical results on the charge asymmetry in $p\bar{p}$ collisions compared with the Tevatron data and in pp collisions at LHC energy.

XVIII International Workshop on Deep-Inelastic Scattering and Related Subjects, DIS 2010

April 19-23, 2010

Firenze, Italy

*Speaker.

1. Introduction

The measure of the asymmetry in the rapidity of charged leptons from W^\pm decay is an important observable at hadron colliders. In $p\bar{p}$ collisions, the total production rates of the W^+ and W^- bosons are equal; nevertheless the W^+ tends to be produced at positive rapidity while the W^- at negative rapidity. In pp collisions, W production is forward–backward symmetric; however, the W^- production rate is smaller than the W^+ production rate and the W^- is mostly produced at central rapidities, while the W^+ is mostly produced at larger rapidities.

These W^+/W^- asymmetries are related to the different proton content of u and d quarks and, in particular, to the fact that u quarks carry, on average, a larger fraction of the proton’s momentum than d quarks. Therefore the W boson charge asymmetries provide important quantitative information on the u and d quark momentum distribution in the proton [1].

At hadron colliders, the produced W bosons are identified by their leptonic decay $W \rightarrow l\nu_l$. Since the longitudinal component of the neutrino momentum is unmeasured, what is actually measured is the rapidity of the charged lepton ¹ and the corresponding lepton charge asymmetry, which is defined as

$$A_{h_1 h_2}(y_l) = \frac{d\sigma_{h_1 h_2}(l^+)/dy_l - d\sigma_{h_1 h_2}(l^-)/dy_l}{d\sigma_{h_1 h_2}(l^+)/dy_l + d\sigma_{h_1 h_2}(l^-)/dy_l}, \quad (1.1)$$

where h_1 and h_2 label the two incoming hadrons and $d\sigma_{h_1 h_2}(l^\pm)/dy_l$ is the charged lepton (l^\pm) rapidity (y_l) distribution.

In actual experimental determinations, the measured rapidity cross sections and charge asymmetries necessary depend on the kinematical cuts and variables that are used to identify and select the observed $W \rightarrow l\nu_l$ events. Our computation of the lepton charge asymmetry is carried out by using the partonic Monte Carlo program DYNNLO [2], which encodes the NNLO radiative corrections to the Drell-Yan process at the fully-differential level ². This allows us to compute the lepton charge asymmetry by including the kinematical cuts applied in experimental analyses [4].

Our N^n LO (with $n = 0, 1, 2$) computation uses the expression of $\alpha_S(\mu_R^2)$ at the n -th order (i.e., we use the μ_R dependence at the level of $(n+1)$ loops), and it is consistently convoluted with parton densities at each corresponding order. The reference value of $\alpha_S(M_Z)$ is fixed at the actual value used in the corresponding set of parton densities. We consider finite-width effects, treating the W boson off shell, and we include the spin correlations of the W leptonic decay. The values of the electroweak parameters used in our computation can be found in Ref. [4].

Since our main purpose is to study the effect of QCD corrections to the lepton asymmetries up to NNLO, we consider only the parton distribution function (PDF) sets of Refs. [5], [6] and [7] which include NNLO parton densities with $N_f = 5$ (effectively) massless quarks. Among these PDFs sets only the global fit of Ref. [5] includes some data on the lepton charge asymmetry at the Tevatron: the ensuing parton densities are thus expected to produce better agreement with available measurements of charge asymmetries.

¹We consider the charged leptons in the massless approximation, thus the lepton rapidity y_l coincides with the pseudorapidity η_l .

²This Monte Carlo program is based on the NNLO subtraction method introduced in Ref. [3].

2. Charged lepton asymmetry at the Tevatron

In this section we present our perturbative QCD calculations up to NNLO, and their comparison with some of the published Tevatron data. Our calculation is invariant under CP transformations, therefore, in $p\bar{p}$ collisions, the charge asymmetry fulfils $A_{p\bar{p}}(-y) = -A_{p\bar{p}}(y)$.

We first consider the electron charge asymmetry in the experimental configuration of the CDF data at the Tevatron Run II [8]. The events are required to have a missing transverse energy $E_T^V > 25$ GeV and a transverse mass $50 \text{ GeV} < M_T < 100$ GeV. The charged lepton must be in the central rapidity region ($|y_l| \leq 2.45$) and must be isolated³ i.e. $E_T^{\text{iso}}/E_T < 0.1$, where E_T^{iso} is the hadronic (partonic) transverse energy in a cone of radius $R = 0.4$ (in pseudorapidity–azimuth space) along the lepton momentum and E_T is the electron transverse energy. Following the experimental analysis we consider two kinematical region of the electron transverse energy: $25 \text{ GeV} < E_T < 35$ GeV (low- E_T) and $35 \text{ GeV} < E_T < 45$ GeV (high- E_T).

In Fig. 1(a) we report the CDF data and present the results of our LO, NLO and NNLO results using the corresponding parton densities from the MSTW2008 Collaboration [5], which include the CDF data [8] in their fit. At small values of η_e , the radiative corrections lead to little effects. In the low- E_T bin, as η_e increases, both the NLO and NNLO effects slightly increase the value of the asymmetry. In the high- E_T bin, as η_e increases, the NLO and NNLO effects are negative and the asymmetry slightly decreases. The NNLO effects are not yet quantitatively relevant in comparison with the size of the experimental uncertainties. The data are well fitted by the MSTW2008 NNLO partons, especially in the low- E_T region, this is confirmed by the values of χ^2 from the comparison of the data with the QCD calculations reported in Fig. 1(a).

We then consider the D0 data for the electron charge asymmetry [9] at the Tevatron Run II. The D0 measurement extends to high rapidities, $|\eta_e| \leq 3.2$, the missing transverse energy is required to be $E_T^V > 25$ GeV and the transverse mass to be $M_T > 50$ GeV. As for lepton isolation the D0 procedure is similar to the CDF one, with the requirement $E_T^{\text{iso}}/E_T < 0.15$. We examine two regions of electron transverse energy: $E_T > 25$ GeV (wide E_T region) and $E_T > 35$ GeV (high- E_T region). Let us note that the D0 electron asymmetry data [9] are not included in the MSTW PDF fit [5]: their inclusion in the global analysis does not permit to obtain a good quality fit, with significant tension between the D0 electron asymmetry data and other data.

In Fig. 1(b) we display the D0 electron data and present the corresponding QCD results at LO, NLO and NNLO using the corresponding MSTW2008 PDFs. In the wide E_T region at high values of the electron rapidity, the effect of the NLO and NNLO corrections is positive and increases the deviation of the QCD results from the D0 data. We see that the agreement between the QCD calculations and the data is poor. This is quantitatively confirmed by the values of χ^2 reported in Fig. 1(b). We note that the value of χ^2 increases in going from LO to NLO and to NNLO. In the high- E_T region (as in the case of the high- E_T bin in Fig. 1(a)) the effect of the NLO and NNLO corrections is negative as the lepton rapidity increases toward high values. This effect reduces the difference between the QCD calculations and the D0 data, although a substantial disagreement between them still persists at the NNLO. The value of χ^2 decreases in going from LO to NLO and to NNLO.

³The charged lepton isolation has a small impact on our results, since it is not effective at the leading order and the radiative corrections are not large.




Figure 1: Electron charge asymmetry up to NNLO compared to a) CDF data in the lower E_T region (upper panel) and higher E_T region (lower panel); b) D0 data in the wide E_T region (upper panel) and (high- E_T) region.

The NNLO results of Fig. 1(b) are reported in Fig. 2(a) by including the PDF errors (68% C.L.) from the MSTW2008 [5], ABKM09 [6] and JR09VF [7] set of PDFs. The ABKM09 PDF errors are slightly smaller than the errors of the MSTW2008 set and the ABKM09 result tends to overshoot the MSTW2008 result. The JR09VF PDF errors are larger than the PDF errors of the MSTW2008 set and the JR09VF result tends to undershoot the MSTW2008 result in most of the rapidity bins. The PDF errors are comparable to (or, larger than) the D0 experimental errors; the inclusion of the PDF errors thus reduces the differences between the NNLO results and the D0 electron data, although a substantial disagreement can be noticed from the value of χ^2 for different sets of PDFs reported in Fig. 2(a).

3. Charged lepton asymmetry at the LHC

We now consider pp collisions and we present results of our QCD calculations at the centre-of-mass energy $\sqrt{s} = 7$ TeV. The lepton charge asymmetry at the LHC is sensitive to parton densities with momentum fractions that are smaller than those probed at the Tevatron. To the purpose of presenting some quantitative, though illustrative, results on the lepton asymmetry at the LHC, we refer to the framework considered in a recent 'Physics Analysis' [10] of the CMS Collaboration. The study of Ref. [10], based on data sets from Monte Carlo simulations, regards the muon rapidity cross sections and charge asymmetry that can be measured with the CMS detector at the LHC. The lepton selection cuts applied in Ref. [10] require the transverse mass and the missing transverse energy to be $M_T > 50$ GeV and $E_T^V > 20$ GeV, respectively. The muons are isolated: the hadronic (partonic) transverse energy E_T^{iso} in a cone of radius $R = 0.3$ is required to fulfil the constraints




Figure 2: a) Electron charge asymmetry at NNLO with PDF uncertainties compared to D0 data in the lower E_T region (upper panel) and higher E_T region (lower panel); b) Muon charge asymmetry up to NNLO at the LHC.

$$E_T^{\text{iso}}/E_T < z/(1-z) \text{ with } z = 0.05, \text{ and } E_T + E_T^{\text{iso}} > E_T^{\text{max}} = 25 \text{ GeV.}$$

The results for the lepton charge asymmetry at $\sqrt{s} = 7$ TeV are presented in Fig. 2(b). The three histograms show the LO, NLO and NNLO results of our calculation with the MSTW2008 parton densities. Considering the effect of the QCD radiative corrections, we see that, the NLO corrections tend to decrease the asymmetry, and the NNLO corrections do not significantly change the NLO result thus indicating a very good quantitative convergence of the truncated perturbative expansion.

References

- [1] E. L. Berger, F. Halzen, C. S. Kim and S. Willenbrock, Phys. Rev. D **40** (1989) 83 [Erratum-ibid. D **40** (1989) 3789]; A. D. Martin, R. G. Roberts and W. J. Stirling, Mod. Phys. Lett. A **4** (1989) 1135.
- [2] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, Phys. Rev. Lett. **103** (2009) 082001.
- [3] S. Catani and M. Grazzini, Phys. Rev. Lett. **98** (2007) 222002.
- [4] S. Catani, G. Ferrera and M. Grazzini, JHEP **1005** (2010) 006 [arXiv:1002.3115 [hep-ph]].
- [5] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, Eur. Phys. J. C **63** (2009) 189.
- [6] S. Alekhin, J. Blumlein, S. Klein and S. Moch, report DESY 09-102 (arXiv:0908.2766 [hep-ph]).
- [7] P. Jimenez-Delgado and E. Reya, report DOT-0913 (arXiv:0909.1711 [hep-ph]).
- [8] D. E. Acosta *et al.* [CDF Collaboration], Phys. Rev. D **71** (2005) 051104.
- [9] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **101** (2008) 211801.
- [10] CMS Collaboration, report CMS PAS Note EWK-09-003 (2009).