

## Hard X-ray Variability of the Brightest Swift/BAT AGN

---

**Claudio Ricci\***

*ISDC Data Centre for Astrophysics  
Geneva Observatory, University of Geneva  
E-mail: claudio.ricci@unige.ch*

**S. Paltani**

*ISDC Data Centre for Astrophysics  
Geneva Observatory, University of Geneva*

**S. Soldi**

*Laboratoire AIM - CNRS - CEA/DSM - Université Paris Diderot (UMR 7158), CEA Saclay,  
DSM/IRFU/SAP, F91191 Gif-sur-Yvette, France*

**T. J.-L. Courvoisier**

*ISDC Data Centre for Astrophysics  
Geneva Observatory, University of Geneva*

Variability is one of the hallmarks of Active Galactic Nuclei. The Burst Alert Telescope onboard of *Swift*, with its homogeneous coverage of the sky is a formidable tool to study variability at hard X-rays. We present here the analysis of the 1-month binned *Swift*/BAT lightcurves of the 20 brightest Active Galactic Nuclei in the hard X-ray sky. The sample consists of 2 blazars, 3 radio galaxies, 6 Seyfert 1/1.5s, 8 Seyfert 2s and 1 Narrow Line Seyfert 1. We found that all the objects show variability, and most of them have a value of the fractional root mean squared variability amplitude of  $F_{\text{var}} \sim 0.2 - 0.3$ . We did not find any significant correlation of  $F_{\text{var}}$  with the column density or the luminosity in our small sample.

*Fast X-ray timing and spectroscopy at extreme count rates  
February 7-11, 2011  
Champéry, Switzerland*

---

\*Speaker.

## 1. Introduction

Active Galactic Nuclei (AGN) are amongst the most luminous X-ray sources in the sky. AGN are thought to be powered by accretion onto supermassive black holes [1], with their X-ray emission probably originating in a hot corona sandwiching the accretion disk [2] in radio-quiet objects, and in the jet in radio-loud AGN (e.g., [3]). Variability is one of the key features of AGN, and it was found to be significant in the X-ray band already in early observations of nearby Seyfert galaxies (Sy) performed by *Ariel V* [4]. The X-ray variability of AGN is aperiodic, and their power spectral density distribution (PSD) can be normally described with a broken power law, with indices ranging between  $-1$  and  $-2$  [5].

The Burst Alert Telescope (BAT) onboard of *Swift* [6] scans continuously the whole sky in the 14–195 keV energy range, and is thus an extremely well suited instrument for studying AGN variability at hard X-rays. Here we report a study of the hard X-ray variability of a small sample of AGN. The sample consists of the 20 brightest AGN detected by *Swift*/BAT, of these 2 are blazars, 2 narrow-line radio galaxies (NLRG), 1 broad-line radio galaxy (BLRG), 6 Seyfert 1/1.5s, 8 Seyfert 2s and 1 Narrow Line Seyfert 1 (NLS1). The 1-month binned light curves have been taken from the NASA *Swift*/BAT 58 months catalog<sup>1</sup> [7].

## 2. Variability estimators

A way to estimate the variability is through the fractional root mean squared (rms) variability amplitude  $F_{\text{var}}$  ([8]), defined as

$$F_{\text{var}} = \sqrt{\frac{S^2 - \overline{\sigma_{\text{err}}^2}}{\bar{x}^2}}. \quad (2.1)$$

Where the sample variance  $S^2$  is given by

$$S^2 = \frac{1}{N-1} \sum_{i=1}^N (x_i - \bar{x})^2, \quad (2.2)$$

while the mean square error  $\overline{\sigma_{\text{err}}^2}$  by

$$\overline{\sigma_{\text{err}}^2} = \frac{1}{N} \sum_{i=1}^N \sigma_{\text{err},i}^2. \quad (2.3)$$

The error of  $F_{\text{var}}$  is given by

$$\text{err}(F_{\text{var}}) = \sqrt{\left( \sqrt{\frac{1}{2N}} \cdot \frac{\overline{\sigma_{\text{err}}^2}}{\bar{x}^2 F_{\text{var}}} \right)^2 + \left( \sqrt{\frac{\overline{\sigma_{\text{err}}^2}}{N}} \cdot \frac{1}{\bar{x}} \right)^2}. \quad (2.4)$$

In the following we will use  $F_{\text{var}}$  to characterize the variability of the objects in our sample.

<sup>1</sup><http://swift.gsfc.nasa.gov/docs/swift/results/bs58mon/>


**Table 1:** Properties of the sources of our sample: (1) detection significances, (2) luminosities in the 14–195 keV energy range, (3) fractional rms variability amplitudes on a timescale of 30 days, (4) hydrogen column densities.

Source	(1) Det. Significance [ $\sigma$ ]	(2) $\log L_{14-195\text{keV}}$ [ $\text{ergs}^{-1}$ ]	(3) $F_{\text{var}} (30\text{-days})$	(4) $N_{\text{H}}$ [ $\text{cm}^{-2}$ ]	Type
Cen A	428.7	44.01	$0.399 \pm 0.002$	$12^a$	NLRG
NGC 4151	275.0	44.11	$0.280 \pm 0.004$	$6.9^b$	Sy 1.5
3C 273	156.8	47.47	$0.31 \pm 0.01$	$0.5^b$	Blazar
NGC 4388	110.7	44.64	$0.31 \pm 0.01$	$27^b$	Sy 2
Mrk 421	109.5	45.46	$0.96 \pm 0.01$	$0.1^b$	Blazar
Circinus Galaxy	101.7	43.09	$0.13 \pm 0.01$	$360^b$	Sy 2
IC 4329A	101.1	45.22	$0.19 \pm 0.02$	$0.4^b$	Sy 1
NGC 2110	98.1	44.60	$0.32 \pm 0.01$	$4.3^c$	Sy2
NGC 5506	95.0	44.31	$0.27 \pm 0.02$	$3.4^c$	NLS1
MCG–05–23–016	90.4	44.50	$0.21 \pm 0.01$	$1.6^c$	Sy2
IGR J21247+5058	83.3	45.25	$0.31 \pm 0.01$	$0.6^b$	BLRG
NGC 4945	76.1	43.37	$0.35 \pm 0.02$	$400^b$	Sy2
Mrk 348	70.4	44.91	$0.28 \pm 0.02$	$30^c$	Sy2
NGC 3783	68.7	44.60	$0.26 \pm 0.02$	$0.1^c$	Sy1.5
NGC 4507	64.6	44.77	$0.29 \pm 0.02$	$29^c$	Sy 2
NGC 3516	62.3	44.33	$0.30 \pm 0.02$	$4^c$	Sy 1.5
NGC 7172	60.1	44.46	$0.35 \pm 0.04^*$	$9^c$	Sy 2
NGC 3227	56.2	43.57	$0.16 \pm 0.21$	$6.8^c$	Sy 1.5
Cyg A	54.0	46.01	$0.34 \pm 0.02$	$11^b$	NLRG
MCG +08–11–011	49.0	45.09	$0.35 \pm 0.03$	$0.2^c$	Sy 1.5
Crab	7496	–	$0.0215 \pm 0.0004$	–	–

**Notes.** <sup>a</sup> [9], <sup>b</sup> [10] and references therein, <sup>c</sup> [11] and ref. therein. \* "Flare" of January 2008 removed.

### 3. Results

In Table 1 are listed the values of the fractional rms variability amplitude of the objects of our sample, and as a check, the value obtained from the light curve of the Crab ( $F_{\text{var}} \sim 0.02$ ). The value of the Crab can be associated to the systematic error of the *Swift*/BAT data. In Fig. 1, we show the light curves of 6 out of the 20 sources of our sample. All the sources of our sample show hard X-ray variability on the time-scale of one month. As it can also be seen from Fig. 2, the value of the fractional rms variability amplitude is  $F_{\text{var}} \sim 0.2 - 0.3$  for most of the objects of the sample, with the average value being  $\overline{F_{\text{var}}} = 0.32$ . The blazar Mrk 421 shows a much stronger variability ( $F_{\text{var}} \sim 0.96$ ) than the average value of the sample. Amongst the radio-quiet NGC 7172 is the most variable, with a value of  $F_{\text{var}} \sim 0.48$ . This is due to what would appear to be a flare, registered in January 2008. Excluding this outlier point NGC 7172 shows a variability consistent with the




**Figure 1:** *Swift*/BAT 30-days binned light curves in the 14–195 keV band. The dotted horizontal lines represent the average value for each object.


average of our sample ( $F_{\text{var}} = 0.35 \pm 0.04$ ). At the other end of the distribution, the Compton-thick Seyfert 2 Circinus Galaxy and the Seyfert 1 IC 4329A show the smallest amounts of variability ( $F_{\text{var}} \sim 0.13$  and  $F_{\text{var}} \sim 0.19$ , respectively). The low value of  $F_{\text{var}}$  of Circinus Galaxy is very likely related to the reflection-dominated nature of its hard X-ray spectrum.

#### 4. Variability vs Luminosity and Column density

An inverse correlation between the variability amplitude in the X-rays and the X-ray luminosity of AGN was found by [12] using *EXOSAT* data. More recently, [13] studied the hard X-ray variability of the 44 brightest AGN detected by BAT after 9 months of operations, and found that possibly this anti-correlation is extended to the hard X-ray band (see also [14]). They also found a possible correlation between the hydrogen column density  $N_{\text{H}}$  and the variability amplitude. We investigated the existence of these two correlations in our sample (see Figs. 3 and 4). The variability amplitudes in our sample are confined in a small range of values, and no correlation with other



**Figure 2:** Distribution of the fractional rms variability amplitude for the 20 sources of our sample.




**Figure 3:**  $F_{\text{var}}$  versus luminosity for the sources of our sample.

parameters is evident. A Spearman rank test gives a correlation coefficient between  $F_{\text{var}}$  and the luminosity of  $r_s = 0.27$ , while it is  $r_s = 0$  between  $F_{\text{var}}$  and  $N_{\text{H}}$ . These values correspond to a probability of correlation of 78% in the first case, and of 0% in the second case. Similar results are obtained also considering Mrk 421 as an outlier. No significant correlation is found also dividing the sample in three categories (Sy 1/1.5, Sy2, radio-loud). The lack of correlations might be due to the limited sample we used, and further studies are needed to better probe it.

## 5. Conclusions

Studying the 1-month binned *Swift*/BAT light-curves of the 20 brightest objects after 58 months



**Figure 4:**  $F_{\text{var}}$  versus hydrogen column density for the sources of our sample.

of observations, we found that all the objects in our sample show variability, ranging between  $F_{\text{var}} = 0.13$  and  $F_{\text{var}} = 0.96$ , for Circinus galaxy and Mrk 421, respectively. The average value of the variability amplitude is  $F_{\text{var}} \sim 0.3$ . We did not find any significant correlation of the variability amplitude with the luminosity and the hydrogen column density.

## References

- [1] Rees, M. J. 1984, ARA&A 22, 471
- [2] Haardt, F., & Maraschi, L. 1991, ApJ 380, L51
- [3] Boettcher, M., arXiv:1006.5048
- [4] Marshall, N., Warwick, R. S., Pounds, K. A. 1981, MNRAS, 194, 987
- [5] McHardy, I. M., & Czerny, B. 1987, Nature, 325, 696
- [6] Barthelmy, S. D., Barbier, L. M., Cummings, J. R., et al. 2005, SSRv, 120, 143
- [7] Baumgartner et al, 2011 ApJS, submitted
- [8] Edelson, R. A.; Krolik, J. H.; Pike, G. F. 1990, ApJ 359, 86
- [9] Beckmann, V.; Jean, P.; Lubinski, P., et al. 2011, A&A in press, arXiv:1104.4253v2
- [10] Beckmann, V.; Soldi, S.; Ricci, C., et al. 2009, A&A 505, 417
- [11] Ricci, C.; Walter, R.; Courvoisier, T. J.-L.; et al. 2011, A&A in press, arXiv:1104.3676
- [12] Barr, P., & Mushotzky, R. F. 1986, Nature, 320, 421
- [13] Beckmann, V.; Barthelmy, S. D.; Courvoisier, T. J.-L.; et al. 2007, A&A 475, 827
- [14] Soldi, S.; Ponti, G.; Beckmann, V.; et al. 2010, arXiv1001.4348S