

CMS ridge effect at the LHC as a manifestation of Bremsstrahlung of gluons off quarks accelerated in a strong color field

Boris Arbuzov*

SINP MSU

E-mail: arbuzov@theory.sinp.msu.ru

The recently reported effect of long-range near-side angular correlations at LHC occurs for large multiplicities of particles with $1 \text{ GeV} < p_T < 3 \text{ GeV}$. To understand the effect several possibilities have been discussed. In the talk we propose [1] a simple qualitative mechanism which corresponds to gluon bremsstrahlung of quarks moving with acceleration defined by the string tension. The smallness of azimuth angle difference $\Delta\phi$ along with large $\Delta\eta$ at large multiplicities in this interval of p_T are natural in the mechanism. The mechanism predicts also bremsstrahlung photons with mean values of $p_T \approx 2.9 \times 2.5/m_u (\text{MeV})$ and $0.72 \text{ GeV} \times 5/m_d (\text{MeV})$.

*The XXth International Workshop High Energy Physics and Quantum Field Theory
September 24 - October 1, 2011
Sochi, Russia*

*Speaker.

In paper [2] the effect in proton-proton collisions at LHC is reported for existence of a ridge in the plot of data for two-particle correlations versus pseudo-rapidity difference $\Delta\eta$ and azimuth angle $\Delta\phi$ plane. This ridge means essential excess of events with $\Delta\phi$ close to zero and large $\Delta\eta$. It is important to emphasize, that the effect is observed under condition that the accompanying charged particles have multiplicity > 100 and are each situated in restricted region of transverse momentum $1\text{ GeV} < p_T < 3\text{ GeV}$.

The result already causes discussion devoted to possible interpretation of the data [3] – [15]. In note [3] possibilities of interpretation of the effect in terms of quark-anti-quark strings are discussed. The first point is that a string is formed being stretched close to the direction of pp collision and thus might decay with $\Delta\phi \approx 0$ and $\approx \pi$. However it is emphasized in [3], that there is no reason, why the effect is observed only for very high multiplicity $N > 100$, while multiple production of hadrons via string decomposition leads to randomization and thus simulations show no ridge.

In the present talk [1] we would try to explain why even for high multiplicity the correlation persists when we take into account radiation of gluons in the process of QCD quark-anti-quark string formation in proton-proton collision at very high energies. As a matter of fact we mean that in the course of collision a quark is moving with acceleration in the strong colour field which in particular may be described in terms of formation of a string between either quark from the first proton and anti-quark from the second one or vice versa. However we do not use some particular string model (see e.g. [16]). We qualitatively assume a formation of some extended object of cigar-like shape inside which quark and anti-quark interact with each other due to strong colour field which accelerate both of them. We shall use simple qualitative considerations based on the classical theory of radiation which is described e.g. in book [17] and then we perform quasi-classical estimates.

Each quark (anti-quark) moves considerable time in a very strong colour field which we describe in terms of the string tension A . The direction of the acceleration is evidently close to the direction of momenta of the colliding protons. In this case an acceleration has almost the same direction as the velocity and thus we may use the well-known classical expression for dipole electromagnetic radiation of electric charge e moving with acceleration being parallel to velocity of the motion [17]

$$\frac{dE}{dt} = \frac{2\alpha w^2}{3}; \tag{1}$$

where w is an acceleration and α is the fine structure constant. We take initial expression (1) in an accompanying reference frame. For strongly interacting quarks we change (1) for the following relation

$$\frac{dE}{dt} = \frac{\alpha_s}{9} \left(\frac{A^2}{m} \right)^2; \tag{2}$$

where acceleration $w = A^2/m$ with A and m being the string tension and a light quark mass. We also change $\alpha \rightarrow \alpha_s$ with the evident colour factor. In view to make estimates we take the following values for fundamental quantities entering (2)

$$m_u = 2.5\text{ MeV}; m_d = 5\text{ MeV}; A = 420\text{ MeV}; \tag{3}$$

where light quark masses are chosen to be in the middle of interval of their possible values: $1.7\text{ MeV} < m_u < 3.3\text{ MeV}$; $4.1\text{ MeV} < m_d < 5.8\text{ MeV}$ [18].

So quarks are moving with acceleration and thus radiate gluons. Let us obtain simple quasi-classical estimate of the mean energy of radiated gluons in an accompanying reference frame. In view of this we rewrite expression (2) in the following form

$$\frac{\Delta E}{\Delta t} = \frac{\alpha_s}{9} \left(\frac{A^2}{m} \right)^2; \quad (4)$$

Now to obtain the quasi-classical estimates we use the well-known uncertainty relation

$$\Delta E \Delta t = 1. \quad (5)$$

Finally we have for the mean energy of a gluon

$$\Delta E = \sqrt{\frac{\alpha_s}{9} \frac{A^2}{m}}. \quad (6)$$

Then we use the standard one loop expression for α_s at scale ΔE

$$\alpha_s(\Delta E) = \frac{12\pi}{(33 - 2N_f) \ln \left(\frac{\Delta E^2}{\Lambda_{QCD}^2} \right)}; \quad (7)$$

We have for one loop expression (7) with $N_f = 4$, $\Lambda_{QCD} \approx 190 \text{ MeV}$ ¹. Then with this result the solution of relations (6, 7) under conditions (3) gives us the following estimates for radiation off quarks u and d

$$\Delta E_u \approx 11.2 \text{ GeV}; \quad \Delta E_d \approx 5.6 \text{ GeV}. \quad (8)$$

The result (8) gives an estimate for mean energies of the bremsstrahlung gluons in an accompanying reference frame.

First of all let us consider an explanation of large differences in pseudo-rapidity $\Delta\eta$ along with small differences in azimuth angle $\Delta\phi$. Here we are to take into account both quarks constituting the extended object (a cigar). Namely let "the cigar" be produced with some overall momentum k while its position remains being (almost) parallel to the line of pp collision. Such situation is presented in 1.




Figure 1: The string moving with momentum k from the point of collision of two protons, ψ_1 , ψ_2 are angles and q_1 , q_2 are momenta of the quarks.

¹We normalize α_s at the point of τ -lepton mass due to better precision of data here.

Then velocities of quarks are not parallel to the direction of acceleration, but constitute some angles ψ_1, ψ_2 with this direction in laboratory reference frame. When a velocity and an acceleration are not parallel $\mathbf{v} \cdot \mathbf{w} = v w \cos \psi$ and there are two accelerated quarks we have the following angular distribution [17]

$$\begin{aligned} \frac{dE}{dt'} &= \frac{\alpha_s}{24\pi} \left(\frac{A^2}{m} \right)^2 \times \\ &\left(\Phi(\psi_1, \theta, \phi, v_1) + \Phi(\psi_2, \theta, \phi, v_2) \right) d\Omega; \end{aligned} \quad (9)$$

$$\begin{aligned} \Phi(\psi, \theta, \phi, v) &= \frac{X + v^2 Y}{Z^5} \\ X &= \sin^2 \theta - 2v \sin \psi \sin \theta \cos \phi \\ Y &= \cos^2 \theta \sin^2 \psi + \sin^2 \theta \sin^2 \psi \cos^2 \phi \\ Z &= 1 - v(\cos \psi \cos \theta + \sin \psi \sin \theta \cos \phi); \end{aligned}$$

where t' is a time with account of a retardation [17], ψ_1, ψ_2 are respectively angles for the first and the second quark. Small $\Delta\phi$ along with the wide spread in pseudo-rapidity in the effect [2] is connected with the same sign of $\sin \psi$, because quarks are directed to one side from the line of collision (see 1). Then the distribution in polar angle θ has two maxima divided by some significant interval $\Delta\theta$, while distribution in azimuth angle ϕ is again close to zero. After integration of (9) by ϕ and θ correspondingly we have these distributions. The situation is illustrated in 2 and 3, in which we present normalized distribution in rapidity η and normalized angular distribution in ϕ in ϕ for $\psi_1 = 0.1, \psi_2 = \pi - 0.1, v_1 = v_2 = 0.999$. From 2, 3 we see that $\Delta\eta$ may be quite significant while $\Delta\phi$ is small. One should note that the peaks in 2 and 3 become narrower with increasing of speed and with increasing of ψ . Emphasize that the effect of a peak around $\phi = 0$ is connected with transverse movement of "the cigar" as it is illustrated in 1. The more is transverse momentum k , i.e. angles ψ_i , the narrower becomes the distribution in ϕ . Provided transverse momentum k being very small the distribution in ϕ becomes practically isotropic.

$$\begin{aligned} \frac{dE(\eta)}{dt'} &= \frac{\alpha_s}{24\pi} \left(\frac{A^2}{m} \right)^2 \Phi(\eta) \frac{d\eta}{\cosh^2 \eta}; \\ \Phi(\eta) &= \int_{-\pi}^{\pi} \Phi_{12}(\psi_i, v_i, \theta, \phi) \cos \theta = f(\eta) d\phi; \end{aligned} \quad (10)$$

$$f(\eta) = \frac{\sinh \eta}{\cosh \eta};$$

$$\begin{aligned} \frac{dE(\phi)}{dt'} &= \frac{\alpha_s}{24\pi} \left(\frac{A^2}{m} \right)^2 \Phi(\phi) d\phi; \\ \Phi(\phi) &= \int_0^{\pi} \Phi_{12}(\psi_i, v_i, \theta, \phi) \sin \theta d\theta; \end{aligned} \quad (11)$$

$$\begin{aligned} \Phi_{12}(\psi_i, v_i, \theta, \phi) &= \Phi(\psi_1, \theta, \phi, v_1) + \\ &\Phi(\psi_2, \theta, \phi, v_2). \end{aligned}$$




Figure 2: Behavior of $\Phi_N(\eta)$, $v = 0.999$, $\psi_1 = 0.1$, $\psi_2 = \pi - 0.1$, $-5 < \eta < 5$.




Figure 3: Behavior of $\Phi_N(\phi)$, $v = 0.999$, $\psi_1 = 0.1$, $\psi_2 = \pi - 0.1$, $-\pi < \phi < \pi$.

Now let us consider properties of gluon radiation of a single quark. For this purpose we approximately assume the same direction of the velocity and of the acceleration in the actual reference frame of LHC. Angular distribution is described by the following expression [17]

$$\begin{aligned} \frac{dE}{dt'} &= \frac{\alpha_s}{24\pi} \left(\frac{A^2}{m}\right)^2 \frac{\sin^2 \theta'}{(1 - v \cos \theta')^5} d\Omega = \\ &= \frac{\alpha_s}{24\pi} \left(\frac{A^2}{m}\right)^2 \Phi_0(\theta') d\Omega; \\ \Phi_0(\theta') &= \Phi(0, \theta', \phi, v); \end{aligned} \tag{12}$$

where v is a velocity of a quark, θ' is a polar angle and $d\Omega = \sin \theta' d\theta' d\phi$. Using angular distribution of the radiation (12) we estimate the mean p_T of the radiated gluon

$$\begin{aligned}
\langle p_T^g \rangle &= \frac{\Delta E}{\sqrt{1-v^2}} \frac{I_1}{I_2}; \quad I_2 = \int \Phi_0(\theta') d\Omega; \\
I_1 &= \int \Phi_0(\theta') A(v, \theta') \sin \theta' d\Omega; \\
A(v, \theta') &= 1 + \frac{\cos \theta' (1-v^2) - v \sin^2 \theta'}{1-v^2 \cos^2 \theta'};
\end{aligned} \tag{13}$$

where $\Phi_0(\theta')$ is defined in (12). Calculating integrals in (13) with the aid of the following relation valid for $v \rightarrow 1$ and $\rho > \frac{\mu}{2}$

$$\int_0^\pi \frac{\sin^{\mu-1} \theta d\theta}{(1-v \cos \theta)^\rho} = \frac{2^{\mu-\rho} \Gamma\left(\frac{\mu}{2}\right) \Gamma\left(\frac{1}{2}\right) \Gamma(2\rho - \mu)}{(1-v^2)^{\rho-\mu/2} \Gamma(v) \Gamma\left(\frac{1-\mu}{2} + \rho\right)};$$

we obtain for quark u and d respectively with $v \rightarrow 1$

$$\begin{aligned}
\langle p_T^g(u) \rangle &= \frac{9\pi \Delta E_u}{32} \approx 9.9 \text{ GeV}; \\
\langle p_T^g(d) \rangle &= \frac{9\pi \Delta E_d}{32} \approx 4.95 \text{ GeV}.
\end{aligned} \tag{14}$$

We have to bear in mind also that in the process of hadronization a gluon give few ordinary hadrons. We assume that a radiating quark moves together with the proton to which it belongs. Then we estimate the multiplicity for gluon energy in an accompanying reference frame (8) by the following expression valid in the region of few GeV for charged multiplicity [19]

$$\begin{aligned}
\langle N_{ch} \rangle &= a + b \ln \sqrt{s}; \\
a &= -0.43 \pm 0.09; \quad b = 2.75 \pm 0.06.
\end{aligned} \tag{15}$$

Neutral particles have to be also taken into account. In view to estimate the total multiplicity we multiply expression (15) by $\frac{3}{2}$. Then we estimate $\sqrt{s} = \sqrt{2\Delta E M_p + M_p^2}$ and corresponding mean multiplicity

$$\begin{aligned}
u: \sqrt{s} &= 4.15 \text{ GeV}; \quad \langle N \rangle = 5.2; \\
d: \sqrt{s} &= 3.37 \text{ GeV}; \quad \langle N \rangle = 4.3.
\end{aligned} \tag{16}$$

We take values (16) with spread ± 2 and thus obtain estimate for transverse momenta of hadrons $p_T = p_T^g/N$

$$\begin{aligned}
u: 1.3 \text{ GeV} &< p_T < 3.0 \text{ GeV}; \\
d: 0.8 \text{ GeV} &< p_T < 2.0 \text{ GeV}.
\end{aligned} \tag{17}$$

Estimates (17) just correspond to the interval of the ridge effect [2]. Of course, by changing light quark masses in their allowable regions we can move boundaries in (17). However order of magnitude of the effect remains the same.

Next point of our interpretation is that gluons are flying in the narrow cone in the directions of a quark and the average angular spread for the multiple gluon radiation is estimated to be

$$\langle \Delta\bar{\theta} \rangle \simeq \frac{\langle p_T^g \rangle \sqrt{N_g}}{\langle E_g \rangle N_g}; \quad (18)$$

where N_g is the multiplicity of bremsstrahlung gluons in the event. Obtaining (18) we take into account that average transverse momentum squared for N_g produced gluons

$$\langle p_T^2(N_g) \rangle = \langle p_T^g \rangle^2 N_g$$

due to statistical nature of the multiple radiation. We take for transverse momentum of a gluon p_T^g estimates (14) and $\langle E_g \rangle$ is a mean energy of a gluon. Dispersion of polar angle $\Delta\bar{\theta}$ leads to dispersion of angles ψ_i in distributions (10, 11). In this connection a significant increase of $\Delta\bar{\theta}$ leads to widening of distribution in ϕ (11). From (18) we see, that for small multiplicity of gluons $\Delta\bar{\theta}$ increases and this explains why the effect disappears in this case. For estimation of the real experimental situation [2] we replace the denominator in (18) by the energy of partons' collision

$$\langle \Delta\bar{\theta} \rangle \simeq \frac{2 \langle p_T^g \rangle \sqrt{N_g}}{\sqrt{x_1 x_2 s}}; \quad (19)$$

where x_1, x_2 are values of x for quark in the first proton and the anti-quark in the second one. Number of radiated gluons N_g depends on angle ψ and velocity v . Using again formulas from [17] we have the following estimate

$$N_g = \frac{\sqrt{x_1 x_2 s}}{2\Delta E \sqrt{1 + \frac{\sin^2 \psi}{(1-v^2)}}}. \quad (20)$$

For example with $\psi = 0.1$ and $v = 0.999$, average $\Delta E = (\Delta E_u + \Delta E_d)/2 = 8.4 \text{ GeV}$, $\sqrt{s} = 7 \text{ TeV}$ [2] and with average of the product $\langle x_1 x_2 \rangle \approx 0.01$ (see, e.g. [20] and references therein) we have $N_g \simeq 17$. Now in our interpretation one bremsstrahlung gluon gives average number of charged hadrons $N_{ch} \approx 3.2$. Bearing in mind, that our quasi-classical estimate corresponds to non-coherent production of gluons, with $N_g \approx 17$ we estimate total number of charged particles produced by a quark $N_{ch}^q = 54$ that gives just multiplicity ≥ 100 for two radiating quarks. Provided one takes into account bremsstrahlung of gluons off all valence quarks moving in the strong colour field the multiplicity of charged particles can only increase, that corresponds to results [2] for lower limit of multiplicity $N_{ch}^q \geq 100$. So our mechanism does not contradict to real experimental situation [2].

However our qualitative considerations do not take into account possible correlations between gluons, that may lead to decreasing of the estimate under discussion.

Now with $N_g = 17$, $\sqrt{s} = 7 \text{ TeV}$, average $\langle p_T^g \rangle = 7.4 \text{ GeV}$ and $\langle x_1 x_2 \rangle = 0.01$ we have from (19)

$$\langle \Delta\bar{\theta} \rangle \approx 0.09; \quad (21)$$

This angular spread (21) actually gives widening of distributions (10, 11) in η and ϕ . The resulting $\Delta\phi$ is to be obtained by simultaneous account of (21) and of $\Phi(\phi)$ width (11). Let us also draw attention to widening of the ridge with \sqrt{s} decreasing. E.g. for $\sqrt{s} = 0.9 \text{ TeV}$ in accordance with (19) $\langle \Delta\bar{\theta} \rangle = 0.8$, that means vanishing of the effect because distribution (11) becomes very broad.

At this point we would emphasize, that with energy increasing on the one side angle distributions (10, 11) become narrower and thus the effect becomes more prominent and on the other side number of radiated gluons in the strong field increases that leads to increasing of the multiplicity. These properties just correspond qualitatively to the essence of effect [2].

Thus, one can conclude the simple mechanism of gluon bremsstrahlung off quarks moving in a strong colour field describes qualitatively the CMS ridge effect. Of course, a real situation could be much more involved. In particular, other colour configurations, as was pointed out in various studies (see, for example, [13]), may play a significant role. Our consideration based on simple quasi-classical estimations shows that constituting string configurations may lead to basic features of the ridge effect, namely, correlations in particular kinematic region at very high multiplicities. Obviously, in order to show more accurate properties of proposed mechanism one should elaborate in more detail corresponding model and develop corresponding event generator to perform more realistic simulations.

Let us note, that the accelerated quarks radiate photons as well. The same quasi-classical estimation gives for radiated photons two values of mean p_T for two values of (anti-)quark charge (shown in brackets)

$$\begin{aligned} \left(\frac{2e}{3}\right) : p_T &\approx 2.9 \text{ GeV} \times \frac{2.5}{m_u(\text{MeV})}; \\ \left(\frac{e}{3}\right) : p_T &\approx 0.72 \text{ GeV} \times \frac{5}{m_d(\text{MeV})}. \end{aligned} \quad (22)$$

It seems to be interesting to check these predictions with CMS data in the region of the ridge. In case of confirmation of the effect, measurement of p_T of the photons may give useful information on current masses of light quarks m_u, m_d . Let us remind, that for the moment these parameters are known with considerable uncertainty.




Figure 4: Data driven signal isolation distribution for leading photon obtained using the photon candidates (solid circles) or extrapolated from electrons (continuous line) (extracted from [21] Fig. 2b).

Note that quite recent data from ATLAS [21] may contain hints for this effect. Namely single photon E_T distributions show unexpected fluctuation at $E_T \simeq 5 \text{ GeV}$ (see 4), which could be related

to the first value of (22) (u-quark) with $m_u \simeq 1.45 \text{ MeV}$. Such value of the u-quark mass for energy scale $\simeq 300 \text{ GeV}$ is quite natural due to its RG evolution ²

$$m_u(E) = \frac{m_u(E_0)}{\left(1 + \frac{7\alpha_s(E_0)}{4\pi} \ln\left(\frac{E^2}{E_0^2}\right)\right)^{4/7}}; \quad E_0 = 2 \text{ GeV}. \quad (23)$$

With value $m_u(E_0) = 2.5 \text{ MeV}$ (3) for $E = 300 \text{ GeV}$, $\alpha_s(E_0) = 0.3$ we have $m_u(E) = 1.425 \text{ MeV}$, which just corresponds to 5.1 GeV spikes. Recent results of CMS for two photons [22] also may serve as indications on behalf of the present mechanism – excess for small $\Delta\phi$ of the two photons. Just "CMS ridge effect" for photons.

References

- [1] B.A. Arbuzov, E.E. Boos and V.I. Savrin, Eur. Phys. J. C **71**: 1730 (2011); arXiv:1104.1283[hep-ph].
- [2] CMS collaboration, JHEP **1009**, 091 (2010); arXiv:1009.4122[hep-ex].
- [3] E. Shuryak, arXiv:1009.4635[hep-ph].
- [4] S. M. Troshin, N. E. Tyurin, arXiv:1009.5229[hep-ph].
- [5] A. Dumitru *et al.*, Phys. Lett. B **697**, 21 (2011); arXiv:1009.5295[hep-ph].
- [6] P. Bozek, Eur. Phys. J. C **71**, 1530 (2011); arXiv:1010.0405 [hep-ph].
- [7] I. M. Dremin, V. T. Kim, Pisma ZhETF **92**, 720 (2011); arXiv:1010.0918[hep-ph].
- [8] T. A. Trainor, D. T. Kettler, arXiv:1010.3048[hep-ph].
- [9] I. O. Cherednikov, N. G. Stefanis, arXiv:1010.4463[hep-ph].
- [10] G. A. Kozlov, arXiv:1011.6231[hep-ph].
- [11] A. Kovner and M. Lublinsky, Phys. Rev. D **83**, 034017 (2011); arXiv:1012.3398 [hep-ph].
- [12] K. Werner, Iu. Karpenko, T. Pierog, Phys. Rev. Lett. **106**, 122004 (2011); arXiv:1011.0375[hep-ph].
- [13] M Yu. Azarkin, I. M. Dremin, A. V. Leonidov, arXiv:1102.3258[hep-ph].
- [14] R. C. Hwa and C. B. Yang, Phys. Rev. C **83**, 024911 (2011); arXiv:1011.0965 [hep-ph].
- [15] S. M. Troshin and N. E. Tyurin, arXiv:1103.0626 [hep-ph].
- [16] X. Artru, Phys. Rep. **97**, 147 (1983).
- [17] L. D. Landau and E. M. Lifshitz, Field Theory (FIZMATLIT, Moscow, 2001).
- [18] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
- [19] Dubna-Bucharest-Eotves U.-*et al* collaboration, Yad.Fiz. **16**, 989 (1972).
- [20] P. M. Nadolski *et al.*, Phys.Rev. D **78**, 013004 (2008).
- [21] ATLAS collaboration, arXiv:1107.0581 [hep-ex].
- [22] L.Millischer (CMS collaboration), arXiv:1109.3310 [hep-ex].

²The evolution of $m_q(E)$ practically does not influence results on hadrons due to decreasing of $\alpha_s(E)$.