

$|V_{ub}|$ determination in lattice QCD

IFIC/12-77
 CERN-PH-TH/2012-327
 DESY 12-209
 HU-EP-12/47
 LPT-Orsay/12-117
 MS-TP-12-16
 SFB/PPP-12-87

**F. Bahr^a, F. Bernardoni^{*a}, B. Blossier^b, J. Bulava^c, M. Della Morte^h, P. Fritzsche^e,
 N. Garron^f, A. Gérardin^b, J. Heitger^g, G. von Hippel^d, A. Ramos^a, H. Simma^a,
 R. Sommer^a**

^a NIC, DESY, Platanenallee 6, 15738 Zeuthen, Germany

^b Laboratoire de Physique Théorique, CNRS/Université Paris XI, F-91405 Orsay Cedex, France

^c CERN, Physics Department, TH Division, CH-1211 Geneva 23, Switzerland

^d Institut für Kernphysik, University of Mainz, Becher-Weg 45, 55099 Mainz, Germany

^e Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489 Berlin, Germany

^f School of Mathematics, Trinity College, Dublin 2, Ireland

^g Universität Münster, Institut für Theoretische Physik, Wilhelm-Klemm-Str. 9, 48149 Münster, Germany

^h IFIC, c/ Catedrático José Beltrán 2, E-46980 Paterna - España

fabio.bernardoni@desy.de

The 2012 PDG reports a tension at the level of 3σ between two exclusive determinations of $|V_{ub}|$. They are obtained by combining the experimental branching ratios of $B \rightarrow \tau\nu$ and $B \rightarrow \pi l\nu$ (respectively) with a theoretical computation of the hadronic matrix elements f_B and the $B \rightarrow \pi$ form factor $f_+(q^2)$. To understand the tension, improved precision and a careful analysis of the systematics involved are necessary. We report the results of the ALPHA collaboration for f_B from the lattice with 2 flavors of $O(a)$ improved Wilson fermions. We employ HQET, including $1/m_b$ corrections, with pion masses ranging down to ≈ 190 MeV. Renormalization and matching were performed non-perturbatively, and three lattice spacings reaching $a^{-1} \approx 4.1$ GeV are used in the continuum extrapolation. We also present progress towards a computation of $f_+(q^2)$, to directly compare two independent exclusive determinations of $|V_{ub}|$ with each other and with inclusive determinations. Additionally, we report on preliminary results for f_{B_s} , needed for the analysis of $B_s \rightarrow \mu^+\mu^-$.

36th International Conference on High Energy Physics,
 July 4-11, 2012
 Melbourne, Australia

*Speaker.

1. Motivation

The precise determination of the CKM matrix elements is a key for testing the Standard Model. Violations of CKM unitarity or discrepancies between independent determinations of the same matrix element can provide hints of New Physics. At the time when we started our work, a tension at the level of 3σ between two exclusive determinations of $|V_{ub}|$ existed, as reported e.g. in the PDG of 2012. These determinations use the branching ratio (BR) for the processes $B \rightarrow \pi l \nu$ and $B \rightarrow \tau \nu$ from experiment combined with the form factor $f_+(q^2)$ and the B decay constant f_B , respectively, from the lattice. Also an inclusive determination, based on a perturbative expansion in α_s and an expansion in $1/m_b$, is possible [1]. The results reported by the PDG 2012, computed before ICHEP 2012, can be summarized as follows [2]:

$$\begin{aligned} |V_{ub}| &= 0.00323(31) & (B \rightarrow \pi l \nu), & & |V_{ub}| &= 0.00510(47) & (B \rightarrow \tau \nu), & & (1.1) \\ |V_{ub}| &= 0.00441(34) & (\text{inclusive}). & & & & & & \end{aligned}$$

At ICHEP 2012 the Belle collaboration reported a new result for $\text{BR}(B \rightarrow \tau \nu)$ [3] based on a new set of data, obtained with a more sophisticated tagging of the B. This result, taken alone, would yield a value for $|V_{ub}|$ that is consistent with the exclusive determination from $B \rightarrow \pi$. However, more data and a careful inspection of the systematics involved are needed to draw more definitive conclusions.

While the experimental precision in the differential decay rate for $B \rightarrow \pi l \nu$ has by now reached good precision, $B \rightarrow \tau \nu$ events are more difficult to reconstruct and there is an error of the order of 20% on the branching ratio. The situation on the theoretical side is the opposite: the lattice computation of a form factor is more challenging than that of a decay constant.

At this conference we have presented the results for the determination of f_B by the ALPHA collaboration which use fully non-perturbative renormalization and matching, and CLS configurations with two degenerate dynamical quarks in the sea. A parallel effort to determine $f_+(q^2)$ in the same setup is ongoing: we have presented the progress reached so far, and the precision that we expect to achieve, once the full non-perturbative renormalization and matching at order $1/m_b$ have been completed. The comparison of these two exclusive predictions, in which the relevant hadronic parameters have been computed in the same setup, will provide a test as free as possible from systematics.

Recently LHCb has presented the first evidence for $B_s \rightarrow \mu^+ \mu^-$, with a decay rate compatible with the Standard Model [4]. We have presented the determination by the ALPHA collaboration of the B_s decay constant, f_{B_s} , that enters in the theoretical prediction of this decay.

2. HQET

Our computations were performed on CLS configurations, which have two degenerate $O(a)$ improved Wilson quarks. The ensembles used in this work have pion masses m_π in the range $190 \text{ MeV} \lesssim m_\pi \lesssim 450 \text{ MeV}$ at three lattice spacings a , namely $a \in \{0.078, 0.065, 0.045\} \text{ fm}$. All of them have a spatial extent L such that $m_\pi L > 4$, so that volume effects are expected to be very small.

Even for our finest lattice spacing the b quark cannot be simulated directly, given that $am_b > 1$.

However, for the low energy processes we are interested in, the m_b scale can be integrated out. Our approach is to use HQET, which is an expansion of the QCD Lagrangian in powers of $1/m_b$. At leading order the b quark is static, i.e. the Lagrangian involves no space derivatives. If we include terms up to order $1/m_b$ the Lagrangian becomes:

$$\begin{aligned}\mathcal{L}_{\text{HQET}}(x) &= \bar{\psi}_h(x) D_0 \psi_h(x) - \omega_{\text{kin}} \mathcal{O}_{\text{kin}}(x) - \omega_{\text{spin}} \mathcal{O}_{\text{spin}}(x), \\ \mathcal{O}_{\text{kin}}(x) &= \bar{\psi}_h(x) \mathbf{D}^2 \psi_h(x), \quad \mathcal{O}_{\text{spin}}(x) = \bar{\psi}_h(x) \boldsymbol{\sigma} \cdot \mathbf{B} \psi_h(x).\end{aligned}\tag{2.1}$$

The corresponding expansion of the time component A_0 of the heavy-light current (at $p = 0$) is

$$A_{0,R}^{\text{HQET}} = Z_A^{\text{HQET}} \left[A_0^{\text{stat}} + c_A^{(1)} A_0^{(1)} \right], \quad A_0^{\text{stat}} = \bar{\psi}_1 \gamma_0 \gamma_5 \psi_h, \quad A_0^{(1)} = \bar{\psi}_1 \gamma_5 \gamma_i \frac{1}{2} (\overleftrightarrow{\nabla}_i^S - \overleftarrow{\nabla}_i^S) \psi_h.\tag{2.2}$$

Since in HQET the $\mathcal{O}(1/m_b)$ terms appear only as insertions in correlation functions, HQET is renormalizable order by order in $1/m_b$ because the static theory is. Once the HQET parameters $\omega_i \in \{m_{\text{bare}}, Z_A^{\text{HQET}}, c_A^{(1)}, \omega_{\text{kin}}, \omega_{\text{spin}}\}$ have been determined using *non-perturbative* matching [5], the continuum limit can therefore be taken safely.

3. Matching

The matching was performed in the Schrödinger Functional scheme in a small volume $L_1 \approx 0.4$ fm, where $am_b \ll 1$, and relativistic b-quarks can be simulated [6]. The HQET parameters can then be fixed by imposing the matching conditions for suitable observables Φ

$$\Phi^{\text{HQET}}(L_1, M, a) = \Phi^{\text{QCD}}(L_1, M, 0), \quad \Phi^{\text{QCD}}(L_1, M, 0) = \lim_{a \rightarrow 0} \Phi^{\text{QCD}}(L_1, M, a),\tag{3.1}$$

and hence the ω_i inherit their dependence on the heavy-quark mass M from QCD. In particular here M is the RGI mass of the b-quark [7]. Using finite-size scaling recursively, we can then take the step $L_1 \rightarrow L_2 = 2L_1$, and finally connect with large volumes $L_\infty \gtrsim \max(2 \text{ fm}, 4/m_\pi)$.

Having performed this matching procedure for each of the lattice spacings used in our large-volume simulations, we know the corresponding $N_f = 2$ HQET parameters $\omega_i(L_1, M, a)$ non- for a range of values of M in the neighbourhood of the b quark mass.

4. Results

The HQET energies and matrix elements are extracted at large Euclidean time t separations. The effects of excited states are exponentially suppressed like $\sim e^{-(E_2 - E_1)t}$, where E_2 and E_1 are the energies of the first excited state and the ground state, respectively. To achieve a better suppression, we solve the Generalized Eigenvalue Problem (GEVP) [8]

$$C(t)v_n(t, t_0) = \lambda_n(t, t_0)C(t_0)v_n(t, t_0), \quad t_0 < t < 2t_0,\tag{4.1}$$

for an $N \times N$ correlator matrix $C(t)$ with $N = 3$. Each entry of the matrix corresponds to a different Gaussian smearing level of the light quark field in the B-meson interpolating quark bilinear. The corrections to the energies and matrix elements so obtained behave like $\propto \exp\{-(E_{N+1} - E_1)t\}$ and




Figure 1: *Left:* HM χ PT extrapolation of f_B ; *centre/right:* linear extrapolation of f_B and f_{B_s} . The blue, red and green points correspond to ensembles at $a = 0.075$ fm, 0.065 fm and 0.048 fm, respectively. The fit formulae evaluated at each given lattice spacing are shown in colour. The black curve is the chiral and continuum extrapolation. It is a fit to all the points shown in the figure plus other at the same pion masses but obtained using a different discretization for the heavy quark action.

$\propto \exp\{-(E_{N+1} - E_1)t_0\} \times \exp\{-(E_2 - E_1)(t - t_0)\}$, respectively. The residual systematic errors are kept under control by requiring $\sigma_{\text{stat}} \gtrsim 3\sigma_{\text{sys}}$.

In phenomenological predictions, we need to know the HQET parameters at the physical mass of the b quark $\omega_i(L_1, M_b, m_\pi^{\text{exp}}, a = 0) \equiv m_B^{\text{exp}} = 5279.5$ MeV. The mass of the B at physical pion mass m_π^{exp} is obtained through a chiral and continuum extrapolation [9]:

$$m_B(z, m_\pi, a, n) = B(z) + Cm_\pi^2 - \frac{3\hat{g}^2}{16\pi f_\pi^2} m_\pi^3 + D_n a^2, \quad \hat{g} = 0.51(2) [10]. \quad (4.2)$$

In HQET the B-meson mass is given by $m_B = m_{\text{bare}} + E^{\text{stat}} + \omega_{\text{kin}} E^{\text{kin}} + \omega_{\text{spin}} E^{\text{spin}}$, so that

$$M_b = 6.56(15)(06)_z \text{ GeV}, \quad \text{or equivalently} \quad \bar{m}_b^{\overline{MS}}(\bar{m}_b) = 4.22(10)(4)_z \text{ GeV}. \quad (4.3)$$

For the following analyses, we use the values of HQET parameters obtained from an interpolation to $z \equiv z_b$.

4.1 B and B_s decay constants

Using the HQET formulas to combine the matrix elements and energies with the matching parameters at $z \equiv z_b$ we can compute f_B and f_{B_s} through a chiral and continuum extrapolation dictated by HMChPT [11, 12]:

$$f_B(m_\pi, a, n) = b \left[1 - \frac{3}{4} \frac{1 + 3\hat{g}^2}{(4\pi f_\pi)^2} m_\pi^2 \ln(m_\pi^2) \right] + cm_\pi^2 + d_n a^2. \quad (4.4)$$

Our analysis for ICHEP 2012 gives

$$f_B = 193(9)_{\text{stat}}(4)_\chi \text{ MeV}, \quad f_{B_s} = 219(12)_{\text{stat}} \text{ MeV},$$

where the error coming from the chiral extrapolation is determined by comparing to a linear extrapolation in m_π (see Fig. 1). For f_{B_s} not all ensembles are analysed yet. For more details, see [13]. Our value for f_B is compatible with the values found by other collaborations [14, 15, 16, 17].

4.2 $B \rightarrow \pi$ form factor

The form factor $f_+(q^2)$ is defined through the Lorentz decomposition of the matrix element:

$$\langle \pi(p_\pi) | V^\mu | B(p_B) \rangle = f_+(q^2) \left[p_B^\mu + p_\pi^\mu - \frac{m_B^2 - m_\pi^2}{q^2} q^\mu \right] + f_0(q^2) \frac{m_B^2 - m_\pi^2}{q^2} q^\mu, \quad (4.5)$$

where $q^\mu = p_B^\mu - p_\pi^\mu$. To extract this matrix element from lattice simulations, we consider the ratio:

$$R(t_\pi, t_B) \equiv \frac{\sum_{\vec{x}_\pi, \vec{x}_B} e^{-i\vec{p} \cdot \vec{x}_\pi} \langle P_{ll}(t_\pi + t_B, \vec{x}_\pi) V^\mu(t_B, \vec{x}_B) P_{hl}(0) \rangle}{\sqrt{\sum_{\vec{x}_\pi} e^{-i\vec{p} \cdot \vec{x}_\pi} \langle P_{ll}(x_\pi) P_{ll}(0) \rangle \sum_{\vec{x}_B} \langle P_{hl}(x_B) P_{hl}(0) \rangle}},$$

$$\langle \pi(p_\pi) | V^\mu | B(p_B) \rangle = \lim_{T, t_B, t_\pi \rightarrow \infty} R(t_\pi, t_B) e^{E_\pi t_\pi / 2 + m_B t_B / 2} \quad (4.6)$$

where P_{ll} and P_{hl} are interpolating operators for the π and the B meson, respectively.

One additional difficulty in comparison to the extraction of decay constants is the presence of large finite T effects, where T is the temporal extension of the lattice. So far we have restricted to few lattices and to the static limit to demonstrate the feasibility of this computation in our setup. Our results show that finite T effects can be understood in the transfer matrix formalism (see Fig. 2) and that a precision of 5% is achievable.

Combining our result at largest q^2 , $q = p_\pi - p_B$, with the experimental data available and a parametrization of the q^2 dependence based on very general properties like unitarity and analyticity [18], $|V_{ub}|$ is obtained with a 15% precision. This does not include the necessary extrapolation in the light quark mass and the lattice spacing. These missing steps will soon be coming out. For more details, see [19].

Acknowledgments

This work is supported in part by the SFB/TR 9 and grant HE 4517/2-1 (P.F. and J.H.) of the Deutsche Forschungsgemeinschaft and by the European Community through EU Contract MRTN-CT-2006-035482, ‘‘FLAVIANet’’. We thank our colleagues in the CLS effort for the joint production and use of gauge configurations. We gratefully acknowledge the computer resources provided within the Distributed European Computing Initiative by the PRACE-2IP, with funding from the European Community’s Seventh Framework Programme (FP7/2007-2013) under grant agreement RI-283493, by the Grand Équipement National de Calcul Intensif at CINES in Montpellier, and by the John von Neumann Institute for Computing at FZ Jülich, at the HLRN in Berlin, and at DESY, Zeuthen.




Figure 2: t_π dependence for the considered ratio eq. (4.6) on one of our lattices with $m_\pi \approx 300$ MeV. We use a smeared interpolating operator for the B and the π . The red curve is obtained by fitting the finite T effects with the prediction from transfer matrix theory: $R(t_\pi, t_B) = \frac{A + B t_B e^{-E_\pi(T-2t_\pi)}}{\sqrt{1 + e^{-E_\pi(T-2t_\pi)}}$ with A and B as parameters.

References

- [1] I. I. Bigi, M. A. Shifman, N. Uraltsev, and A. I. Vainshtein, “QCD predictions for lepton spectra in inclusive heavy flavor decays,” *Phys.Rev.Lett.*, vol. 71, pp. 496–499, 1993.
- [2] J. Beringer *et al.*, “Review of Particle Physics (RPP),” *Phys.Rev.*, vol. D86, p. 010001, 2012.
- [3] Y. Y. for the Belle collaboration, “Leptonic and semileptonic B decays at Belle,” *talk at ICHEP*, 2012.
- [4] R. Aaij *et al.*, “First evidence for the decay $B_s \rightarrow \mu^+ \mu^-$,” 2012.
- [5] J. Heitger and R. Sommer, “Nonperturbative heavy quark effective theory,” *JHEP*, vol. 0402, p. 022, 2004.
- [6] B. Blossier *et al.*, “Parameters of Heavy Quark Effective Theory from Nf=2 lattice QCD,” *JHEP*, vol. 1209, p. 132, 2012.
- [7] P. Fritzsch, J. Heitger, and N. Tantalo, “Non-perturbative improvement of quark mass renormalization in two-flavour lattice QCD,” *JHEP*, vol. 1008, p. 074, 2010.
- [8] B. Blossier, M. Della Morte, G. von Hippel, T. Mendes, and R. Sommer, “On the generalized eigenvalue method for energies and matrix elements in lattice field theory,” *JHEP*, vol. 0904, p. 094, 2009.
- [9] F. Bernardoni, P. Hernandez, and S. Necco, “Heavy-light mesons in the epsilon-regime,” *JHEP*, vol. 1001, p. 070, 2010.
- [10] J. Bulava, M. Donnellan, and R. Sommer, “The $B^* B \pi$ Coupling in the Static Limit,” *PoS*, vol. LATTICE2010, p. 303, 2010.
- [11] S. R. Sharpe and Y. Zhang, “Quenched chiral perturbation theory for heavy - light mesons,” *Phys.Rev.*, vol. D53, pp. 5125–5135, 1996.
- [12] J. Goity, “Chiral perturbation theory for SU(3) breaking in heavy meson systems,” *Phys.Rev.*, vol. D46, pp. 3929–3936, 1992.
- [13] F. Bernardoni, B. Blossier, J. Bulava, M. Della Morte, P. Fritzsch, *et al.*, “B-physics from HQET in two-flavour lattice QCD,” 2012.
- [14] H. Na, C. J. Monahan, C. T. Davies, R. Horgan, G. P. Lepage, *et al.*, “The B and B_s Meson Decay Constants from Lattice QCD,” *Phys.Rev.*, vol. D86, p. 034506, 2012.
- [15] A. Bazavov *et al.*, “B- and D-meson decay constants from three-flavor lattice QCD,” *Phys.Rev.*, vol. D85, p. 114506, 2012.
- [16] C. McNeile, C. Davies, E. Follana, K. Hornbostel, and G. Lepage, “High-Precision f_{B_s} and HQET from Relativistic Lattice QCD,” *Phys.Rev.*, vol. D85, p. 031503, 2012.
- [17] P. Dimopoulos *et al.*, “Lattice QCD determination of m_b , f_B and f_{B_s} with twisted mass Wilson fermions,” *JHEP*, vol. 1201, p. 046, 2012.
- [18] C. Bourrely, I. Caprini, and L. Lellouch, “Model-independent description of $B \rightarrow \pi l \nu$ decays and a determination of $|V_{ub}|$,” *Phys.Rev.*, vol. D79, p. 013008, 2009.
- [19] F. Bahr, F. Bernardoni, A. Ramos, H. Simma, R. Sommer, *et al.*, “ $B \rightarrow \pi$ form factor with 2 flavours of $O(a)$ improved Wilson quarks,” 2012.