

## Analysis of the new INTEGRAL Earth observations to measure the cosmic X-ray background

---

M. Türler\*, N. Produit, L. Pavan, C. Ferrigno, and P. Bordas†

*ISDC & Observatory of the University of Geneva, ch. d'Ecogia 16, 1290 Versoix, Switzerland*

*E-mail: marc.turler@unige.ch*

A new series of Earth occultation observations have been started in 2012 to refine the determination of the cosmic X-ray background by the *INTEGRAL* mission. We show here that the new detector lightcurves in the 3 to 160 keV range differ from the ones obtained in 2006. Instead of the expected modulation induced by the passage of the Earth through the field of view of the JEM-X, IBIS/ISGRI and SPI instruments, we record unrelated variability on shorter timescales. We discuss the differences obtained with the datasets of 2006 and 2012 in view of the changes in pointing direction, spacecraft orbit and solar cycle phase. We conclude that the Earth occultation signal in 2012 is likely blended by radioactive decay resulting from the activation of the spacecraft when crossing the proton radiation belt at perigee passage. The observed variability, on the other hand, results most likely from the current solar maximum. In addition to a variable particle environment from inhomogeneities of the solar wind, we also find evidence for hard X-ray auroral emission. While the former can be traced by SPI/ACS counts, the latter – by enhancing unpredictably the Earth emission – is a major disturbance for measuring the diffuse X-ray background through occultation by the Earth.

*"An INTEGRAL view of the high-energy sky (the first 10 years)" 9th INTEGRAL Workshop and celebration of the 10th anniversary of the launch,*

*October 15-19, 2012*

*Bibliothèque Nationale de France, Paris, France*

---

\*Speaker.

†also at: Institut für Astronomie und Astrophysik, Universität Tübingen, Sand 1, 72076 Tübingen, Germany

## 1. Introduction

The idea to use the Earth disk as a shield to occult the cosmic X-ray background (CXB) was evaluated early in the *INTEGRAL* mission and was implemented in January and February 2006. At the start of four spacecraft revolutions, the Earth was allowed to cross the field of view (FoV) of the instruments aimed at a fixed position in the sky. The occultation of the CXB by the Earth passage resulted in a characteristic signal, a hollow, in the evolution of the detector count rates. The depth of the signal modulation in different energy bands can then be used to extract the spectrum of the CXB. Based on the successful performance and analysis of these very unconventional observations [1, 2], it has been proposed in 2010 to repeat the experiment.


The new set of observations was aimed at deriving the spectrum of the CXB in the hard X-ray domain with improved statistics and less systematics. The idea was to gain a factor of two in statistical uncertainties by multiplying by four the number of occultations. The gain in systematics was expected by performing them in a FoV not affected by the presence of the Galactic ridge and bright point sources as it was the case in 2006. As the bombardment of the Earth by cosmic rays is reduced during solar maximum, it was also anticipated to have a lower emission by the Earth atmosphere above 70 keV in 2012–2014 with respect to the solar minimum of 2006 [4]. This strong scientific case and the evaluation of the feasibility of the observations by the staff of the Mission Operation Centre [3] led to the recommendation by the *INTEGRAL* Users Group to perform the proposed observation program.

Four new Earth occultation observations have been conducted from May 2012 to September 2012. We present here these observations for all three high-energy instruments aboard *INTEGRAL*. The obtained data presented in Sect. 2 do unfortunately not match the expectations for various reasons that we discuss in Sect. 3.

## 2. Data

The first four new *INTEGRAL* Earth-occultation observations (EOs) of 2012 have been conducted at the start of the spacecraft revolutions 1169 (May 10), 1197 (Aug. 01), 1206 (Aug. 28) and 1216 (Sep. 27). We refer to them as EO 2.1, 2.2, 2.3 and 2.4, respectively. We extracted full detector lightcurves for all three high-energy instruments aboard *INTEGRAL* [5]. We used version 10 of the Off-line Scientific Analysis (OSA 10) provided by the ISDC [6] with a small change for the JEM-X instrument to allow the analysis of slew observations rather than pointings. For IBIS/ISGRI we analysed the data until the deadtime correction level and extracted the lightcurves with the `ii_light` executable included in the OSA package in time bins of 300 s. For SPI, we simply selected the events of all live detectors based on time and energy.

Figure 1 shows one of the new sets of detector lightcurves (EO 2.2) compared to those obtained the same way using the data of EO 1.1 in revolution 401 (Jan. 2006). The drop in the count rates resulting from the passage of the Earth through the FoV of the instruments is clearly visible in the lowest energy lightcurves of each instrument for EO 1.1, but is not seen for EO 2.2. For the latter observation, a CXB occultation might be present in JEM-X at the very beginning, before a strong emission bump arises in the signal. The SPI and IBIS lightcurves at energies below 60 keV show correlated variability which is not matching the evolution of the SPI/ACS counts.




**Figure 1:** Comparison of JEM-X, IBIS/ISGRI and SPI detector lightcurves for Earth observations EO 1.1 (upper panel) and EO 2.2 (lower panel). For each instrument, we show six average-subtracted lightcurves in different energy bands with energies increasing from the top curve (red) to the bottom curve (magenta). The SPI lightcurves have also been divided by the square-root of the average count rate to better match JEM-X and IBIS rates. JEM-X lightcurves are the average of JEM-X 1 and 2 in the energy bands: 3–5, 5–8, 8–12, 12–18, 18–24, 24–30 keV. ISGRI and SPI bands are: 20–28, 28–38, 38–52, 52–72, 72–100, 100–160 keV and 25–30, 30–35, 35–40, 40–45, 45–50, 50–55 keV, respectively. The simultaneous SPI/ACS lightcurve variability is also shown as a black line for comparison with IBIS counts.

The detector lightcurves of the three other EOs of 2012 do also not display the expected occultation modulation, as shown in Fig. 2. While the lightcurves of EO 2.3 are relatively stable, those of EO 2.1 and EO 2.4 are affected by strong variability also visible in the SPI/ACS count rates.

### 3. Discussion

The diversity of the detector lightcurves of the EOs of 2012 and the absence of the expected CXB occultation signal present in the data of 2006 is puzzling. In order to understand the problem,



**Figure 2:** Comparison of JEM-X (upper panels) and IBIS/ISGRI (lower panels) detector lightcurves for Earth observations EO 2.1, EO 2.3 and EO 2.4. The energy bands and colour coding is as in Fig. 1.

we discuss below the possible effect of three main differences between the EOs of 2012 and the ones of 2006: a change in pointing direction, in spacecraft orbit, and in the solar cycle phase.

### 3.1 Effect of the pointing direction

The EOs of 2006 were directed at a Galactic longitude  $l \approx -30^\circ$  not far from the Galactic bulge and including the Galactic ridge X-ray emission (GRXE) in the partially coded FoV of IBIS. This was not very favorable, as it resulted also in the occultation of the GRXE and of bright point sources in addition to the CXB occultation. The new observations are still pointing close to the Galactic plane, but at  $l \approx -110^\circ$  where the GRXE is much reduced [7].

This better configuration with respect to the background sky could however imply an increase of the albedo of the Earth by atmospheric reflection of the Galactic bulge X-ray emission. The time-averaged contribution of all point sources in the central radian of the Milky Way ( $|l| < 30^\circ$  and  $|b| < 15^\circ$ ) has been derived from SPI observations to be a powerlaw above 20 keV with a photon index  $\Gamma = 2.67$  and a flux at 50 keV of  $2.47 \times 10^{-3} \text{ ph cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$  [8]. As this normalization is just about the flux of the Crab Nebula at 50 keV – which is located in the opposite direction on the sky – the Earth albedo due to point sources should be roughly constant for any pointing direction. Point sources contribute thus only an additional  $\sim 2\text{--}3\%$  (1/40) to the albedo of the Earth from CXB reflexion, as estimated with the Crab flux for the EOs of 2006 [1].

### 3.2 Effect of the spacecraft orbit

Another difference with respect to the EOs of 2006, is the change in the orbit of the spacecraft. The perigee height during the first four EOs of 2012 was only between  $\sim 3300$  km and  $\sim 3800$  km, whereas it was at  $12'600$  km for the EOs of 2006. As a consequence, *INTEGRAL* is currently passing through the proton radiation belt at perigee, which leads to activation of the spacecraft material that is then radioactively decaying during the post-perigee EOs. This leads to an exponential decay of the instrumental background that is clearly present in the SPI/ACS detector count rates and in the highest energy bands of IBIS/ISGRI (see Figs.1 & 2).

In parallel to this change of perigee height, the orbit precessed such that in 2012 the exit of the electron radiation belt is at about twice higher altitude,  $\sim 50'000$  km, than the belt entrance height, whereas in 2006 we had the opposite situation with an exit at  $\sim 35'000$  km and an entrance at  $\sim 60'000$  km. The higher exit height in 2012 delays the start of the EOs after perigee and thus results in a slightly smaller Earth size and a slower passage of the Earth through the FoV of the instruments. As a consequence, the EOs of 2012 do only start when the Earth is already close to the centre of the IBIS FoV and their duration is about twice longer ( $\sim 16$ h) than in 2006 ( $\sim 8$ h).

The overall consequence of this change of orbit for post-perigee EOs is that the expected Earth modulation signal in IBIS and SPI is basically a continuous increase of the count rates during the observations. This rise of detector counts can be easily blended by the opposite decrease of counts resulting from the radioactive decay of the spacecraft. We would need to analyse several beginning of revolutions – preferentially in clear extragalactic fields – to evaluate if the contribution of the radioactive decay of the spacecraft to the detector lightcurves in different energy bands can be uniquely assessed and removed from the EOs. This would also need to be done for JEM-X which is known to have a variable gain in the first few science windows after perigee passage. The possibility of performing pre-perigee observations would solve all these issues.

### 3.3 Effect of the solar cycle

A third difference with respect to the EOs of 2006, is that we are now close to solar maximum. This was initially seen positively (see Sect. 1), but turns out to result in complex variability in the detector counts. Variations in the solar wind experienced by *INTEGRAL* result in a variable instrumental background that can be traced by the SPI/ACS detector and the IREM electron count rates. This kind of variability does strongly affect EOs 2.1 and 2.4 (see Fig. 2).

The variability seen in EO 2.2 (Fig. 1) is likely of another kind. In the JEM-X bands it seems quite clear that we observe an event of strong auroral activity [9] with a soft spectrum creating the bump of emission when the Earth polar regions are in the FoV of the instrument. The slight enhancement seen quasi-simultaneously in the lowest energy bands of IBIS and SPI might be related. The other variability event in the second part of the EO is only seen in IBIS and SPI, but not in the SPI/ACS and in JEM-X, suggesting that it is not instrumental, but rather also auroral emission that occurred when the Earth has already left the smaller FoV of JEM-X. We note that the recent EO 2.5 (not presented here) shows a similar behaviour with much higher amplitude. It might be of interest to study hard X-ray auroral events from space, although their variability and spectral characteristics are already well known from balloon experiments back in the 1960s [10].

Concerning the extraction of the CXB spectrum, aurorae are clearly a serious nuisance, especially if their variability timescale is of the order of several hours, as is the occultation signal.

#### 4. Conclusion

The detector lightcurves of the four first EOs of 2012 do not show the expected modulation from the passage of the Earth through the FoV of the instruments, thus preventing us from extracting the spectrum of the CXB. The occultation signal is likely blended by radioactive spacecraft decay after radiation belt exit. This issue could be solved by performing pre-perigee observations. Unfortunately, the data are also contaminated by variations due to the current strong solar activity. This has both an effect on the instrumental background due to a variable solar wind, and on the auroral X-ray emission of the Earth. The latter effect might be interesting in itself, but is a major issue for determining the CXB spectrum by Earth occultations at solar maximum.

#### Acknowledgments

We are very grateful to the *INTEGRAL* staff at ISOC and MOC for scheduling and performing these unconventional observations.

#### References

- [1] Churazov, E., Sunyaev, R., Revnivtsev, M., et al., *INTEGRAL observations of the cosmic X-ray background in the 5-100 keV range via occultation by the Earth*, 2007, A&A, 467, 529
- [2] Türler, M., Chernyakova, M., Courvoisier, T. J.-L., et al., *INTEGRAL hard X-ray spectra of the cosmic X-ray background and Galactic ridge emission*, 2010, A&A, 512, A49
- [3] Hübner, J. M., Lozano, C., & Southworth, R., *INTEGRAL Earth occultation observations revealing the soft gamma ray background – The operational view*, in proceedings of *The Extreme and Variable High Energy Sky 2011*, PoS (Extremesky 2011) 059
- [4] Sazonov, S., Churazov, E., Sunyaev, R., & Revnivtsev, M., *Hard X-ray emission of the Earth's atmosphere: Monte Carlo simulations*, 2007, MNRAS, 377, 1726
- [5] Winkler, C., Courvoisier, T. J.-L., Di Cocco, G., et al., *The INTEGRAL mission*, 2003, A&A, 411, L1
- [6] Courvoisier, T. J.-L., Walter, R., Beckmann, V., et al., *The INTEGRAL Science Data Centre (ISDC)*, 2003, A&A, 411, L53
- [7] Krivonos, R., Tsygankov, S., Revnivtsev, M., et al., *INTEGRAL constraints on the Galactic hard X-ray background from the Milky Way anticenter*, 2012, A&A, 537, 92
- [8] Bouchet, L., Jourdain, E., Roques, J.-P., et al., *INTEGRAL SPI All-Sky View in Soft Gamma Rays: A Study of Point-Source and Galactic Diffuse Emission*, 2008, ApJ, 679, 1315
- [9] Bhardwaj, A., Elsner, R. F., Randall Gladstone, G., et al., *X-rays from solar system objects*, 2007, P&SS, 55, 1135
- [10] Barcus, J. R., & Rosenberg, T. J., *Energy Spectrum of Auroral-Zone X Rays*, 1966, J. Geophys. Res., 71, 803