

Charm quark mass and D-meson decay constants from two-flavour lattice QCD



Jochen Heitger*

Westfälische Wilhelms-Universität Münster, Institut für Theoretische Physik,
Wilhelm-Klemm-Straße 9, D-48149 Münster, Germany

E-mail: heitger@uni-muenster.de

Georg M. von Hippel

PRISMA Cluster of Excellence,
Johannes Gutenberg Universität Mainz, Institut für Kernphysik,
Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany

E-mail: hippel@kph.uni-mainz.de

Stefan Schaefer†

CERN, Physics Department, TH Unit, CH-1211 Geneva 23, Switzerland

E-mail: stschaef@mail.cern.ch

Francesco Virota

NIC, DESY, Platanenallee 6, D-15738 Zeuthen, Germany

We present a computation of the charm quark's mass and the leptonic D-meson decay constants f_D and f_{D_s} in two-flavour lattice QCD with non-perturbatively $O(a)$ improved Wilson quarks. Our analysis is based on the CLS configurations at two lattice spacings ($a = 0.065$ and 0.048 fm, where the lattice scale is set by f_K) and pion masses ranging down to ~ 190 MeV at $Lm_\pi \gtrsim 4$, in order to perform controlled continuum and chiral extrapolations with small systematic uncertainties.

CLS
based

31st International Symposium on Lattice Field Theory LATTICE 2013

July 29 - August 3, 2013

Mainz, Germany

*Speaker.

†Address since 1. November 2013: NIC, DESY, Platanenallee 6, D-15738 Zeuthen, Germany

id	L/a	β	a [fm]	κ_l	κ_s	m_π [MeV]	# cfigs
E5	32	5.3	0.065	0.136250	0.135777(17)	440	2×199
F6	48			0.136350	0.135741(17)	310	250
F7	48			0.136380	0.135730(17)	270	547
G8	64			0.136417	0.135705(17)	190	820
N5	48	5.5	0.048	0.136600	0.136262(08)	440	238
N6	48			0.136670	0.136250(08)	340	1000
O7	64			0.136710	0.136243(08)	270	486

Table 1: Parameters of the ensembles used: ensemble label, spatial extent of the lattice in lattice units, bare coupling $\beta = 6/g_0^2$, lattice spacing a , hopping parameters of the sea and (valence) strange quarks, the mass of the sea pion and the number of configurations employed. All lattices have dimensions $T \times L^3$ with $T = 2L$.

1. Introduction

The decay constants f_D and f_{D_s} entering the Standard Model expression for the leptonic decay widths of charged D-mesons are given by the non-perturbative QCD matrix elements

$$\langle 0 | \bar{q} \gamma_\mu \gamma_5 c | D_q(p) \rangle = i f_{D_q} p_\mu, \quad q = d, s. \quad (1.1)$$

Besides the importance of f_{D_q} in extracting $|V_{cq}|$ and overconstraining the CKM matrix, there is continuing phenomenological interest in these decays, because a significant deviation between experimental and lattice results for f_{D_q} could hint at New Physics in the flavour sector [1]. Extending our previous work [2] with an expanded set of ensembles with higher statistics, lighter sea quarks and largely reduced systematics (such as the uncertainty in the lattice scale), we here report results of a preliminary analysis of $f_{D(s)}$ and of the charm quark mass.

2. Computational setup and techniques

Our measurements were performed on a subset of the Coordinated Lattice Simulations (CLS) gauge field ensembles, which are characterized by the Wilson plaquette gauge action and a sea of $N_f = 2$ mass degenerate flavours of non-perturbatively $O(a)$ improved Wilson quarks. The sea quark masses span a range corresponding to pion masses ($190 \lesssim m_\pi \lesssim 440$) MeV, while the strange valence quark is fixed to its physical value and we scan a range of charm valence quark masses around the physical charm quark mass. Two values of the lattice spacing are considered: $a \in \{0.065 \text{ fm}, 0.048 \text{ fm}\}$. Suppression of finite-size effects is ensured by restricting ourselves to ensembles with $Lm_\pi \gtrsim 4$. To generate the gauge configurations, either the DD-HMC [3–6] or the MP-HMC based on mass preconditioning [7, 8] was employed. The parameters of the ensembles used are summarized in table 1. Statistical errors are estimated via a jackknife procedure, but will be double-checked in the final analysis by the method of [9] studying autocorrelation functions.

The lattice spacings a , pion masses m_π , pion decay constants f_π and values of the (quenched) strange quark's hopping parameter κ_s are inferred from [10], where the scale is set through f_K at the “physical point” defined by $m_{\pi,\text{phys}} = 134.8 \text{ MeV}$, $m_{K,\text{phys}} = 494.2 \text{ MeV}$ and $f_{K,\text{phys}} = 155 \text{ MeV}$ in the isospin-symmetric limit with QED effects removed. For two ensembles (G8,N6), we had to perform a short linear interpolation in $1/\kappa_s$ (at fixed charmed κ_c) to the physical value.

As for the hopping parameter of the (quenched) charm quark, κ_c , we fix it by requiring the D_s -meson mass to acquire its physical value, $m_{D_s} = m_{D_s,\text{phys}} = 1968 \text{ MeV}$, irrespective of the sea quark mass. To this end, after choosing a few values in the vicinity of this target, $(am_{D_s})^2$ was interpolated linearly in $1/\kappa_c$ to $(am_{D_s,\text{phys}})^2$.

3. Observables and analysis details

For two mass non-degenerate valence quarks r and s , we compute correlators of the pseudoscalar density $P^{rs} = \bar{\psi}_r \gamma_5 \psi_s$ and the time component of the axial vector current $A_0^{rs} = \bar{\psi}_r \gamma_0 \gamma_5 \psi_s$ as

$$f_{\text{PP}}^{rs}(x_0) = -a^3 \sum_{\vec{x}} \langle P^{rs}(x) P^{sr}(0) \rangle, \quad f_{\text{AP}}^{rs}(x_0) = -a^3 \sum_{\vec{x}} \langle A_0^{rs}(x) P^{sr}(0) \rangle. \quad (3.1)$$

These are evaluated using 10 $U(1)$ noise sources $\eta_t(x) = \delta_{t,x_0} \exp(i\phi(\vec{x}))$ located on randomly chosen time slices t [11, 12] so that solving the Dirac equation once for each noise vector $\zeta_t^r = Q^{-1}(m_{0,r})\eta_t = a^{-1}(D + m_{0,r})^{-1}\gamma_5\eta_t$ suffices to estimate the two-point functions projected onto zero momentum:

$$a^3 f_{\text{XP}}^{rs}(x_0) = \sum_{\vec{x}} \langle [\zeta_t^r(x_0 + t, \vec{x})]^\dagger \Gamma \zeta_t^s(x_0 + t, \vec{x}) \rangle, \quad \Gamma = \mathbf{1}, \gamma_0 \text{ for } X = \text{P, A}, \quad (3.2)$$

where the average is over noise sources and gauge configurations.

The $O(a)$ improved effective average PCAC quark mass of flavours r and s is now defined as

$$\frac{1}{2} (m_{rr} + m_{ss})(x_0) = m_{rs}(x_0) = \frac{\frac{1}{2} (\partial_0 + \partial_0^*) f_{\text{AP}}(x_0) + c_A a \partial_0^* \partial_0 f_{\text{PP}}(x_0)}{2 f_{\text{PP}}(x_0)}, \quad (3.3)$$

where the improvement coefficient c_A is non-perturbatively known from [13]. For sufficiently large x_0 , $m_{rs}(x_0)$ exhibits a plateau, over which we take a timeslice average to calculate m_{rs} . Examples for two representative ensembles and various valence κ -combinations are shown in figure 1.

The renormalized PCAC mass is then given by

$$m_{\text{R}}^{rs} = \frac{Z_A (1 + \bar{b}_A a m_{\text{sea}} + \tilde{b}_A a m_{rs})}{Z_P (1 + \bar{b}_P a m_{\text{sea}} + \tilde{b}_P a m_{rs})} \times m_{rs}, \quad m_{\text{sea}} = m_{\text{ll}} \text{ with } l = \text{l(ight)}, \quad (3.4)$$

where Z_A and Z_P assume values from the non-perturbative determinations in [10, 14–16]). The b -coefficients, multiplying (very small) improvement terms, are known in 1-loop perturbation theory [10, 17]; in particular, $\bar{b}_A = \bar{b}_P = 0$ holds at this order.

Expressions for the pseudoscalar (PS) meson mass and its decay constant arise from the spectral decomposition for infinite T ,

$$f_{\text{PP}}(x_0) = \sum_{i=1}^{\infty} c_i e^{-E_i x_0}, \quad E_1 = m_{\text{PS}}, \quad E_{i \geq 2}: \text{excited states contributions}, \quad (3.5)$$




Figure 1: Local PCAC quark masses for representative charm-strange (upper 2 panels) and charm-light (lower 2 panels) κ -combinations for the G8 and O7 ensembles. Our bare quark mass estimates are obtained as averages over the plateau regions.

which decays exponentially for large time separations. In this asymptotic regime, the decay constant is thus given by:

$$f_{\text{PS}} = Z_A (1 + \bar{b}_A am_{\text{sea}} + \tilde{b}_A am_{\text{rs}}) \times f_{\text{PS}}^{\text{bare}}, \quad f_{\text{PS}}^{\text{bare}} = 2\sqrt{2c_1} m_{\text{rs}} m_{\text{PS}}^{-3/2}. \quad (3.6)$$

Since in the actual analysis we face finite time extents T and separations x_0 , and hence particles running backwards in time and excited states, we employ the following two-step procedure to fix the region $x_0 \in [x_0^{\text{min}}, T - x_0^{\text{min}}]$, in which the excited state contribution to f_{PP} can be neglected: 1.) Determine x_0^{min} as the smallest x_0 , where the excited state, estimated by a 2-state fit (including finite- T effects) to $f_{\text{PP}}(x_0) = c_1 [e^{-E_1 x_0} + e^{-E_1(T-x_0)}] + c_2 [e^{-E_2 x_0} + e^{-E_2(T-x_0)}]$, contributes less than 1/4 of the statistical uncertainty on the effective mass $M_{\text{eff}}(x_0)$; here the effective PS meson mass M_{eff} is defined as $\cosh[M_{\text{eff}}(x_0 - T/2)] / \cosh[M_{\text{eff}}(x_0 + 1 - T/2)] = f_{\text{PP}}(x_0) / f_{\text{PP}}(x_0 + 1)$. 2.) Perform a 1-state fit of the asymptotic exponential decay, restricted to this region, to extract $m_{\text{PS}} = E_1$ and the leading coefficient c_1 , eventually entering the evaluation of the decay constants according to eq. (3.6). This is illustrated for our most chiral ensemble in figure 2.

4. Preliminary results

Unphysical pion masses and non-zero lattice spacings in our data are accounted for by employing joint chiral ($m_\pi \rightarrow m_{\pi, \text{phys}} = 134.8 \text{ MeV}$) and continuum limit ($a \rightarrow 0$) extrapolations.

Assuming a linear dependence on the squared (sea) pion mass, our fit ansatz for the renormalized PCAC quark mass composed of a charm and a light valence flavour ($s = s, l$) reads

$$m_{\text{R}}^{\text{cs}}(m_\pi, a) = B + Cm_\pi^2 + Da^2. \quad (4.1)$$

In addition, we also consider the definition via the bare subtracted quark mass, $m_{\text{q},c} = m_{0,c} - m_{\text{crit}}$, so that we have three ways to obtain the renormalization group invariant (RGI) charm quark mass,




Figure 2: The effective mass in the pseudoscalar channel extracted from f_{PP} for a representative charm-strange (left) and charm-light (right) κ -combination for the G8 ensemble. A two-state fit to data outside the shaded area, where the fit function describes the data well given their accuracy, determines x_0^{\min} . The result of the final one-state fit is given by the error band.

M_c [18, 19]:

$$\frac{1}{2}(M_c + M_s) = \frac{M}{\bar{m}} m_R^{cs}; \quad M_c = \frac{M}{\bar{m}} \frac{Z_A}{Z_P} Z \left(1 + b_m am_{q,c}\right) m_{q,c}, \quad m_{q,c} = \frac{1}{2} \left(\frac{1}{\kappa_c} - \frac{1}{\kappa_{\text{crit}}} \right). \quad (4.2)$$

The universal factor M/\bar{m} , which translates the running mass at a given scale to the RGI one, as well as the other renormalization and improvement factors entering here, are non-perturbatively known from [10, 14–16]. The combined m_π^2 - and a^2 -dependence of the three definitions in eq. (4.2) is shown in the left panel of figure 3 to lead to consistent results in the joint chiral and continuum limit. A more careful error analysis still to come, we consider the spread of these values as an upper limit for the overall uncertainty and quote as preliminary estimate for the charm quark’s mass

$$M_c = 1.51(4) \text{ GeV} \quad \Rightarrow \quad \bar{m}_c^{\overline{\text{MS}}}(\bar{m}_c^{\overline{\text{MS}}}) = 1.274(36) \text{ GeV}, \quad (4.3)$$

where M_s from [10] and in the conversion to the $\overline{\text{MS}}$ scheme the known 4-loop anomalous dimensions of quark masses and coupling [20, 21] together with $\Lambda_{\overline{\text{MS}}}$ from [10] were used.

For the D-meson decay constants, we adopt again fit ansätze linear in m_π^2 (and a^2), while for f_D we also model the sea quark dependence in a fit form inspired by partially quenched heavy meson chiral perturbation theory (HM χ PT) [22, 23], treating the charm quark as heavy, viz.

$$f_{D(s)}(m_\pi, a) = b_{(s)} + c_{(s)} m_\pi^2 + d_{(s)} a^2, \quad (4.4)$$

$$f_D(m_\pi, a) = b' \left[1 - \frac{3}{4} \frac{1 + 3\hat{g}^2}{(4\pi f_\pi)^2} m_\pi^2 \ln(m_\pi^2) \right] + c' m_\pi^2 + d' a^2; \quad (4.5)$$

m_s -terms are assumed to be absorbed into the $O(m_\pi^0)$ constants, because we work at fixed physical strange quark mass, and $\hat{g} = g_{D^*D\pi} = 0.6$ [24] is the $D^*D\pi$ -coupling. These combined chiral and




Figure 3: *Left:* Joint chiral and continuum extrapolation of the RGI mass of the charm quark, using the definitions via the heavy-light PCAC relation (where $s = s, 1$) and via the bare subtracted mass, eq. (4.2). *Right:* Joint chiral and continuum extrapolations to the physical point of f_D and f_{D_s} to the fit ansätze in eqs. (4.4) and (4.5). Labels “coarse” and “fine” refer to $\beta = 5.3$ and 5.5, respectively, and for $f_{D(s)}$ also fits neglecting cutoff effects are displayed.

continuum extrapolations to the physical point are depicted in the right panel of figure 3. As can be seen from the fits, our data for f_D are best described by the linear extrapolation along eq. (4.4), and we do not see any evidence for the significance of the chiral logarithm-term in eq. (4.5). Therefore, we take the linear extrapolations as the central values to arrive at our present results

$$f_{D_s} = 247(5)_{\text{stat}}(5)_{\text{syst}} \text{ MeV} , \quad f_D = 216(7)_{\text{stat}}(5)_{\text{syst}} \text{ MeV} , \quad f_{D_s}/f_D = 1.14(2)_{\text{stat}}(3)_{\text{syst}} \quad (4.6)$$

and the difference to the HM χ PT fit to account for a part of the systematic error of f_D . Apart from the statistical errors, the quoted uncertainties also contain a conservative estimate of the contribution from the scale setting.

5. Conclusions and outlook

The results for the charm quark mass and the D-meson decay constants from our analysis are very well in line with computations of other groups, see, e.g., the recent summaries in [25, 26]. Note that by setting the scale through f_K we effectively compute $f_{D(s)}/f_K$, where Z_A in eq. (3.6) (and thus also its error) drops out, but that it still re-enters indirectly by also fixing κ_s, κ_c through f_K [10]. This uncertainty, estimated conservatively so far, will likely decrease in the final analysis.

Acknowledgments. We thank Rainer Sommer for useful discussions and Nazario Tantalo for his contribution at an early stage of this project. This work is supported by the grant HE 4517/3-1 (J. H.) of the Deutsche Forschungsgemeinschaft. We are indebted to our colleagues in CLS for the joint production and use of the $N_f = 2$ gauge configurations. Most of our numerical simulations have been performed on the computers of the John von Neumann Institute for Computing at Forschungszentrum Jülich (under project ID “HCH09”), HLRN in Berlin and DESY, Zeuthen, and we thank these institutions for allocating computer time for this project and the computer center’s staff for their technical support. In particular, we also gratefully acknowledge the granted access to the HPC resources of the Gauss Center for Supercomputing at Forschungszentrum Jülich, Germany, made available within the Distributed European Computing Initiative by

the PRACE-2IP, receiving funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement RI-283493.

References

- [1] J. L. Rosner and S. Stone, published in J. Beringer et al. (Particle Data Group), *The review of particle physics*, *Phys. Rev.* **D86** (2012) 010001 [[arXiv:1201.2401](#)].
- [2] ALPHA, G. von Hippel, R. Sommer, J. Heitger, S. Schaefer and N. Tantalo, *PoS LATTICE2008* (2008) 227 [[arXiv:0810.0214](#)].
- [3] M. Lüscher, *Comput. Phys. Commun.* **156** (2004) 209 [[hep-lat/0310048](#)].
- [4] M. Lüscher, *Comput. Phys. Commun.* **165** (2005) 199 [[hep-lat/0409106](#)].
- [5] M. Lüscher, *J. High Energy Phys.* **0712** (2007) 011 [[arXiv:0710.5417](#)].
- [6] M. Lüscher, <http://luscher.web.cern.ch/luscher/DD-HMC/index.html>.
- [7] M. Hasenbusch, *Phys. Lett.* **B519** (2001) 177 [[hep-lat/0107019](#)].
- [8] ALPHA, M. Marinkovic and S. Schaefer, *PoS LATTICE2010* (2010) 031 [[arXiv:1011.0911](#)].
- [9] ALPHA, U. Wolff, *Comput. Phys. Commun.* **156** (2004) 143 [[hep-lat/0306017](#)].
- [10] ALPHA, P. Fritzsche, F. Knechtli, B. Leder, M. Marinkovic, S. Schaefer, R. Sommer and F. Virotta, *Nucl. Phys.* **B865** (2012) 397 [[arXiv:1205.5380](#)].
- [11] R. Sommer, *Nucl. Phys. Proc. Suppl.* **42** (1995) 186 [[hep-lat/9411024](#)].
- [12] UKQCD, M. Foster and C. Michael, *Phys. Rev.* **D59** (1999) 074503 [[hep-lat/9810021](#)].
- [13] ALPHA, M. Della Morte, R. Hoffmann and R. Sommer, *J. High Energy Phys.* **0503** (2005) 029 [[hep-lat/0503003](#)].
- [14] ALPHA, M. Della Morte, R. Hoffmann, F. Knechtli, J. Rolf, R. Sommer, I. Wetzorke and U. Wolff, *Nucl. Phys.* **B729** (2005) 117 [[hep-lat/0507035](#)].
- [15] ALPHA, M. Della Morte, R. Sommer and S. Takeda, *Phys. Lett.* **B672** (2009) 407 [[arXiv:0807.1120](#)].
- [16] ALPHA, P. Fritzsche, J. Heitger and N. Tantalo, *J. High Energy Phys.* **1008** (2010) 074 [[arXiv:1004.3978](#)].
- [17] ALPHA, S. Sint and P. Weisz, *Nucl. Phys.* **B502** (1997) 251 [[hep-lat/9704001](#)].
- [18] ALPHA, J. Rolf and S. Sint, *J. High Energy Phys.* **0212** (2002) 007 [[hep-ph/0209255](#)].
- [19] ALPHA, J. Heitger and A. Jüttner, *J. High Energy Phys.* **0905** (2009) 101 [[arXiv:0812.2200](#)]. Erratum: *ibid.* 1101 (2011) 036.
- [20] K. G. Chetyrkin and A. Retey, *Nucl. Phys.* **B583** (2000) 3 [[hep-ph/9910332](#)].
- [21] K. Melnikov and T. van Ritbergen, *Phys. Lett.* **B482** (2000) 99 [[hep-ph/9912391](#)].
- [22] J. L. Goity, *Phys. Rev.* **D46** (1992) 3929 [[hep-ph/9206230](#)].
- [23] S. R. Sharpe and Y. Zhang, *Phys. Rev.* **D53** (1996) 5125 [[hep-lat/9510037](#)].
- [24] CLEO, A. Anastassov *et al.*, *Phys. Rev.* **D65** (2002) 032003 [[hep-ex/0108043](#)].
- [25] FLAG Working Group, S. Aoki *et al.*, [arXiv:1310.8555](#).
- [26] A. X. El Khadra, *these proceedings*.