

The NESSiE Concept for Sterile Neutrinos

Luca Stanco*

I.N.F.N. Padova, Via Marzolo, 8 I-35131 Padova, Italy

E-mail: luca.stanco@pd.infn.it

Neutrino physics is nowadays receiving more and more attention as a possible source of information for the long-standing problem of new physics beyond the Standard Model. The recent measurement of the third mixing angle θ_{13} in the standard mixing oscillation scenario encourages us to pursue the still missing results on leptonic CP violation and absolute neutrino masses. However, several puzzling measurements exist, which deserve an exhaustive evaluation.

The NESSiE Collaboration has been setup to undertake a definitive experiment to clarify the muon disappearance measurements at small L/E , which will be able to put severe constraints to any model with more than the three-standard neutrinos, or even to robustly measure the presence of a new kind of neutrino oscillation for the first time. Within the context of the current CERN project, aimed to revitalize the neutrino field in Europe, we will illustrate the achievements that can be obtained by a double muon-spectrometer system, with emphasis on the search for sterile neutrinos.

*XV Workshop on Neutrino Telescopes,
11-15 March 2013
Venice, Italy*

*For the NESSiE Collaboration.

1. Introduction

The unfolding of the physics of the neutrino is a long and exciting story spanning the last 80 years. Over this time the interchange of theoretical hypotheses and experimental facts has been one of the most fruitful demonstrations of the progress of knowledge in physics. The work of the last decade and a half finally brought a coherent picture within the Standard Model (SM) (or some small extensions of it), namely the mixing of three neutrino flavour states with three ν_1 , ν_2 and ν_3 mass eigenstates. The last unknown mixing angle, θ_{13} , was recently measured [1] but still many questions remain unanswered to completely settle the scenario: the absolute masses, the Majorana/Dirac nature and the existence and magnitude of leptonic CP violation. Answers to these questions will beautifully complete the (standard) three–neutrino model but they will hardly provide an insight into new physics Beyond the Standard Model (BSM). Many relevant questions will stay open: the reason for neutrinos, the relation between the leptonic and hadronic sectors of the SM, the origin of Dark Matter and, overall, where and how to look for BSM physics. Neutrinos may be an excellent source of BSM physics and their story is supporting that at length.

There are actually several experimental hints for deviations from the “coherent” picture described above. Many unexpected results, not statistically significant on a single basis, appeared also in the last decade and a half, bringing attention to the hypothesis of *sterile neutrinos* [2]. A recent White Paper [3] contains a comprehensive review of these issues. Here we focus on one of the most intriguing and long–standing unresolved result: the unexpected oscillation of neutrinos at relatively small values of the ratio L/E (distance in km, energy in GeV), corresponding to a scale of $\mathcal{O}(1)$ eV^2 , incompatible with the much smaller values related to the atmospheric $|\Delta m_{32}^2| \simeq 2.4 \times 10^{-3}$ eV^2 and to the solar $\Delta m_{21}^2 \simeq 8 \times 10^{-5}$ eV^2 scales.

The first unexpected measurement came from an excess of $\bar{\nu}_e$ originating from an initial $\bar{\nu}_\mu$ beam from Decay At Rest (LSND [4]). The LSND experiment saw a 3.8σ effect. The subsequent experiment with ν_μ ($\bar{\nu}_\mu$) beam from accelerator, MiniBooNE [5], although confirming an independent 3.8σ effect after sustained experimental work, was unable to draw conclusive results on the origin of the LSND effect having observed an excess at higher L/E in an energy region where background is high.

In recent years many phenomenological studies were performed by analyzing the LSND effect together with similar unexpected results coming from the measurement of lower than expected rates of $\bar{\nu}_e$ and ν_e interactions (*disappearance*), either from **(a)** near-by nuclear reactors ($\bar{\nu}_e$) [6] or **(b)** from Mega-Curie K-capture calibration sources in the solar– ν_e Gallium experiments [7]. These ν_e ($\bar{\nu}_e$) disappearance measurements, all at the statistical level of 3–4 σ , could also be interpreted [8] as oscillations between neutrinos at large $\Delta m^2 \simeq 1$ eV^2 . Several attempts were then tried to reach a coherent picture in terms of mixing between active and sterile neutrinos, in $3 + 1$ and $3 + 2$ [9] or even $3 + 1 + 1$ [10] or $3 + 3$ [11] models, as extensions of the standard three–neutrino model. We refer to [12, 13, 14] as the most recent and industrious works where a very crucial issue is raised: “a consistent interpretation of the global data in terms of neutrino oscillations is challenged by the non-observation of a positive signal in ν_μ disappearance experiments” [13]. In fact, since the appearance probability of ν_e from ν_μ , in the two-flavour limit, is given by:

$$P(\nu_\mu \rightarrow \nu_e)_{SBL}^{3+1} = 4|U_{\mu 4}|^2|U_{e 4}|^2 \sin^2 \frac{\Delta m_{41}^2 L}{4E},$$

dependence on both the mixing elements with the sterile state, $|U_{e4}|$ and $|U_{\mu4}|$, rises up. Therefore, the presence of additional sterile states introduces quite naturally appearance and disappearance phenomena involving the flavour states in all production channels. In particular, a ν_μ disappearance effect has to be present and possibly measured. It turns out that only old experiments and measurements are available for Charged Current (CC) ν_μ interactions at small L/E [15]. The CDHS experiment reported in 1984 the non-observation of ν_μ oscillations in the Δm^2 range 0.3 eV^2 to 90 eV^2 . Their analyzed region of oscillation did not span however low values of the mixing parameter down to around 0.1 in $\sin^2(2\theta)$. More recent results are available on ν_μ disappearance from MiniBooNE [16], a joint MiniBooNE/SciBooNE analysis [17, 18] and the Long-Baseline MINOS experiment [19]. These results slightly extend the ν_μ disappearance exclusion region, however still leaving out the small-mixing region. Similar additional constraints on ν_μ disappearance could possibly come from the analysis of atmospheric neutrinos in IceCube [20].

Despite this set of measurements being rather unsatisfactory when compared with the corresponding LSND allowed region that lies at somewhat lower values of the mixing angle, they are still sufficient to introduce tensions in all the phenomenological models developed so far (see e.g. [3, 12, 13, 14, 21] for comprehensive and recent reviews). Therefore it is mandatory to setup a new experiment able to improve the small-mixing angle region exclusion by at least one order of magnitude with respect to the current results. In such a way one could also rule out the idea that the mixing angle extracted from LSND is larger than the true value due to a data over-fluctuation. Once again, the ν_μ disappearance channel should be the optimal one to perform a full disentangling of the mechanism given the strong tension between the ν_e appearance and ν_μ disappearance around $\Delta m^2 \simeq 1 \text{ eV}^2$. In fact, whereas the LSND effect may be confirmed by a more accurate ν_e oscillation measurement, only the presence of a ν_μ oscillation pattern could shed more light on the nature and the interpretation of the effect.

A very recent paper by us [22] depicted the phenomenology of the analysis that can be performed with a NESSiE-like experiment in the newly proposed Neutrino Platform at CERN, in case a neutrino-beam will be available. An extension of more than one order of magnitude in the sensitivity to the mixing parameter will be reached. This report, although corresponding to the presentation given at the Neutrino-Telescope Workshop in March 2013, is largely based on the more recent work of [22] and the up-to-date development of the NESSiE experiment. In particular, the re-use of available hardware from the OPERA experiment is reported.

2. The CERN experimental proposal

The need for a definitive clarification on the possible existence of a neutrino mass scale around 1 eV has brought up several proposals and experimental suggestions exploiting the sterile neutrino option by using different interaction channels and refurbished experiments. In the light of the considerations discussed in the previous section there are essentially two sets of experiments which must be redone: a) the measurement of $\bar{\nu}_e$ neutrino fluxes at reactors (primarily the ILL one) together with refined and detailed computations of the flux simulations [23]; b) the appearance/disappearance oscillation measurements at low L/E with a standard muon neutrino beam with its intrinsic electron neutrino component.

There is actually another interesting option which comes from the Neutrino Factory studies and the very recently submitted EOI from ν STORM [24]. We refer to [25] for a comprehensive review of the corresponding possible ν_e and ν_μ disappearance effects. It is interesting to note that the figures of merit about ν_μ disappearance as obtained in [22] are rather similar to or even slightly more competitive than those illustrated in [25] (e.g compare the exclusion regions in Fig. 6 of [25] and those in Fig. 5 of [22] despite the use of different C.L.), not forgetting the rather long time needed to setup the ν STORM project.

Coming to experiments with standard beams, investigations are underway at CERN where two Physics Proposals [26, 27] were submitted in October 2011 and later merged into a single Technical Proposal (ICARUS-NESSiE, [28]). CERN has subsequently set up working groups for the proton beam extraction from the SPS, the secondary beam line and the needed infrastructure/buildings for the detectors. The work was reported in a recent LOI and at this Conference [29].

The experiment is based on two identical Liquid Argon (LAr)–Time Projection Chambers (TPC) [27] complemented by magnetized spectrometers [26] detecting electron and muon neutrino events at far and near positions, 1600 m and 460 m away from the proton target, respectively. The project will exploit the ICARUS T600 detector, the largest LAr–TPC ever built of about 600 ton mass, now presently in the LNGS underground laboratory where it was exposed to the CNGS beam. It is supposed to be moved at the CERN “far” position. An additional 1/4 of the T600 detector (T150) would be constructed from scratch as a clone of the original one, except for the dimensions, and located in the near site. Two spectrometers would be placed downstream of the two LAr–TPC detectors to greatly enhance the physics reach. The spectrometers will exploit a bipolar magnet with instrumented iron slabs, and a newly designed air–core magnet, to perform charge identification and muon momentum measurements in an extended energy range (from 0.5 GeV or less to 10 GeV), over a transverse area larger than 50 m².

While the LAr–TPCs will mainly perform a direct measurement of electron neutrinos [31] the spectrometers will allow an extended exploitation of the muon neutrino component, with neutrino/antineutrino discrimination on an event-by-event basis [22].

2.1 The neutrino beam

The proposed new neutrino beam will be constructed in the SPS North Area [29]. The setup is based on a 100 GeV proton beam with a fast extraction scheme providing about $3.5 \cdot 10^{13}$ protons/pulse, in two pulses of 10.5 μ s durations¹ separated by 50 ms, that corresponds to about $4.5 \cdot 10^{19}$ protons on target (p.o.t.) per year. A target station will be located next to the so called TCC2 target zone, 11 m underground, followed by a cylindrical He-filled decay pipe with a length of about 110 m and a diameter of 3 m. The beam dump of 15 m in length, will be composed of iron blocks with a graphite inner core. Downstream of the beam dump a set of muon chambers stations will act as beam monitors. The beam will point upward, with a slope of about 5 mrad, resulting in a depth of 3 m for the detectors in the far site.

The current design of the focusing optics includes a pair of pulsed magnetic horns operated at relatively low currents. A graphite target of about 1 m in length is deeply inserted into the first horn

¹Pulses of 10.5 μ s duration are normally put in coincidence with the fast response of the spectrometers' detectors and efficiently used to reject the cosmic ray background. See e.g. [30] where a time resolution of less than 2 ns is reported for the detectors used in the OPERA experiment.

allowing a large acceptance for the focusing of low momentum pions emitted at large angles. This design allows production of a spectrum peaking at about 2 GeV thus matching the most interesting domain of Δm^2 with the detector locations at 460 and 1600 m from the target.

The charged current event rates for ν_μ and $\bar{\nu}_\mu$ at the near and far detectors are shown in Fig. 1 for the positive and negative focusing configuration.




Figure 1: Expected neutrino CC interactions in the no-oscillation hypothesis for positive polarity (left) and negative polarity (right) for the new proposed CERN neutrino beam (elaborated from [29]) and for an integrated luminosity of 1 year. In the first (second) row the rates are shown at the near (far) position. Note that the distributions of the antineutrino rates in case of positive polarity are multiplied by a factor 5 to allow a better visual inspection.

A relevant contamination of ν_μ in the negative polarity configuration is visible especially at high energy. This component arises as a result of the decays of high energy poorly de-focused mesons produced at small angles. The charge discrimination of the magnetic system described below will allow an efficient discrimination of these two components with a charge confusion below or of the order of 1% from sub-GeV (0.3–0.5 GeV) up to momenta around 8–10 GeV [26].

3. Spectrometer requirements

The main purpose of the spectrometers placed downstream of the LAr-TPC is to provide charge identification and momentum reconstruction for the muons produced in neutrino interactions occurring in the LAr volume or in the magnetized iron of the spectrometers. In order to perform this measurement with high precision and in a wide energy range, from sub-GeV to multi-GeV, a massive iron-core dipole magnet (ICM) is coupled to an air-core magnet (ACM) in front of it [26]. Low momentum muons will be measured by the ACM while the ICM will be employed at higher momenta.

As considered in the previous sections the definition of two sites, near and far, constitutes a fundamental issue for each physics program which aims to perform any sterile neutrino search. The two layouts have to be as similar as possible in order to allow an almost complete cancellation of the systematic uncertainties when comparing the measurements made at the near and far sites. Hence the near spectrometer will be an exact clone of the far one, with identical thickness along the beam axis but a scaled transverse size. A sketch of the possible far site NESSiE detector is shown in Fig. 2.




Figure 2: Sketch of the far site spectrometer that could be built by extensively reusing materials available from the OPERA spectrometers [32]. The basic elements of the new ACM concept are also depicted. Neutrinos are traveling from right to left.

The key feature of the ACM is the large geometric and momentum acceptance. The need of a low momentum threshold for muon reconstruction can be met using a magnet in air. The only dead material along the muon path is given by the conductors needed to generate the magnetic field and the position detectors instrumenting the magnet itself. For the conductors the use of aluminum instead of copper is preferable due to the lower Z and a density lower by a factor 3. The magnetic field needed in the low momentum range covered by the ACM is in the range 0.1 - 0.15 T. A spatial resolution in the range of 0.1 - 1 mm can be reached using drift tubes [33] as high precision trackers in combination with scintillator strip detectors [34]. These could provide the external trigger needed by the drift tubes and a coarse spatial measurement in the non-bending direction i.e. the direction parallel to the drift tubes. Silicon photomultiplier (SiPM) devices may eventually be used to read out some scintillator planes embedded in the magnetic field.

The general layout of the two OPERA [32] iron spectrometers fulfills the requirements of the ICM detector, and they could eventually be used for the CERN project putting the two magnets one after the other in order to obtain a total of 2.5 m longitudinal thickness of iron.

The OPERA spectrometers are built assembling vertical iron plates (slabs) in a planar structure of 875 cm (width) \times 800 cm (height). Each passive plane is made out of seven adjacent iron

slabs. Resistive Plate Chambers (RPC) [35] are sandwiched between iron planes; 21 RPC detectors, arranged in seven rows and three columns, are used in each active plane. Each RPC detector has a rectangular shape and covers an area of about 3.2 m^2 . The RPCs provide tracking measurements with about 1 cm resolution using the digital read-out of strips with a 2.6 cm pitch in the bending direction and 3.5 cm in the non-bending direction. The magnet is made of two arms with 22 RPC layers alternating with iron layers. Top and bottom iron yokes are connecting the arms. Copper coils surrounding the yokes are used to generate a magnetic field of about 1.5 T in the iron circuit. A total of 924 RPC chambers are needed to instrument each spectrometer. They could be recovered from the OPERA spectrometers and re-used. These RPC chambers are standard 2 mm gap chambers with bakelite electrodes with resistivity in the range from 10^{11} to $5 \times 10^{12} \Omega \cdot \text{cm}$ at $T = 20^\circ\text{C}$. In OPERA they are operated in the streamer regime with a gas mixture made of $\text{Ar}/\text{C}_2\text{H}_2\text{F}_4/\text{I-C}_4\text{H}_{10}/\text{SF}_6$ in the volume ratios of 75.4/20/4/0.6 at five refills/day. The high-amplitude streamer signals produced by charged tracks crossing the gas volume allow to house the Front-End (FE) discriminators in racks placed on top of the spectrometer. A totally new FE electronic for a surface and closeness to the neutrino production, i.e. at a very larger rate than that available at LNGS, has been already developed by the NESSiE Collaboration.




Figure 3: Sensitivity on the charge-ID and the momentum resolution for muon tracks coming from CC interactions. The full reconstruction procedure has been applied. On the left figure, the blue line corresponds to the mis-ID due only to the ICM part, while the red line is for the ACM part. The combination of the two magnetic system allow to achieve a mis-ID below 2% up to 5 GeV in the muon momentum. The right figure shows the relative resolution on the muon momentum. There are two regions: the first up to 3.5 GeV with a very good resolution of about 5% and the second region that corresponds to muons with momentum greater than 3.5 GeV. The first region corresponds to the momentum measured by range by the ICM. The second region is the resolution obtained by analyzing the curvature either in the ICM or the ACM (similar response by the two system is observed).

4. Perspectives

The magnetic detector system that has to be developed for the ν_μ disappearance measurement should take into account all the considerations depicted in previous sections. The system developed by the NESSiE Collaboration [26] is actually well suited since it couples a very powerful high- Z magnet for the momentum measurement via range to a low- Z magnet, to extend the useful muon momentum interval as low as possible, to allow charge discrimination on an event-by-event basis

and to allow NC event measurement whether coupled to an adequate (but not necessarily highly performant and with large mass) detector to identify NC events. The sensitivities that can be obtained are depicted in Fig. 3 for the charge mis-identification and the momentum resolution of the muon track. Full simulation and reconstruction of the neutrino-interaction events have been applied.

For one year of operation at the CERN beam, either with negative or positive polarity beam, Table 1 reports the expected interaction rates in the LAr-TPCs at the Near (fiducial 119 t) and Far locations (fiducial 476 t), and the expected rates of fully reconstructed events in the NESSiE spectrometers at the Near (fiducial 241 t) and Far locations (fiducial 661 t), with and without LAr contribution. Significant results will be achievable. Both ν_e and ν_μ disappearance modes will be used, in addition to ν_e appearance mode, to add conclusive information on the sterile mixing angles, either in the $3 + 1$ or the $3 + 2$ scenarios.

	NEAR (Neg. foc.)	NEAR (Pos. foc.)	FAR (Neg.)	FAR (Pos.)
$\nu_e + \bar{\nu}_e$ (LAr)	35 K	54 K	4.2 K	6.4 K
$\nu_\mu + \bar{\nu}_\mu$ (LAr)	2000 K	5200 K	270 K	670 K
App. Test Point	590	1900	360	910
ν_μ CC (NESSiE+LAr)	230 K	1200 K	21 K	110 K
ν_μ CC (NESSiE alone)	1150 K	3600 K	94 K	280 K
$\bar{\nu}_\mu$ CC (NESSiE+LAr)	370 K	56 K	33 K	6.9 K
$\bar{\nu}_\mu$ CC (NESSiE alone)	1100 K	300 K	89 K	22 K
Disapp. Test Point	1800	4700	1700	5000

Table 1: The expected rates of interaction (LAr) and reconstructed (NESSiE) events for 1 year of operation. Values for Δm^2 around 2 eV^2 are reported as example.




Figure 4: Disappearance signals as extracted from only one year of running in antineutrino mode. The intrinsic contribution of neutrinos in the running mode is sufficient to observe the possible disappearance signal in the neutrino channel, too. Only statistical errors are shown.

The powerfulness of the statistical sample that can be collected in just one year of running at the proposed CERN neutrino beam is demonstrated in Fig. 4. Again for one year of running in negative focussing the disappearance signals are reported. In the negative focussing running mode both

the contributions of neutrinos and anti-neutrinos are sufficiently high to allow for measurements at the same time.

Finally, we outline in Fig. 5 the estimation evaluated in the paper [22] (Fig. 7 of [22]). The estimated limits at 95% C.L. on ν_μ disappearance that can be achieved via the *Far/Near* estimator are shown for different data periods (3, 5 and 10 years, corresponding to $13.5 \cdot 10^{19}$, $22.5 \cdot 10^{19}$ and $45.0 \cdot 10^{19}$ p.o.t., respectively). The different results for ν_μ and $\bar{\nu}_\mu$ beams were evaluated using the two variables, p and $\log_{10}(1/p)$. In negative polarity runs the muon charge identification allows an independent, simultaneous and similar-sensitivity measurements of the $\bar{\nu}_\mu$ and ν_μ disappearance rates, due to the large ν_μ contamination in the $\bar{\nu}_\mu$ beam.




Figure 5: The estimated limits at 95% C.L. (taken from [22]) for ν_μ disappearance at a Short-Baseline beam at CERN for several luminosity running periods and different beam polarities, with a two-site massive spectrometer (770 tons and 330 tons, respectively) with 90% inner fiducial volume. The top figure refers to the positive polarity beam. The continuous (dashed) lines correspond to the sensitivity limits obtained with the $\log_{10}(1/p)$ (p) variable (used in [22]). 3 years correspond to $13.5 \cdot 10^{19}$ p.o.t., 5 years to $22.5 \cdot 10^{19}$ p.o.t. and 10 years to $45.0 \cdot 10^{19}$ p.o.t. The exclusion limit from combined MiniBooNE and SciBooNE ν_μ disappearance result at 90% C.L. from Ref. [17] is shown for comparison by the black curve in the right. The bottom figure refers to the negative polarity beam. Sensitivity limits are evaluated with the $\log_{10}(1/p)$ variable. Clearly the negative polarity run allows the contemporary analysis of the $\bar{\nu}_\mu$ and ν_μ disappearance exclusion regions thanks to the disentangling of the muon charge on an event-by-event-basis. The black curve in the right shows for comparison the central value of the sensitivity at 90% C.L. from combined MiniBooNE and SciBooNE $\bar{\nu}_\mu$ disappearance result (Ref. [18]).

5. Conclusions

Neutrino physics is receiving more and more attention as a venue for the long standing search for new physics beyond the Standard Model. The current anomalies which do not fit into the established standard scenario with 3 neutrinos deserve refined studies and experiments. The CERN proposal for a new Short-Baseline experimental project is a very valuable one. We illustrated the current critical tensions in the muon-neutrino disappearance field and the achievements that can be obtained within the CERN project and the experiment proposed by the NESSiE Collaboration. Specifically, by considering $\mathcal{O}(1)$ kton massive spectrometers, an improvement by an order of magnitude can be obtained in the sensitivity to the mixing parameter space between standard neutrinos

and sterile ones with respect to today's limits. Conversely, a possible ν_μ disappearance signal will be essential to measure the relevant physical parameters and to fully disentangle the different sterile models. An effective analysis can be performed with a two-site experiment by using muon spectrometers with a low-Z part that allows clean charge identification on an event-by-event basis, and with a massive part allowing clean momentum measurement through range. Such a kind of spectrometer is under study by the NESSiE Collaboration and might be available with a limited investment using the OPERA spectrometers. The performances of these kinds of detectors are suitable to put a definitive result on the sterile neutrino issue at the eV mass scale.

Acknowledgements

We warmly thank the Organizers of the Neutrino Telescopes Workshop for the kind invitation and the relevance they endowed to our project. We would also like to thank the CERN and INFN managements for their strong support towards the Nessie Collaborations in view of the development of the foreseen CERN Neutrino-Platform.

References

- [1] DAYA-BAY Collaboration, F. An et al., "*Observation of electron-antineutrino disappearance at Daya Bay*", Phys. Rev.Lett. 108, 171803 (2012), arXiv:1203.1669.
RENO Collaboration, J. Ahn et al., "*Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment*", Phys. Rev.Lett. 108, 191802 (2012), arXiv:1204.0626.
DOUBLE-CHOOZ Collaboration, Y. Abe et al., "*Indication of Reactor $\bar{\nu}_e$ Disappearance in the Double Chooz Experiment*", Phys. Rev. Lett. 108, 131801 (2012), arXiv:1207.6632.
T2K Collaboration, K. Abe et al., "*Indication of Electron Neutrino Appearance from an Accelerator-produced Off-axis Muon Neutrino Beam*", Phys. Rev.Lett. 107, 041801 (2011), arXiv:1106.2822.
- [2] B. Pontecorvo, "*Neutrino Experiments and the Problem of the Conservation of Leptonic Charge*", Zh. Eksp. Teor. Fiz. 53, 1717 (1967) [Sov. Phys. JETP 26, 984 (1968)].
- [3] K.N. Abazajian et al., "*Light Sterile neutrinos: a White Paper*", (2012), arXiv:1204.5379.
- [4] LSND Collaboration, A. Aguilar et al., "*Evidence for neutrino oscillations from the observation of $\bar{\nu}_e$ appearance in a $\bar{\nu}_\mu$ beam*", Phys. Rev. D 64, 112007 (2001), hep-ex/0104049.
- [5] MiniBooNE Collaboration, A. A. Aguilar-Arevalo et al., "*A Search for electron neutrino appearance at the $\Delta m^2 \sim 1 \text{ eV}^2$ scale*", Phys.Rev.Lett. 98, 231801 (2007), arXiv:0704.1500;
MiniBooNE Collaboration, A. A. Aguilar-Arevalo et al., "*Event Excess in the MiniBooNE Search for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ Oscillations*", Phys. Rev. Lett. 105, 181801 (2010), arXiv:1007.1150;
MiniBooNE Collaboration, A. A. Aguilar-Arevalo et al., "*Improved statistics in the search for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations in MiniBooNE*", Nucl. Phys. B, Proc. Suppl. 235-236, 207 (2013);
C. Polly (MiniBooNE Collaboration), "*Results from MiniBooNE*", Talk at Neutrino2012, 3-9 June 2012, Kyoto, Japan, 2012, Nucl. Phys. B 235-236, vol. 1, 207, (2013).;
MiniBooNE Collaboration, A. A. Aguilar-Arevalo et al., "*Improved Search for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ Oscillations in the MiniBooNE Experiment*", Phys. Rev. Lett. 110, 161801 (2013), arXiv:1303.2588.
- [6] G. Mention, et al., Phys.Rev. D83 (2011) 073006, arXiv:1101.2755, and references therein.
D. Lhuillier, "*Reactor flux calculations*", Nucl. Phys. B, Proc. Suppl. 235-236, 11 (2013);
T. Lasserre, "*Testing the Reactor and Gallium Anomalies with Intense (Anti)Neutrino Emitters*", Nucl. Phys. Proc. Suppl. 235-236, 214 (2013), arXiv:1209.5090.

- [7] J. N. Abdurashitov et al. (SAGE Collaboration), Phys. Rev. C 80, 015807 (2009); J. N. Abdurashitov et al. (SAGE Collaboration), Phys. Rev. Lett. 77, 4708 (1996); J. N. Abdurashitov et al. (SAGE Collaboration), Phys. Rev. C 59, 2246 (1999); J. N. Abdurashitov et al., Phys. Rev. C 73, 045805 (2006); F. Kaether, W. Hampel, G. Heusser, J. Kiko, and T. Kirsten, Phys. Lett. B 685, 47 (2010); P. Anselmann et al. (GALLEX Collaboration), Phys. Lett. B 342, 440 (1995); W. Hampel et al. (GALLEX Collaboration), Phys. Lett. B 420, 114 (1998).
- [8] C. Giunti and M. Laveder, “*Statistical Significance of the Gallium Anomaly*”, Phys. Rev. C 83, 065504 (2011), arXiv:1006.3244.
- [9] Some relevant papers are the following: J. Kopp, M. Maltoni, and T. Schwetz, “*Are there Sterile-Neutrinos at the eV scale?*”, Phys. Rev. Lett. 107, 091801 (2011), arXiv:1103.4570; Giunti, C. and Laveder, M., “*3+1 and 3+2 Sterile Neutrino Fits*”, Phys. Rev. D 84, 073008 (2011), arXiv:1107.1452.
- [10] E. Kuflik, S. D. McDermott and K. M. Zurek, “*Neutrino Phenomenology in a 3+1+1 Framework*”, Phys. Rev. D 86, 033015 (2012), arXiv:1205.1791; A. E. Nelson, “*Effects of CP Violation from Neutral Heavy Fermions on Neutrino Oscillations, and the LSND/MiniBooNE Anomalies*”, Phys. Rev. D 84, 053001 (2011), arXiv:1010.3970.
- [11] J. M. Conrad, C. M. Ignarra, G. Karagiorgi, M. H. Shaevitz and J. Spitz, “*Sterile Neutrino Fits to Short-Baseline Neutrino Oscillation Measurements*”, Adv. High Energy Phys. 2013, 163897 (2013), arXiv:1207.4765.
- [12] J. Kopp, P. A.N. Machado, M. Maltoni, T. Schwetz, “*Sterile Neutrino Oscillations: The Global Picture*”, JHEP 05, 050 (2013), arXiv:1303.3011.
- [13] T. Schwetz, “*Status of sterile neutrino oscillations*”, Nucl. Phys. B, Proc. Suppl. 235-236, 229 (2013).
- [14] C. Giunti, M. Laveder, Y. F. Li and H. W. Long, “*Update of short-baseline electron neutrino and antineutrino disappearance*”, Phys. Rev. D 86, no. 11, 113014 (2012), arXiv:1308.5288.
- [15] CDHS Collaboration, F. Dydak et al., “*A Search for ν_μ Oscillations in the Δm^2 range $0.3 \div 90 \text{ eV}^2$* ”, Phys. Lett. B 134, 281 (1984).
- [16] MiniBooNE Collaboration, A. A. Aguilar-Arevalo et al., “*A search for muon neutrino and antineutrino disappearance in MiniBooNE*”, Phys. Rev. Lett. 103, 061802 (2009), arXiv:0903.2465.
- [17] MiniBooNE and SciBooNE Collaborations, K. B. M. Mahn et al., “*Dual baseline search for muon neutrino disappearance at $0.5 \text{ eV}^2 < \Delta m^2 < 40 \text{ eV}^2$* ”, Phys. Rev. D 85, 032007 (2012), arXiv:1106.5685.
- [18] MiniBooNE and SciBooNE Collaborations, G. Cheng et al., “*Dual baseline search for muon antineutrino disappearance at $0.1 \text{ eV}^2 < \Delta m^2 < 100 \text{ eV}^2$* ”, Phys. Rev. D 86, 052009 (2012), arXiv:1208.0322.
- [19] MINOS Collaboration, P. Adamson et al., “*Search for sterile neutrino mixing in the MINOS long-baseline experiment*”, Phys. Rev. D 81, 052004 (2010), arXiv:1001.0336.
- [20] S. Razzaque, A. Y. Smirnov, “*Searches for sterile neutrinos with IceCube DeepCore*”, Phys. Rev. D 85, 093010 (2012), arXiv:1203.5406; A. Esmaili, F. Halzen, O. Peres, “*Constraining Sterile Neutrinos with AMANDA and IceCube Atmospheric Neutrino Data*”, Jour. Cosm. and Astr. Phys. 11, 041 (2012), arXiv:1206.6903.
- [21] A. Palazzo, “*Phenomenology of light sterile neutrinos: a brief review*”, Modern Phys. Lett. A 28, no. 7 (2013), arXiv:1302.1102.

- [22] L. Stanco et al., “An Appraisal of Muon Neutrino Disappearance at Short Baseline”, Adv. High Energy Phys. 2013, 948626 (2013), arXiv:1306.3455v2.
- [23] P. Huber, “On the determination of anti-neutrino spectra from nuclear reactors”, Phys. Rev. C84 024617 (2011), arXiv: 1106.0687;
P. Huber, “Reactor neutrinos fluxes”, Talk at NeuTel2012, 11-15 March 2013, Venice, Italy, 2013.
- [24] ν STORM Collaboration, D. Adey et al., “Neutrinos from Stored Muons (ν STORM): Expression of Interest”, arXiv:1305.1419.
- [25] W. Winter, “Optimization of a Very Low Energy Neutrino Factory for the Disappearance Into Sterile Neutrinos”, Phys. Rev. D85, 113005 (2012), arXiv:1204.2671.
- [26] NESSiE Collaboration, P. Bernardini et al., “Prospect for Charge Current Neutrino Interactions Measurements at the CERN-PS”, SPSC-P-343 (2011), arXiv:1111.2242.
- [27] C. Rubbia et al., “A comprehensive search for “anomalies” from neutrino and anti-neutrino oscillations at large mass differences ($\Delta m^2 \sim 1 \text{ eV}^2$) with two LAr-TPC imaging detectors at different distances from the CERN-PS”, SPSC-P-345 (2011).
- [28] ICARUS and NESSiE Collaborations, M. Antonello et al., “Search for anomalies from neutrino and anti-neutrino oscillations at $\Delta m^2 \sim 1 \text{ eV}^2$ with muon spectrometers and large LAr-TPC imaging detectors”, SPSC-P-347 (2012), arXiv:1203.3432.
- [29] M. Nessi et al., “Letter of Intent for the new CERN Neutrino Facility (CENF)”, <https://edms.cern.ch/nav/P:CERN-0000077383:V0/P:CERN-0000096728:V0/TAB3>.
M. Nessi, “The New CERN Neutrino Facility”, Talk at NeuTel2012, 11-15 March 2013, Venice, Italy, 2013.
- [30] E. Balsamo et al., “The OPERA RPCs front end electronics; a novel application of LVDS line receiver as low cost discriminator”, JINST 7, P11007 (2012).
- [31] C. Rubbia, A. Guglielmi, F. Pietropaolo, P. Sala, “Sterile neutrinos: the necessity for a 5 sigma definitive clarification”, arXiv:1304.2047.
- [32] R. Acquafredda et al., “The OPERA experiment in the CERN to Gran Sasso neutrino beam”, JINST 4, P04018 (2009).
- [33] R. Zimmermann et al., “The precision tracker of the OPERA detector”, Nucl. Instrum. Meth. A555, 435 (2005), [Erratum-ibid. A557, 690 (2006)].
- [34] T. Adam et al., “The OPERA experiment target tracker”, Nucl. Instrum. Meth. A 577, 523 (2007).
- [35] A. Bergnoli et al. “Tests of OPERA RPC Detectors”, IEEE Trans. Nucl. Sci. 52, 2963 (2005).
M. Ambrosio et al. “The OPERA muon spectrometer tracking electronics”, Nucl. Instrum. Meth. A 533, 173 (2004).