

## Four-loop on-shell integrals: $\overline{\text{MS}}$ -on-shell relation and g-2

---

**P. Marquard\***

*Deutsches Elektronen Synchrotron DESY, Platanenallee 6, D15738 Zeuthen, Germany*

*E-mail: Peter.marquard@desy.de*

We present first results towards a full four-loop calculation for both the anomalous magnetic moment of the muon and the  $\overline{\text{MS}}$ -on-shell relation. The calculation requires the detailed study of an up to now not considered class of diagrams, so-called on-shell diagrams, at four-loop order.

*11th International Symposium on Radiative Corrections (Applications of Quantum Field Theory to Phenomenology)*

*22-27 September 2013*

*Lumley Castle Hotel, Durham, UK*

---

\*Speaker.

## 1. Introduction

The anomalous magnetic moments of electron and muon have been measured with unrivaled precision. In the case of the muon the experimental value<sup>1</sup>

$$a_{\mu}^{exp} = 1.16592080(54)(33)[63] \cdot 10^{-3} \quad (1.1)$$

has to be compared with the theory prediction

$$a_{\mu}^{theo} = 1.16591790(65) \cdot 10^{-3}. \quad (1.2)$$

The discrepancy between the two values of about 3 standard deviations is approximately of the same order as the four-loop QED corrections. The four- and five-loop corrections have been calculated in Ref.[1] and have not been verified by an independent calculation. In this paper we will present first steps to such an independent calculation to verify the results in Ref.[1]. An object that is technically related is the  $\overline{\text{MS}}$ -on-shell relation for quark masses in QCD. One of the main motivations, why it is of importance to know the  $\overline{\text{MS}}$ -on-shell relation with four-loop accuracy, is the planned determination of the top-quark mass at a future linear collider. The precision reached at such an experiment requires an equally precise knowledge of the  $\overline{\text{MS}}$ -on-shell relation.

In the following we will review recent results for both the anomalous magnetic moment and the  $\overline{\text{MS}}$ -on-shell relation.

## 2. Calculation and Results

For both, the calculation of the  $\overline{\text{MS}}$ -on-shell relation and the anomalous magnetic moment of the muon, the evaluation of on-shell integrals is necessary. To be more precise, to obtain the  $\overline{\text{MS}}$ -on-shell relation the quark propagator has to be evaluated on its mass shell, while for the anomalous magnetic moment the magnetic form factor of the muon-photon vertex has to be calculated for vanishing photon momentum, thus also leading to an on-shell diagram. Typical diagrams appearing in the calculation are shown in Fig 1 and 2. On-shell integrals have not been studied at four loops in any detail. Thus one of the main obstacle for the calculation lies in the calculation of the missing master integrals.


The calculation is set up as follows, the Feynman diagrams are generated using QGRAF [2], its output is then converted into FORM [3] input using q2e and exp [4, 5]. Suitable projectors are applied and the resulting scalar integrals are reduced to master integrals using integration-by-parts identities implemented in CRUSHER [6] and FIRE [7]. The reduction leads to the master integrals shown in Figs. 3 and 4. The integrals in Fig. 3 can be expressed in closed form in terms of Gamma functions while the ones in Fig. 4 have been calculated in an expansion in  $\epsilon = (d - 4)/2$  using the method of dimensional recurrence and analyticity [8]. As an example we show the result for master

---


<sup>1</sup>The errors indicated in brackets denote the statistical and systematic ones. The error given in square brackets is obtained by adding them in quadrature.

integral  $M_3$  (cf. Fig. 4)


$$\begin{aligned}
e^{4\epsilon\gamma_E} M_3 = & -\frac{1}{6\epsilon^4} - \frac{7}{6\epsilon^3} - \left(\frac{10}{3} + \frac{13\pi^2}{18}\right)\epsilon^{-2} - \left(-\frac{61}{6} + \frac{73\pi^2}{18} + \frac{118\zeta_3}{9}\right)\epsilon^{-1} - \left(-\frac{851}{4} + \frac{83\pi^2}{18} \right. \\
& + \left.\frac{637\zeta_3}{9} + \frac{37\pi^4}{10}\right)\epsilon^0 - \left(-\frac{14861}{8} - \frac{3467\pi^2}{36} + \frac{1003\zeta_3}{18} + \frac{1121\pi^4}{60} + \frac{1894\pi^2\zeta_3}{27} \right. \\
& + \left.\frac{16018\zeta_5}{15}\right)\epsilon^1 - \left(-\frac{613975}{48} - \frac{25981\pi^2}{24} - \frac{68293\zeta_3}{36} + \frac{83\pi^4}{24} + \frac{9559\pi^2\zeta_3}{27} + \frac{79891\zeta_5}{15} \right. \\
& + \left.\frac{59501\pi^6}{2835} + \frac{17704\zeta_3^2}{27}\right)\epsilon^2 - \left(-\frac{7539347}{96} - \frac{382349\pi^2}{48} - \frac{482627\zeta_3}{24} - \frac{426659\pi^4}{720} \right. \\
& + \frac{3757\pi^2\zeta_3}{54} + \frac{2525\zeta_5}{6} + \frac{43201\pi^6}{420} + \frac{88585\zeta_3^2}{27} + \frac{17204\pi^4\zeta_3}{45} + \frac{206434\pi^2\zeta_5}{45} \\
& \left. + \frac{1267243\zeta_7}{21}\right)\epsilon^3 + O(\epsilon^4). \tag{2.1}
\end{aligned}$$




**Figure 1:** Feynman diagrams for the calculation of the  $\overline{\text{MS}}$ -on-shell relation.



**Figure 2:** Feynman diagrams for the calculation of the anomalous magnetic moment.



**Figure 3:** Simple master integrals. They can be expressed in closed form in terms of Gamma functions.



**Figure 4:** Non-trivial master integrals. They can be calculated in an expansion in  $\varepsilon = (4-d)/2$ .

## 2.1 $\overline{\text{MS}}$ -on-shell relation

The  $\overline{\text{MS}}$ -on-shell relation can be written as a power series in the strong coupling constant  $\alpha_s$

$$\begin{aligned}
 z_m^{\text{OS}}(\mu) &= \frac{\bar{m}_q(\mu)}{M_q} = \frac{Z_m^{\text{OS}}}{Z_m^{\overline{\text{MS}}}} \\
 &= 1 + \frac{\alpha_s(\mu)}{\pi} \delta z_m^{(1)} + \left( \frac{\alpha_s(\mu)}{\pi} \right)^2 \delta z_m^{(2)} + \left( \frac{\alpha_s(\mu)}{\pi} \right)^3 \delta z_m^{(3)} + \left( \frac{\alpha_s(\mu)}{\pi} \right)^4 \delta z_m^{(4)} \\
 &\quad + \mathcal{O}(\alpha_s^5)
 \end{aligned} \tag{2.2}$$

and labeling contributions from massless and massive quark loops by  $n_l$  and  $n_h$ , respectively, we obtain the result for contributions from diagrams with at least two massless quark loops

$$\begin{aligned}
 z_m^{\text{OS}} &= 1 - A_s 1.333 + A_s^2 (-14.229 - 0.104 n_h + 1.041 n_l) \\
 &\quad + A_s^3 (-197.816 - 0.827 n_h - 0.064 n_h^2 + 26.946 n_l - 0.022 n_h n_l - 0.653 n_l^2) \\
 &\quad + A_s^4 (-43.465 n_l^2 - 0.017 n_h n_l^2 + 0.678 n_l^3 + \dots) + \mathcal{O}(A_s^5),
 \end{aligned} \tag{2.3}$$

with  $A_s = \alpha_s(m_q)/\pi$ .

## 2.2 Anomalous magnetic moment of the muon

In the approximation of a massless electron only the leading term including the logarithms can be obtained. For the sub-leading contributions a proper asymptotic expansion has to be performed. Expanding  $a_\mu$  in a power series in the fine structure constant  $\alpha$

$$a_\mu = 1 + \frac{\alpha}{\pi} a_\mu^{(1)} + \left(\frac{\alpha}{\pi}\right)^2 a_\mu^{(2)} + \left(\frac{\alpha}{\pi}\right)^3 a_\mu^{(3)} + \left(\frac{\alpha}{\pi}\right)^4 a_\mu^{(4)} + \mathcal{O}(\alpha^5) \quad (2.4)$$

and marking contributions from electron loops by  $n_l$

$$a_\mu^{(4)} = n_l^3 a_\mu^{(43)} + n_l^2 a_\mu^{(42)} + \dots \quad (2.5)$$

we obtain the result for contributions with three electron loops

$$a_\mu^{(43)} = \frac{1}{54} L_{\mu e}^3 - \frac{25}{108} L_{\mu e}^2 + \left(\frac{317}{324} + \frac{\pi^2}{27}\right) L_{\mu e} - \frac{2\zeta_3}{9} - \frac{25\pi^2}{162} - \frac{8609}{5832} \approx 7.19666, \quad (2.6)$$

where  $L_{\mu e} = \ln(M_\mu^2/M_e^2)$ . The result for diagrams with two electron loops can be further split into a contribution with and without an additional muon loop,  $a_\mu^{(42)b}$  and  $a_\mu^{(42)a}$ , respectively,

$$a_\mu^{(42)} = a_\mu^{(42)a} + a_\mu^{(42)b},$$

with

$$\begin{aligned} a_\mu^{(42)a} &= L_{\mu e}^2 \left[ \pi^2 \left( \frac{5}{36} - \frac{a_1}{6} \right) + \frac{\zeta_3}{4} - \frac{13}{24} \right] + L_{\mu e} \left[ -\frac{a_1^4}{9} + \pi^2 \left( -\frac{2a_1^2}{9} + \frac{5a_1}{3} - \frac{79}{54} \right) \right. \\ &\quad \left. - \frac{8a_4}{3} - 3\zeta_3 + \frac{11\pi^4}{216} + \frac{23}{6} \right] - \frac{2a_1^5}{45} + \frac{5a_1^4}{9} + \pi^2 \left( -\frac{4a_1^3}{27} + \frac{10a_1^2}{9} \right. \\ &\quad \left. - \frac{235a_1}{54} - \frac{\zeta_3}{8} + \frac{595}{162} \right) + \pi^4 \left( -\frac{31a_1}{540} - \frac{403}{3240} \right) + \frac{40a_4}{3} + \frac{16a_5}{3} - \frac{37\zeta_5}{6} \\ &\quad + \frac{11167\zeta_3}{1152} - \frac{6833}{864} \\ &\approx -3.62427, \end{aligned} \quad (2.7)$$

$$\begin{aligned} a_\mu^{(42)b} &= \left( \frac{119}{108} - \frac{\pi^2}{9} \right) L_{\mu e}^2 + \left( \frac{\pi^2}{27} - \frac{61}{162} \right) L_{\mu e} - \frac{4\pi^4}{45} + \frac{13\pi^2}{27} + \frac{7627}{1944} \\ &\approx 0.49405. \end{aligned} \quad (2.8)$$

Our results for  $a_\mu^{(43)}$  and  $a_\mu^{(42)b}$  agree with the results given in Refs. [12, 9]. The result for  $a_\mu^{(42)a}$  can be compared with the result from Refs. [1, 10]

$$a_\mu = -3.64204(112). \quad (2.9)$$

Our new result confirms the previously obtained results, the small discrepancy is due to missing terms in the expansion in  $m_e/m_\mu$ .

To finish, let us also present the result of a recent calculation of the contribution from  $\tau$ -leptons. Using asymptotic expansion in Ref. [11] the four-loop corrections due to  $\tau$ -leptons was obtained. The new result

$$A_{2,\mu}^{(8)}(M_\mu/M_\tau) = 0.0421670 + 0.0003257 + 0.0000015 = 0.0424941(2)(53) \quad (2.10)$$

is more precise and in full agreement with previous evaluations [1].

### 3. Conclusions

We presented results for both the  $\overline{\text{MS}}$ -on-shell relation and the anomalous magnetic moment of the muon at four-loop order. These results comprise a first step towards the full four-loop calculation and confirm the results known in the literature.

### Acknowledgements

We like to thank A. Kurz, R. Lee, T. Liu, A.V. Smirnov, V.A. Smirnov and M. Steinhauser for collaboration on the topics presented in this proceedings article. P.M. has been supported in part by DFG Sonderforschungsbereich Transregio 9, Computergestützte Theoretische Teilchenphysik, and by the EU Network LHCPHENOnet PITN-GA-2010-264564.

### References

- [1] Tatsumi Aoyama, Masashi Hayakawa, Toichiro Kinoshita, and Makiko Nio. Complete tenth-order QED contribution to the muon  $g-2$ . *Phys.Rev.Lett.*, 109:111808, 2012.
- [2] P. Nogueira. Automatic feynman graph generation. *J. Comput. Phys.*, 105:279–289, 1993.
- [3] J.A.M. Vermaseren. New features of FORM. math-ph/0010025.
- [4] T. Seidensticker. Automatic application of successive asymptotic expansions of Feynman diagrams. hep-ph/9905298.
- [5] R. Harlander, T. Seidensticker, and M. Steinhauser. Complete corrections of  $\mathcal{O}(\alpha\alpha_s)$  to the decay of the Z boson into bottom quarks. *Phys.Lett.*, B426:125–132, 1998.
- [6] P. Marquard and D. Seidel. Crusher. unpublished.
- [7] A.V. Smirnov. Algorithm FIRE – Feynman Integral REduction. *JHEP*, 0810:107, 2008.
- [8] R.N. Lee. Space-time dimensionality D as complex variable: Calculating loop integrals using dimensional recurrence relation and analytical properties with respect to D. *Nucl.Phys.*, B830:474–492, 2010.
- [9] Jean-Philippe Aguilar, David Greynat, and Eduardo De Rafael. Muon Anomaly from Lepton Vacuum Polarization and The Mellin-Barnes Representation. *Phys.Rev.*, D77:093010, 2008.
- [10] Toichiro Kinoshita and M. Nio. Improved  $\alpha^4$  term of the muon anomalous magnetic moment. *Phys.Rev.*, D70:113001, 2004.
- [11] Alexander Kurz, Tao Liu, Peter Marquard, and Matthias Steinhauser. Anomalous magnetic moment with heavy virtual leptons. arXiv:1311.2471.
- [12] S. Laporta. The analytical contribution of some eighth order graphs containing vacuum polarization insertions to the muon ( $g-2$ ) in QED. *Phys.Lett.*, B312:495–500, 1993.