

Analytic Continuation of Real Loop Quantum Gravity : Lessons from black hole thermodynamics

Jibril Ben Achour*

Laboratoire APC – Astroparticule et Cosmologie, Université Paris Diderot Paris 7, 75013 Paris, France

benachou@apc.univ-paris7.fr

Karim Noui

Laboratoire de Mathématiques et Physique Théorique, Université François Rabelais, Parc de Grandmont, 37200 Tours, France

Laboratoire APC – Astroparticule et Cosmologie, Université Paris Diderot Paris 7, 75013 Paris, France

karim.noui@lmpt.univ-tours.fr

This contribution is devoted to summarize the recent results obtained in the construction of an “analytic continuation” of Loop Quantum Gravity (LQG). By this we mean that we construct analytic continuation of physical quantities in LQG from real values of the Barbero-Immirzi parameter γ to the purely imaginary $\gamma = \pm i$. This should allow us to define a quantization of gravity with self dual Ashtekar variables. We first realized in [1] that this procedure, when applied to compute the entropy of a spherical black hole in LQG for $\gamma = \pm i$, allows to reproduce exactly the Bekenstein Hawking area law at the semi classical limit. The rigorous construction of the analytic continuation of spherical black hole entropy has been done in [2]. Here we start with a review of the main steps of this construction: we recall that our prescription turns out to be unique (under natural assumptions) and leads to the right semi-classical limit with its logarithmic quantum corrections. Furthermore, the discrete and γ -dependent area spectrum of the black hole horizon becomes continuous and obviously γ -independent. Then, we review how this analytic continuation could be interpreted in terms of the analytic continuation from the compact gauge group $SU(2)$ to the non compact gauge group $SU(1, 1)$ relying on an analysis of three dimensional quantum gravity.

Frontiers of Fundamental Physics 14 - FFP14,

15-18 July 2014

Aix Marseille University (AMU) Saint-Charles Campus, Marseille

*Speaker.

1. Motivation : Getting rid of γ

The Barbero-Immirzi parameter γ seems to play apparently a paradoxical role in LQG. Whereas it is totally irrelevant in the classical theory, it enters into the expressions of “physical” quantities like eigenvalues of geometric operators in the kinematical sector, the maximal density of the universe in quantum cosmology or the black hole entropy. Nonetheless, many observations (from black holes physics [1, 2] and three dimensional quantum gravity [16, 17]) indicate that γ should somehow “disappear” from the quantum theory in the sense that it should take the natural complex value $\gamma = \pm i$ and not a real value. We review here some of these observations focussing mainly on the case of the black hole entropy. We finish with a discussion where we quickly review the state of the art in three dimensions.

2. Analytic continuation of black hole entropy

2.1 Real γ Black Holes

In the framework of LQG, a black hole is defined as a boundary in space-time which satisfies the constraints of an isolated horizon [3]. Those constraints impose that the black hole degrees of freedom are encoded into the phase space of an $SU(2)$ Chern Simons theory defined on a punctured two-sphere S^2 as a canonical surface. The presence of the gauge group $SU(2)$ derives from the phase space of gravity in the bulk expressed in terms of Ashtekar-Barbero variables. The Chern-Simons level k is proportional to the horizon area a_H and depends on γ according to (2.1)¹. The punctures originate from the spin networks (the quantum states of the gravitational field) defined in the bulk which pierce the horizon. They are viewed as the fundamental excitations of the black hole and each puncture carries a quantum of area a_l which contributes to the macroscopic area a_H in the usual “real γ ” picture according to (n labels the number of punctures)

$$a_H = \frac{2\pi\gamma}{(1-\gamma^2)}k = \sum_{l=1}^n a_l \quad a_l = 8\pi l_P^2 \gamma \sqrt{j_l(j_l+1)}. \quad (2.1)$$

As usual in LQG, the spin $j_l \in \mathbb{N}/2$ labels an $SU(2)$ unitary irreducible representation (irrep). For a fixed n , a microscopic state of the black hole is defined by an ordered² family of spins $\mathcal{P} = (j_1, \dots, j_n)$.

The degeneracy of a configuration \mathcal{P} is given by the dimension of the Chern-Simons Hilbert space $\mathcal{H}_k(S^2; j_1, \dots, j_n)$ which is well-known to be defined by the space of $U_q(su(2))$ invariant tensors³ in the tensor product $\otimes_l V_l$. Here V_l are $U_q(su(2))$ modules labelled by the spins j_l whose dimension is denoted $d_l = 2j_l + 1$. The quantum parameter is a root of unity defined by $q = \exp(i\pi/(k+2))$ where the level k is necessarily integer. The dimension $g_k(d_l)$ of this Hilbert space

¹Variant expressions exist but the precise dependence on γ is not important for our purpose. The main point is that k is large when the area a_H is macroscopic.

²In the usual real picture, the punctures are distinguishable. For this reason, we consider a priori an ordered family of spins. In the complex picture, we relax the distinguishability.

³The space of invariant tensors is endowed with the quantum Haar measure when viewed as the space of linear forms on $SU_q(2)$, the polynomials of the quantum deformation of $SU(2)$.

is easily computed [4] and can be expressed as the following sum over the integer d

$$g_k(d_l) = \frac{2}{2+k} \sum_{d=1}^{k+1} \sin^2\left(\frac{\pi d}{k+2}\right) \prod_{l=1}^n \frac{\sin\left(\frac{\pi}{k+2} d d_l\right)}{\sin\left(\frac{\pi}{k+2} d\right)}. \quad (2.2)$$

This Verlinde formula allows to recover at the semi-classical limit (a_H large in Planck units) the Bekenstein-Hawking law for the black hole entropy provided that γ is fixed to a peculiar value [5, 6]. This is easily seen in the simplest model where $d_l = d$ for any l is fixed and n becomes large at the semi-classical limit. Even if this result is certainly an important success of LQG, it has risen important questions concerning the role of γ and the validity of the computation we have just sketched. Since then, different interpretations of γ have been discussed but none are totally convincing (see [7] and references therein).

2.2 Complex $\gamma = \pm i$ Black Holes

The last couple of years, a new road towards the understanding of the role of γ has emerged. In that new picture, the Barbero-Immirzi parameter is viewed as a “regulator” which should be sent back to its “original” imaginary value $\gamma = \pm i$. To clarify this point of view, let us recall that γ has been first introduced to overcome the problem of working with complex variables, to circumvent the resolution of the reality conditions and then to start the loop quantization of gravity. It is important to notice that such a strategy has been successful because it has led to a very beautiful picture of the quantum (kinematical) geometry at the Planck scale. However, for solving the quantum dynamics, the real Ashtekar-Barbero connection doesn’t seem to be well suited anymore. Already at the classical level, it is well known that this connection doesn’t transform properly under timelike diffeomorphisms, and this might be the reason why γ remains in the theory at least at the kinematical level. This fact is enhanced by a series of recent works which all point towards the need to come back to the (anti) self-dual variables [1, 8, 9, 10]. One of the most striking result [1, 2] in that respect has been obtained in the context of black hole physics: the analytic continuation of the formula (2.2) for the Chern-Simons Hilbert space dimension to the imaginary value $\gamma = \pm i$ allows to reproduce the expected semi-classical Bekenstein-Hawking law for the black hole entropy. We are going to briefly recall how this works following the construction of [2]. Details can be found in the original paper [2].

First of all, we immediately notice that taking $\gamma = \pm i$ leads to a complex value of the Chern-Simons level which becomes $k = i\lambda$ with $\lambda \in \mathbb{R}$. From the LQG point of view, this is an immediate consequence of (2.1). From the Chern-Simons theory point of view, this shift from $k \in \mathbb{N}$ to $k \in i\mathbb{R}$ can be interpreted by the fact that one works now with a complex $SL(2, \mathbb{C})$ connection rather than a compact real $SU(2)$ connection [11]. Unfortunately, Chern-Simons theory with complex gauge group and complex level is poorly understood at the quantum level and the only one serious candidates for its quantization deeply relies on analytic continuation techniques [11]. However, the process of analytic continuation is rather subtle even in the construction of the analytic continuation of the Hilbert space dimension (2.2). Indeed, since k enters in the upper bound of the sum, the expression (2.2) is not really convenient for analytic continuation purposes even if we used it formally in the first proposal [1]. It is much more convenient to view (2.2) as a sum of residues of an analytic function in order to write it as an integral in the complex plane along a contour \mathcal{C} (see figure 1) which encompass the imaginary axis between $[0, i\pi]$:




Figure 1: The contour $\mathcal{C} = \mathcal{C}_+ + \mathcal{C}_-$ in the complex plane encloses the imaginary axis between $[0, i\pi]$.

$$g_k(d_l) = \frac{i}{\pi} \oint_{\mathcal{C}} dz \sinh^2(z) \prod_{l=1}^n \frac{\sinh(d_l z)}{\sinh(z)} \coth((k+2)z). \quad (2.3)$$

To simplify the discussion below, we introduce the notation $G(z) = \coth((k+2)z)$ while the remaining part of the integrand will be denoted $F(z)$. When k and d_l are integers, the poles of the integrand are the poles of $G(z)$ which are located on the imaginary axis: $z_p = \frac{i\pi p}{k+2}$ for $p \in \mathbb{N}^*$. This justifies the choice of the contour \mathcal{C} .

Now, it makes sense to consider $k = i\lambda$ as G is an analytic function of k . Let us discuss what happens to the integral (2.3) when one performs such an analytic continuation.

- If the dimensions d_l remain integer, the poles of the integrand are located on the real axis: the poles of $G(z)$ are $z_p = -\frac{\pi p}{\lambda}$ with $p \in \mathbb{N}^*$ whereas $F(z)$ still has no pole. In this case, a contour \mathcal{C} close enough to the complex axis doesn't enclose any poles and the integral vanishes. As a consequence, analytic continuing only the Chern Simons level from $k \in \mathbb{N}^*$ to $k \in i\mathbb{R}$ keeping d_l unchanged leads to inconsistent physical results.
- If $d_l = is_l \in i\mathbb{R}$, the analytic continuation is much more interesting. The location of the poles of $G(z)$ is unchanged compared to the previous case but the novelty is that $F(z)$ admits new poles on the imaginary axis at $z_m = i\pi m$ with $m \in \mathbb{N}^*$. Among all these new poles, we are more interested in the one located at $i\pi$ which must be enclosed by the contour \mathcal{C} in order for the integral (2.3) to be non-trivial. Therefore, such a continuation leads to a non-trivial result for the black hole Hilbert space that we define to be the dimension⁴ of the black hole Hilbert space when $\gamma = \pm i$.

⁴To be interpreted as a dimension, g_k must necessary be a non-negative real number. This is asymptotically the case, i.e. when the horizon area becomes large in Planck units and under some conditions satisfied by the number n of punctures. For non large area, g_k is in general complex but we can argue that we have to consider $|g_k|$ or $\Re(g_k)$ as the dimension even if this aspect deserves to be studied deeper. In [2], we considered the modulus of g_k .

To summarize, the dimension of the black hole Hilbert space for $\gamma = \pm i$ is defined from the analytic continuation of (2.3) when \mathcal{C} encloses the point $i\pi$ and

$$k = i\lambda \quad , \quad d_l = is_l \Leftrightarrow j_l = \frac{1}{2}(-1 + is_l) \quad \lambda, s_l \in \mathbb{R}^+. \quad (2.4)$$

The fact that λ and s_l are non negative is not restrictive. At this point, it is important to explain the choice $d_l \in i\mathbb{R}$. From a physical point of view, this is the only consistent choice which leaves the area spectrum real when $\gamma = \pm i$ as seen from

$$A(j_l) = 8\pi l_p^2 \gamma \sqrt{j_l(j_l + 1)} \xrightarrow{\gamma = \pm i} A(s_l) = 4\pi l_p^2 \sqrt{s_l^2 + 1} \quad (2.5)$$

where we choose the square root of -1 such that the area is non negative. From a mathematical point of view, changing j_l to $\frac{1}{2}(-1 + is_l)$ amounts to considering $SU(1, 1)$ irreps instead of $SU(2)$ irreps for coloring the punctures.

2.3 Semi-classical limit: area law and logarithmic corrections

To simplify the study of the semi-classical limit, we consider the model where all the punctures carry the same color $s_l = s$. This corresponds to the one color model in the following. We first impose that k is large in (2.3) and we obtain the following expression for the candidate to the dimension of the one-color black hole Hilbert space

$$g_\infty(s, n) = \frac{i}{\pi} \oint_{\mathcal{C}} dz \sinh^2(z) e^{nS(z)} \quad S(z) = \log\left(\frac{\sinh(sz)}{\sinh(z)}\right). \quad (2.6)$$

In the semi-classical limit, the black hole area $a_H = 4\pi l_p^2 ns$ is large, which means that the product ns becomes large. It has been argued in [12] that the semi-classical regime corresponds to both n and s large. Then, the form (2.6) of the integral is well suited for the study of the thermodynamical limit. When n is large, this integral can be estimated using the stationary phase method. The study of the critical points reveals that there are two critical points, $z_c = 0$ and $z_c = i(\pi + \frac{1}{s}) + o(\frac{1}{s})$ for s large. Only the later contributes to the saddle point approximation which finally leads to

$$S_m = \log(g_\infty(s, n)) = \frac{a_H}{4l_p^2} + S_{cor} \quad (2.7)$$

for the black hole microcanonical entropy S_m . The leading term reproduces the expected Bekenstein-Hawking area law without any fine tuning and S_{cor} are quantum corrections. At this point, the quantum corrections scale in general as $\sqrt{a_H}$ in Planck units, and then are much larger than logarithmic corrections.

In the grand canonical ensemble, the situation concerning the quantum corrections is more satisfying. Using the local framework developed in [13, 14], we have a notion of energy (measured by an observer located at a "small" distance L of the horizon compared to the black hole radius) which allows to compute the canonical and grand canonical partition functions. In this approach, the black hole is viewed as a gas of indistinguishable punctures which has been studied first in [12]. If we assume in addition that the punctures admit a non-vanishing chemical potential μ and satisfy the Maxwell-Boltzmann statistics, we can show that, at the semi-classical limit, the black

hole temperature approaches the Unruh temperature $T_U = 1/\beta_U$ for the local observer and the black hole mean area \bar{a}_H , the mean number of punctures \bar{n} and the mean color \bar{s} scale as follows

$$\bar{a}_H = 4\pi l_p^2 \frac{z}{2x^2} \left(1 - \frac{3}{2z}x\right) \quad x = \frac{2L}{l_p^2}(\beta - \beta_U) \quad \bar{n} \propto \sqrt{\bar{a}_H}, \quad \bar{s} \propto \sqrt{\bar{a}_H} \quad (2.8)$$

where $z = \exp(\beta\mu)$ is the fugacity. As a consistency check, we recover that n and s are large in the semi-classical regime. It is easy to compute from this analysis the semi-classical expansion of the grand canonical entropy and we deduce the expression

$$S_{gc} = \frac{\bar{a}_H}{4l_p^2} - \frac{3}{2} \log \left(\frac{\bar{a}_H}{l_p^2} \right) + \frac{z_U}{2} (2 - \mu\beta_U) \left(\frac{\bar{a}_H}{l_p^2} \right)^{1/2} + o \left(\log \left(\frac{\bar{a}_H}{l_p^2} \right) \right). \quad (2.9)$$

We recover therefore the expected logarithmic quantum corrections supplemented with larger quantum corrections $\propto \sqrt{\bar{a}_H}$ which vanish when the chemical potential is fixed to $\mu = 2T_U$. A physical interpretation of such a result is still missing. Note however that the same value of the chemical potential is also found to cancel the too large quantum correction for the real black hole ($\gamma \in \mathbb{R}$) with the Maxwell Boltzman statistic [15]. Therefore, this behaviour of the quantum correction $S_{cor} \propto \sqrt{\bar{a}_H}$ is not specific to the complex model.

The calculation can be generalized in different ways. First, we can extend the model to a black hole with p colors [2]. In that case, the entropy has the same form as (2.9) with a modified logarithmic correction which depends on p . Interestingly we notice that only the case $p = 1$, i.e. the one color model, which can be interpreted as a kind of spherical symmetric quantum condition, allows to recover the prefactor $-3/2$ for the logarithmic corrections. We could also generalize to the cases where the punctures satisfy a quantum statistic. This has been done in [12, 15] when the area spectrum is discrete. In that context, we showed that assuming the punctures are boson, there exists a semi-classical regime where the gas condensates to spin $1/2$ punctures and also where the quantum corrections are logarithmic⁵. It would be interesting to see whether a similar phenomenon occurs when the spectrum is continuous.

Finally, this prescription was applied to the simplest Loop Quantum Cosmology model in [18]. It was shown that the analytic continuation described above preserves also the bouncing universe scenario. Another study was realized in the context of three dimensional gravity, in order to understand the appearance of the gauge group $SU(1, 1)$. For more details on those results, see [16, 17].

This analytic continuation constitutes a proposal for defining a theory of self-dual quantum gravity in terms of the complex Ashtekar connection and for solving the so-called reality conditions. We expect that the systematic investigation of this analytic continuation in various setups will eventually lead to new insights on the status of the quantum states of complex Ashtekar gravity. The first attempt to define a Wick rotation in the context of Ashtekar gravity was proposed in [19]. Establishing a clear link between the two approaches would inevitably shed some light on the one described in this contribution.

⁵This contrasts with the case of a classical Maxwell-Boltzmann statistics where large spins dominate at the semi-classical limit.

References

- [1] E. Frodden, M. Geiller, K. Noui and A. Perez, Black holes entropy with complex Ashtekar variables, *Europhys.Lett* **107**, 10005 (2012), arXiv:1212.4060 [gr-qc].
- [2] J. Ben Achour, K. Noui and A. Mouchet, Analytic continuation of black hole entropy in loop quantum gravity, (2014), arXiv:1406.6021 [gr-qc].
- [3] A. Ashtekar, A. Corichi and K. Krasnov, Isolated horizons : the classical phase space, *Adv.Theor.Math.Phys* 3 (1999) 419-478 arXiv:9905.089 [gr-qc].
- [4] J. Engle, K. Noui and A.Perez, D. Pranzetti, The SU(2) Black Hole entropy revisited *JHEP*. 1105 (2011) 016 arXiv:1103.2723 [gr-qc].
- [5] C. Rovelli, Black hole entropy and Loop Quantum Gravity, *Phys.Rev.Lett.* 77 (1996) 3288-3291 arXiv:9603.063 [gr-qc].
- [6] A. Ashtekar, J. Baez, A. Corichi, K. Krasnov Quantum geometry and Black hole entropy, *Phys.Rev.Lett.* 80 (1998) 904-907 arXiv:9710.007 [gr-qc].
- [7] M.Geiller and K.Noui, A note on the Holst action, the time gauge, and the Barbero-Immirzi parameter, *Gen.Rel.Grav.* 45 1733-1760 (2013) arXiv:1212.5064 [gr-qc].
- [8] D. Pranzetti and H. Salmahn, Horizon entropy with loop quantum gravity methods (2014), arXiv:1412.7435 [gr-qc].
- [9] D. Pranzetti, Geometric temperature and entropy of isolated horizon *Phys.Rev.* D89 (2014) 104046, arXiv:1309.6714 [gr-qc].
- [10] M.Geiller and K.Noui, Near-Horizon Radiation and Self-Dual Loop Quantum Gravity *Euro.Phys.Lett.* 105 (2014) 60001 arXiv:1402.4138 [gr-qc].
- [11] E. Witten Analytical continuation of Chern Simons theory Conference proceedings. (2010) C09-08-03.10 arXiv:1001.2933 [gr-qc].
- [12] A. Ghosh, K. Noui and A. Perez, Statistics, holography, and black hole entropy in loop quantum gravity, *Phys. Rev. D* **89**, 084069 (2014), arXiv:1309.4563 [gr-qc].
- [13] A. Gosh and A. Perez, Black hole entropy and isolated horizons thermodynamics *Phys. Rev. D* **87**, 121503 (2013), arXiv:1110.4055 [gr-qc].
- [14] E. Frodden, A. Ghosh and A. Perez, Quasilocal first law for black hole thermodynamics, *Phys. Rev. D* **87**, 121503 (2013), arXiv:1110.4055 [gr-qc].
- [15] O. Azin, J. Ben Achour, M. Geiller, K. Noui and A. Perez, Black hole as a gas of punctures with a chemical potential : Bose Einstein condensation and logarithmic corrections to the entropy, (2014) arXiv:1412.5851 [gr-qc], .
- [16] J. Ben Achour, M. Geiller, K. Noui and C. Yu, Testing the role of the Barbero Immirzi parameter and the choice of connection in Loop Quantum Gravity, (2013) arXiv:1306.3241 [gr-qc].
- [17] J. Ben Achour, M. Geiller, K. Noui and C. Yu, Spectra of geometric operators in three dimensional LQG : from discrete to continuous Published in *Phys.Rev. D89* (2014) 064064, arXiv:1306.3246 [gr-qc].
- [18] J. Ben Achour, J. Grain, and K. Noui, Loop Quantum Cosmology with Complex Ashtekar Variables, *Class. Quantum Gravity.* (2014), arXiv:1407.3768 [gr-qc].
- [19] T. Thiemann, Reality conditions inducing transforms for quantum gauge field theory and quantum gravity Published in *Class.Quant.Grav.* 13 (1996) 1383-1404, gr-qc/9511057.