

Neutrino scattering physics with the SHiP Experiment

Antonia DI CRESCENZO*

†

Universita e INFN, Napoli (IT)

E-mail: antonia.di.crescenzo@cern.ch

SHiP (Search for Hidden Particles) is a new general purpose fixed target facility, proposed at the CERN SPS accelerator. In its initial phase the 400 GeV protons beam will be dumped on a heavy target with the aim of integrating 2×10^{20} pot in five years. A dedicated detector downstream the target will allow to probe a variety of models with the light long-lived exotic particles and masses below $O(10)$ GeV/ c^2 . Another dedicated detector will allow the study of active neutrino cross-sections and angular distributions. In particular, the neutrino deep-inelastic cross-sections will be performed with a statistics 1000 times larger than currently available, with the extraction of the F_4 and F_5 structure functions, never measured so far. Tau neutrinos will be distinguished by anti-neutrinos, thus providing the first observation of the tau anti-neutrino. With muon neutrinos it will be possible to study the strangeness content of the nucleon.

*XXIII International Workshop on Deep-Inelastic Scattering
27 April - May 1 2015
Dallas, Texas*

*Speaker.

†on behalf of the SHiP Collaboration.

1. Introduction

The Standard Model of elementary particle physics has provided a consistent description of Nature's fundamental constituents and their interactions. Its predictions have been tested and confirmed by numerous experiments. With the discovery of the Higgs boson [1, 2], all the predicted constituents of the SM have now been observed. At the same time, no significant deviations from the Standard Model were found in direct or in indirect searches for new physics.

For the particular value of the Higgs mass it is possible that the Standard Model remains mathematically consistent and valid as an effective field theory up to a very high energy scale, possibly all the way to the scale of quantum gravity, the Planck scale [3, 4, 5].

However, it is clear that the SM is not a complete theory. It fails to explain a number of observed phenomena in particle physics, astrophysics and cosmology. The major unsolved challenges are: the non-zero neutrino mass, the existence of dark matter, and the baryon asymmetry of the universe. Some yet unknown particles or interactions would be needed to explain these puzzles and to answer these questions. The hypothetical new particles can be searched for either at the energy and at the intensity frontier.

The high energy frontier will be comprehensively investigated in the next few decades, while the searches for alternative low mass *beyond Standard Model* physics at the intensity frontier have been neglected in recent years. Light new particles may have remained undetected by previous experiments because of the very small couplings involved. New data at the intensity frontier will therefore be particularly useful in exploring portal models with light new physics, and in searching for Majorana neutrinos. A new intensity frontier experiment, SHiP, is consequently very timely for direct searches for very weakly interacting new physics.

2. The SHiP experiment

The main physics goal of the SHiP experiment consists in exploring hidden portals and extensions of the Standard Model (SM) which incorporate long-lived and very weakly interacting particles through the direct detection of their decays to SM particles. For this purpose, the 400 GeV protons beam of the SPS will be dumped on a heavy target. Over five years of data taking, 2×10^{20} protons on target are expected.

Hidden particles are predominantly produced in decays of hadrons, in particular in decays of charmed and beauty hadrons above the kaon mass. They have very small coupling with SM particles and are therefore very long-lived. In order to maximise their production, the target is made of molybdenum and tungsten, being these materials characterized by a very short interaction length. The target is followed by an iron absorber, aiming at stopping the hadrons and the electromagnetic radiation emerging from the target.

In order to reduce the large flux of muons produced in the decay of pions and kaons, a 48 m long active muon shield based on magnetic deflection is located immediately downstream of the target. At SPS energies, the D_s meson decays are the principal source of tau neutrinos. The charm hadrons decays are also a source of electron and muon neutrinos. The experimental setup is therefore ideally suited for studying interactions of tau neutrinos and neutrino induced charm production by neutrinos and anti-neutrinos of all flavours. The facility will therefore host a 10 m long tau neu-

trino detector, located upstream of the decay volume of the hidden particles (HS) detector. The HS detector is made by a cylindrical 50 m long decay volume. It is under vacuum and surrounded by a veto system. The full reconstruction of the hidden particle decays is performed by a magnetic spectrometer and a system for particle identification at the end of the decay volume. Figure 1 shows a schematic overview of the SHiP facility from the proton target to the end of the Hidden Sector detector.




Figure 1: Overview of the SHiP facility.

3. The neutrino detector

The tau neutrino detector shown in Figure 2 is located immediately downstream of the active muon shield. It consists of the Neutrino Target in a magnetic field followed by a Muon Magnetic Spectrometer. The Neutrino Target is made of Opera-type modules which employ the Emulsion Cloud Chamber (ECC) technology. The charges of the hadrons produced in the neutrino interactions are identified using the Target Tracker (TT) consisting of active planes which interleave the ECC modules. The magnetic field is provided by the Goliath magnet. The Muon Magnetic Spectrometer of the tau neutrino detector consists of a warm iron dipole magnet instrumented with active layers based on the Opera Resistive Plate Chambers and the Drift Tube Tracker. The Emulsion Cloud Chamber (ECC) structure used in the Neutrino Detector is made of a sequence of passive material plates interleaved with emulsion films. The target is modular and its units are made of two parts as shown in Figure ??: the brick and the Compact Emulsion Spectrometer (CES). The brick, using lead as passive material, combines the micrometric tracking accuracy of nuclear emulsions and the high lead density as required to maximise the number of neutrino interactions in a compact detector. The CES is made of a sandwich of light material plates (e.g. Rohacell) and emulsion films. It is designed to distinguish ν_τ and $\bar{\nu}_\tau$ by performing the electric charge measurement of the τ decay products.

So far only DONUT [6] and OPERA [7] experiments succeeded in detecting a few ν_τ interactions. The ECC structure is well suited for the measurement of charged particles momenta [8] and

for electron identification [9], as largely exploited with the OPERA experiment. The momentum is measured by the multiple coulomb scattering in the lead plates; electrons can be identified by detecting the electromagnetic showers. The ν_τ and $\bar{\nu}_\tau$ interactions are identified through the detection of the τ lepton production and decay. The τ decay channels investigated are the electron ($BR = 17.8\%$), muon ($BR = 17.7\%$), single-hadron ($BR = 49.5\%$) and three-hadron ($BR = 15.0\%$) channels.




Figure 2: Overview of the tau neutrino detector.

4. Tau neutrino physics

The neutrino detector has the unique capability of detecting all three neutrino flavors and of distinguishing neutrinos from anti-neutrinos. The expected number of charged-current deep-inelastic neutrino interactions in a 9.6 ton detector in 5 years run is reported in Table 1: the energy spectra are shown in Figure 3.

	$\langle E \rangle$ (GeV)	Interactions
N_{ν_e}	46	2.5×10^5
N_{ν_μ}	29	1.7×10^6
N_{ν_τ}	59	6.7×10^3
$N_{\bar{\nu}_e}$	46	9.0×10^4
$N_{\bar{\nu}_\mu}$	28	6.7×10^5
$N_{\bar{\nu}_\tau}$	58	3.4×10^3

Table 1: Charged-current deep-inelastic neutrino interactions integrated in the target in five years run.




Figure 3: Energy spectrum of the different neutrino flavors interacting in the neutrino detector. A 0.5 GeV cut is applied for ν_μ and ν_e . The total number of neutrinos is normalized to 100.

The charged-current ν_τ ($\bar{\nu}_\tau$) differential cross-section is represented by a standard set of five structure functions:

$$\frac{d^2\sigma^{\nu(\bar{\nu})}}{dxdy} = \frac{G_F^2 M E_\nu}{\pi(1+Q^2/M_W^2)^2} \left((y^2x + \frac{m_\tau^2 y}{2E_\nu M}) F_1 + \left[(1 - \frac{m_\tau^2}{4E_\nu^2}) - (1 + \frac{Mx}{2E_\nu}) \right] F_2 \right. \\ \left. \pm \left[xy(1 - \frac{y}{2}) - \frac{m_\tau^2 y}{4E_\nu M} \right] F_3 + \frac{m_\tau^2(m_\tau^2 + Q^2)}{4E_\nu^2 M^2 x} F_4 - \frac{m_\tau^2}{E_\nu M} F_5 \right),$$

where $\{x, y, Q^2\}$ are the standard DIS kinematic variables related through $Q^2 = 2M_N E_\nu xy$.

The structure functions F_4 and F_5 , pointed out by Albright and Jarlskog in Ref. [10], are neglected in muon neutrino interactions because of a suppression factor depending on the square of the charged lepton mass divided by the nucleon mass times neutrino energy. Given the higher mass value of the τ lepton, F_4 and F_5 structure functions contribute, instead, to the tau neutrino cross section. At leading order, in the limit of massless quarks and target hadrons, $F_4 = 0$ and $2xF_5 = F_2$, where x is the Bjorken- x variable (Albright-Jarlskog relations). Calculations at NLO show that F_4 is about 1% of F_5 [11].

With the statistics of tau neutrino interactions collected in five years run, the SHiP experiment will have the unique capability of being sensitive to F_4 and F_5 . The hypothesis of $F_4 = F_5 = 0$ would result in an increase of the ν_τ and $\bar{\nu}_\tau$ charged-current deep-inelastic cross sections and consequently, of the number of expected ν_τ and $\bar{\nu}_\tau$ interactions.

The difference between the cross sections in the $F_4 = F_5 = 0$ hypothesis and the SM one is larger for lower neutrino energies. This behavior reflects in the energy dependence of the variable r , defined as the ratio between the cross section in the two hypotheses: it is higher for lower neutrino energies, where the discrepancy of the two curves is larger, and decreases, tending to one, for higher energies, where the contribution of F_4 and F_5 becomes negligible.

The ratio r is reported for $\bar{\nu}_\tau$ in the left plot of Fig. 4. To have evidence of a non-zero value of

F_4 and F_5 , the ratio r is required to be larger than 3σ , being σ the uncertainty on the incoming neutrino flux, amounting to 20%. This condition is satisfied for $E_{\bar{\nu}_\tau} < 38$ GeV, where we expect to observe about 300 $\bar{\nu}_\tau$ interactions.

The ratio r was estimated also for the sum of ν_τ and $\bar{\nu}_\tau$. The right plot of Fig. 4 shows that in this case $r > 3\sigma$ for neutrino energies below 20 GeV, where the number of observed $\nu_\tau + \bar{\nu}_\tau$ interactions, not requiring the leptonic number to be measured, is expected to be about 420.




Figure 4: Energy dependence of the ratio r between the DIS cross section in the $F_4 = F_5 = 0$ hypothesis and the SM prediction, for $\bar{\nu}_\tau$ (left) and for the sum of ν_τ and $\bar{\nu}_\tau$ (right).

5. Charmed hadron physics

The SHiP experiment is suitable also to perform studies of charmed hadron production. The relative charm production yield in muon and electron neutrino and anti-neutrino interactions expected at SHiP energy amounts to 4% and 6%, respectively. In five years run, more than 10^5 neutrino induced charmed hadrons are expected, thus largely exceeding the statistics available in previous experiments by more than one order of magnitude. Therefore all the studies on charm physics performed with neutrino interactions will be revised with improved accuracy and some channels inaccessible in the past will be explored.

Unlike neutrino scattering where the presence of valence quarks favours the d -quark as neutrino target thus compensating the large suppression provided by the Cabibbo angle, charmed hadron production in anti-neutrino interactions selects anti-strange quark in the nucleon. Precise knowledge of the strangeness is an important information for many precision tests of the SM as well as for BSM searches at the LHC.

The potential impact of simulated charm data was assessed by adding them to the NNPDF3.0 NNLO fit [12]. The constraining power of the SHiP pseudodata is shown in Figure 5 (left) for the s^+ variable defined as $s^+ = s(x) + \bar{s}(x)$. The vertical axis reports $1 + \Delta s^+ / s^+$ where Δs^+ is the accuracy on s^+ , in such a way that the difference to unity indicates the accuracy. The horizontal axis is the Bjorken variable x . A significant improvement on this variable is achieved in the x range between 0.03 and 0.35. Figure 5 (right) shows the improvement achieved on the variable $s^- = s(x) - \bar{s}(x)$. versus the x variable. The vertical axis reports $1 + \Delta s^- / s^-$. A significant gain with SHiP data is obtained in the x range between 0.08 and 0.30.




Figure 5: The blue line shows the present accuracy in the distribution of $s^+ = s + \bar{s}$ (left) $s^- = s - \bar{s}$ (right) as a function of x . The red contour shows the improvement obtained with SHiP.

6. Conclusions

The SHiP experiment is a new general-purpose fixed target facility at the CERN SPS to search for hidden particles as predicted by a very large of recently elaborated models of Hidden Sectors which can explain dark matter, neutrino oscillations and the baryon asymmetry in the Universe. The SHiP experiment is also suited to perform active neutrino physics with high statistics. It will have the opportunity observe for the first time the tau anti-neutrino and to perform tau neutrino and tau anti-neutrino cross-section measurements, with the extraction of F_4 and F_5 , never measured so far. The large flux of muon and electron neutrino will allow the study of the charmed hadron production with high statistics and the estimation of the strangeness content of the nucleon.

References

- [1] G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Phys.Lett. **B716** (2012) 1.
- [2] S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Phys.Lett. **B716** (2012) 30.
- [3] D. Buttazzo, G. Degrassi, P. P. Giardino, G. F. Giudice, F. Sala, et al., *Investigating the near-criticality of the Higgs boson*, JHEP **1312** (2013) 089.
- [4] G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, et al., *Higgs mass and vacuum stability in the Standard Model at NNLO*, JHEP **1208** (2012) 098.
- [5] F. Bezrukov, M. Y. Kalmykov, B. A. Kniehl, and M. Shaposhnikov, *Higgs Boson Mass and New Physics*, JHEP **1210** (2012) 140.
- [6] K. Kodama et al., *Observation of tau neutrino interactions*, Phys.Lett. **B504** (2001) 218.
- [7] N. Agafonova et al., *Observation of tau neutrino appearance in the CNGS beam with the OPERA experiment*, PTEP **2014(10)** (2014) 101C01.
- [8] N. Agafonova et al., *Momentum measurement by the Multiple Coulomb Scattering method in the OPERA lead emulsion target*, New J.Phys. **14** (2012) 013026.
- [9] N. Agafonova et al., *Search for $\nu_\mu \rightarrow \nu_e$ oscillations with the OPERA experiment in the CNGS beam*, JHEP **1307** (2013) 004.

- [10] C. H. Albright and C. Jarlskog, Nucl. Phys. B **84** (1975) 467.
- [11] M. H. Reno, Phys. Rev. D **74** (2006) 033001.
- [12] R. D. Ball et al., *Parton distributions for the LHC Run II* (2014).