

The β -function of Quantum Chromodynamics and the effective Higgs-gluon-gluon coupling in five-loop order

P. A. Baikov

*Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University
1(2), Leninskie Gory, Moscow 119991, Russian Federation*

K. G. Chetyrkin*

Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT), Germany

J. H. Kühn

Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT), Germany

We present the result of the analytical computation of the five-loop term in the beta function which governs the running of α_s — the quark-gluon coupling constant in QCD. Using the result, a low-energy theorem and the four-loop QCD decoupling formula, we derive the effective Higgs-gluon-gluon coupling constant in five-loop approximation.

*Loops and Legs in Quantum Field Theory
24-29 April 2016
Leipzig, Germany*

*Speaker.

Asymptotic freedom, manifest by a coupling constant decreasing with increasing energy, is one of the basic predictions of nonabelian gauge theories. It is quantitatively encoded in the behaviour of the β -function of Quantum Chromodynamics (QCD):

$$\beta(a_s) = \mu^2 \frac{d}{d\mu^2} a_s(\mu) = - \sum_{i \geq 0} \beta_i a_s^{i+2} \quad (1)$$

which describes the running of the quark-gluon coupling constant $a_s \equiv \alpha_s/\pi$ as a function of the renormalization scale μ . The four lowest terms were calculated in Refs. [1–8] with the result (we use the $\overline{\text{MS}}$ renormalization scheme throughout the paper):

$$\begin{aligned} \beta_0 &= \frac{1}{4} \left\{ 11 - \frac{2}{3} n_f \right\}, \\ \beta_1 &= \frac{1}{4^2} \left\{ 102 - \frac{38}{3} n_f \right\}, \\ \beta_2 &= \frac{1}{4^3} \left\{ \frac{2857}{2} - \frac{5033}{18} n_f + \frac{325}{54} n_f^2 \right\}, \\ \beta_3 &= \frac{1}{4^4} \left\{ \frac{149753}{6} + 3564 \zeta_3 - \left[\frac{1078361}{162} + \frac{6508}{27} \zeta_3 \right] n_f + \left[\frac{50065}{162} + \frac{6472}{81} \zeta_3 \right] n_f^2 + \frac{1093}{729} n_f^3 \right\}, \end{aligned}$$

where n_f denotes the number of active quark flavors. These results have moved the theory from qualitative agreement with experiment to precise quantitative predictions, covering a wide kinematical range, from τ -lepton decays up to LHC results.

Recently, the five-loop term has been evaluated with the result:

$$\begin{aligned} \beta_4 &= \frac{1}{4^5} \left\{ \frac{8157455}{16} + \frac{621885}{2} \zeta_3 - \frac{88209}{2} \zeta_4 - 288090 \zeta_5 \right. \\ &\quad + n_f \left[-\frac{336460813}{1944} - \frac{4811164}{81} \zeta_3 + \frac{33935}{6} \zeta_4 + \frac{1358995}{27} \zeta_5 \right] \\ &\quad + n_f^2 \left[\frac{25960913}{1944} + \frac{698531}{81} \zeta_3 - \frac{10526}{9} \zeta_4 - \frac{381760}{81} \zeta_5 \right] \\ &\quad + n_f^3 \left[-\frac{630559}{5832} - \frac{48722}{243} \zeta_3 + \frac{1618}{27} \zeta_4 + \frac{460}{9} \zeta_5 \right] \\ &\quad \left. + n_f^4 \left[\frac{1205}{2916} - \frac{152}{81} \zeta_3 \right] \right\}. \quad (2) \end{aligned}$$

As expected from the three and four-loop results, β_4 does not contain the higher transcendentals ζ_3^2 , ζ_6 and ζ_7 which do appear in individual diagrams. Note that the n_f^4 term is in full agreement with the result of [9] derived for a generic gauge group long time ago with a very different technique. Furthermore, quite recently, the n_f^3 term has been confirmed and even extended for a generic gauge group in [10].

In numerical form the coefficients $\beta_0 - \beta_4$ read

$$\begin{aligned}\beta_0 &\approx 2.75 - 0.166667 n_f, \\ \beta_1 &\approx 6.375 - 0.791667 n_f, \\ \beta_2 &\approx 22.3203 - 4.36892 n_f + 0.0940394 n_f^2, \\ \beta_3 &\approx 114.23 - 27.1339 n_f + 1.58238 n_f^2 + 0.0058567 n_f^3, \\ \beta_4 &\approx 524.558 - 181.799 n_f + 17.156 n_f^2 - 0.22586 n_f^3 - 0.0017993 n_f^4.\end{aligned}\quad (3)$$

For the particular cases of $n_f = 3, 4, 5$ and 6 we get:

$$\begin{aligned}\bar{\beta}(n_f = 3) &= 1 + 1.78 a_s + 4.47 a_s^2 + 20.99 a_s^3 + 56.59 a_s^4, \\ \bar{\beta}(n_f = 4) &= 1 + 1.54 a_s + 3.05 a_s^2 + 15.07 a_s^3 + 27.33 a_s^4, \\ \bar{\beta}(n_f = 5) &= 1 + 1.26 a_s + 1.47 a_s^2 + 9.83 a_s^3 + 7.88 a_s^4, \\ \bar{\beta}(n_f = 6) &= 1 + 0.93 a_s - 0.29 a_s^2 + 5.52 a_s^3 + 0.15 a_s^4,\end{aligned}\quad (4)$$

where $\bar{\beta} \equiv \frac{\beta(a_s)}{-\beta_0 a_s^2} = 1 + \sum_{i \geq 1} \bar{\beta}_i a_s^i$. A very modest growth of the coefficients is observed and the (apparent) convergence is better than one would expect from comparison with other examples. The smallness of the five-loop contributions in eqs. (4) implies that the final result is hardly effected when running α_s from the scale of the τ -lepton mass m_τ to the mass of the Z -boson.

The combined uncertainty in $\alpha_s^{(5)}(M_Z)$ induced by running and matching from m_τ to M_Z can be conservatively estimated by the shift in $\alpha_s^{(5)}(M_Z)$ produced by the use of five-loop running (and, consequently) four-loop matching instead of four-loop running (and three-loop matching). It amounts to a $8 \cdot 10^{-5}$ which is by a factor of three less than the similar shift made by the use of four-loop running instead of the the three-loop one. The final value of $\alpha_s^{(5)}(M_Z) = 0.1198 \pm 0.0015$ which follows from $\alpha_s^{(3)}(M_\tau)$ is in remarkably good agreement with the fit to electroweak precision data ([11]) $\alpha_s^{(5)}(M_Z) = 0.1197 \pm 0.0028$.

Our result for the β -function leads to a determination of the effective Higgs-gluon-gluon coupling, improved by one more loop. In the heavy top limit the Higgs boson couples directly to gluons via the effective Lagrangian of the form [12–15]

$$\mathcal{L}_{\text{eff}} = -2^{1/4} G_F^{1/2} H C_1(\mu^2/m_t^2, a_s(\mu)) G_{\nu\rho}^a G_{\nu\rho}^a. \quad (5)$$

The effective coupling constant $C_1(\mu^2/m_t^2, a_s(\mu))$ appears as a common factor in two quantities important for Higgs physics processes, namely, Higgs decay into gluons (one of the main decay channels for the Standard Model Higgs boson) and Higgs production via gluon fusion (the main Higgs production mode on LHC). It is expressible through massive tadpoles and was computed at four loops in 1997 [16] (long before the direct calculation of four-loop generic massive tadpoles started to be technically feasible). This happened to be possible due to a low energy theorem¹ (exact in all orders) [16]

$$C_1 = -\frac{1}{2} m_h^2 \frac{\partial}{\partial m_h^2} \ln \zeta_g^2, \quad (6)$$

¹Very recently the theorem has been redrived with a different technique [17] and even extended on the case of multi-Higgs couplings to gluons in Ref. [18].

which connects C_1 with the corresponding “decoupling” constant ζ_g defined as a conversion factor in the relation

$$\alpha'_s = \alpha'_s(\mu, \alpha_s, m_h) = \zeta_g^2(\mu, \alpha_s, m_h) \alpha_s(\mu), \quad \zeta_g^2 = 1 + \sum_{i \geq 1} a_s^i(\mu) d_{\text{MS},i},$$

where α'_s denotes the coupling constant in the effective QCD with the heavy quark, integrated out and n_l is the number of remaining active quark flavours.

Eq. (6) assumes that we are dealing with QCD with one heavy quark h and n_l massless ones and that Higgs field is coupled with the scalar heavy quark current via

$$-\frac{H}{v} m_h \bar{h} h.$$

The total number of active flavours is $n_f = n_l + 1$. For the case of the SM Higgs one should set $m_h = m_t$ and $n_f = 6$, $n_l = 5$. A more convenient form of the rhs of eq. (6) was found in Ref. [16]

$$C_1 = \frac{\pi}{2\alpha'_s[1 - 2\gamma_m(\alpha_s)]} \left[\beta'(\alpha'_s) - \beta(\alpha_s) \frac{\partial \alpha'_s}{\partial \alpha_s} \right]. \quad (7)$$

By counting the powers of α_s in the rhs of eq. (7) one can easily see that, say, the $(L+1)$ -loop contribution to C_1 is constructed from the $(L+1)$ loop β -function as well as from the quark mass anomalous dimension γ_m and the decoupling constant ζ_g^2 , with the latter two quantities being sufficient to be known at L loops *only*.

The decoupling constant ζ_g^2 is currently known at four loops; the corresponding coefficients d_i read²

$$d_{\text{MS},1} = -\frac{1}{6} \ell_{\mu m}, \quad (8)$$

$$d_{\text{MS},2} = \frac{11}{72} - \frac{11}{24} \ell_{\mu m} + \frac{1}{36} \ell_{\mu m}^2, \quad (9)$$

$$\begin{aligned} d_{\text{MS},3} = & \frac{564731}{124416} - \frac{82043}{27648} \zeta_3 - \frac{955}{576} \ell_{\mu m} + \frac{53}{576} \ell_{\mu m}^2 - \frac{1}{216} \ell_{\mu m}^3 \\ & + n_l \left[-\frac{2633}{31104} + \frac{67}{576} \ell_{\mu m} - \frac{1}{36} \ell_{\mu m}^2 \right], \end{aligned} \quad (10)$$

$$\begin{aligned} d_{\text{MS},4} = & \Delta_{\text{MS},4} + \frac{7391699}{746496} \ell_{\mu m} - \frac{2529743}{165888} \zeta_3 \ell_{\mu m} + \frac{2177}{3456} \ell_{\mu m}^2 - \frac{1883}{10368} \ell_{\mu m}^3 + \frac{1}{1296} \ell_{\mu m}^4 \\ & + n_l \left[-\frac{110341}{373248} \ell_{\mu m} + \frac{110779}{82944} \zeta_3 \ell_{\mu m} - \frac{1483}{10368} \ell_{\mu m}^2 - \frac{127}{5184} \ell_{\mu m}^3 \right] \\ & + n_l^2 \left[\frac{6865}{186624} \ell_{\mu m} - \frac{77}{20736} \ell_{\mu m}^2 + \frac{1}{324} \ell_{\mu m}^3 \right], \end{aligned} \quad (11)$$

²The original result for ζ_g^2 as obtained in Refs. [19, 20] was not completely analytical as a pair of relevant master integrals was then known only numerically. In the formula below we have used finding of Refs. [21, 22] to display the $\Delta_{\text{MS},4}$ in a completely analytical form (it was also done in Ref. [23]).

$$\begin{aligned}
\Delta_{\text{MS},4} = & \frac{291716893}{6123600} - \frac{76940219}{195955200} \pi^4 - \frac{2362581983}{87091200} \zeta_3 - \frac{12057583}{483840} \zeta_5 \\
& + \frac{9318467}{32659200} \pi^4 \ln 2 - \frac{3031309}{1306368} \pi^2 \ln^2 2 + \frac{340853}{816480} \pi^2 \ln^3 2 + \frac{3031309}{1306368} \ln^4 2 \\
& - \frac{340853}{1360800} \ln^5 2 + \frac{3031309}{54432} a_4 + \frac{340853}{11340} a_5 \\
& + n_l \left[-\frac{4770941}{2239488} - \frac{541549}{14929920} \pi^4 + \frac{3645913}{995328} \zeta_3 + \frac{115}{576} \zeta_5 \right. \\
& \quad \left. - \frac{685}{124416} \pi^2 \ln^2 2 + \frac{685}{124416} \ln^4 2 + \frac{685}{5184} a_4 \right] \\
& + n_l^2 \left[-\frac{271883}{4478976} + \frac{167}{5184} \zeta_3 \right]. \tag{12}
\end{aligned}$$

Here $\ell_{\mu m} = \ln \frac{\mu^2}{m^2(\mu)}$, $m(\mu)$ is the (running) heavy quark mass and $a_n = \text{Li}_n(1/2) = \sum_{i=1}^{\infty} 1/(2^i i^n)$.

The quark mass anomalous dimension γ_m is known at four loops since long [24, 25]. Finally, using our result for the β -function, ζ_g and γ_m we arrive at (μ_h is defined as $\mu_h = m_h(\mu_h)$):

$$\begin{aligned}
C_1 = & -\frac{1}{12} a_s(\mu_h) \left\{ 1 + \frac{11}{4} a_s(\mu_h) + a_s^2(\mu_h) \left[\frac{2821}{288} + n_l \left(-\frac{67}{96} \right) \right] \right. \\
& + a_s^3(\mu_h) \left[-\frac{4004351}{62208} + \frac{1305893}{13824} \zeta(3) + n_l \left(\frac{115607}{62208} - \frac{110779}{13824} \zeta(3) \right) + n_l^2 \left(-\frac{6865}{31104} \right) \right] \\
& + a_s^4(\mu_h) \left[\left(-\frac{348837110927}{223948800} + \frac{1959268669}{111974400} \pi^4 + \frac{35074684379}{24883200} \zeta_3 + \frac{127050929}{138240} \zeta_5 \right. \right. \\
& \quad \left. - \frac{121140071}{9331200} \pi^4 \ln 2 + \frac{39407017}{373248} \pi^2 \ln^2 2 - \frac{4431089}{233280} \pi^2 \ln^3 2 - \frac{39407017}{373248} \ln^4 2 \right. \\
& \quad \left. + \frac{4431089}{388800} \ln^5 2 - \frac{39407017}{15552} a_4 - \frac{4431089}{3240} a_5 \right) \\
& + n_l \left(\frac{223712155249}{1567641600} + \frac{190184767}{348364800} \pi^4 - \frac{108268867471}{348364800} \zeta_3 - \frac{22007549}{483840} \zeta_5 \right. \\
& \quad \left. + \frac{9318467}{10886400} \pi^4 \ln 2 - \frac{3896297}{580608} \pi^2 \ln^2 2 + \frac{340853}{272160} \pi^2 \ln^3 2 + \frac{3896297}{580608} \ln^4 2 \right. \\
& \quad \left. - \frac{340853}{453600} \ln^5 2 + \frac{3896297}{24192} a_4 + \frac{340853}{3780} a_5 \right) \\
& + n_l^2 \left(-\frac{8059709}{4478976} - \frac{576757}{4976640} \pi^4 + \frac{455777}{36864} \zeta_3 - \frac{115}{384} \zeta_5 - \frac{685}{41472} \pi^2 \ln^2 2 + \frac{685}{41472} \ln^4 2 + \frac{685}{1728} a_4 \right) \\
& \left. + n_l^3 \left(-\frac{270407}{1492992} + \frac{211}{1728} \zeta_3 \right) \right\}, \tag{13}
\end{aligned}$$

or, numerically,

$$\begin{aligned}
C_1 \approx & -\frac{1}{12} a_s(\mu_h) \left[1 + 2.75 a_s(\mu_h) + (9.795 - 0.698 n_l) a_s^2(\mu_h) \right. \\
& \quad \left. + (49.183 - 7.774 n_l - 0.221 n_l^2) a_s^3(\mu_h) \right. \\
& \quad \left. + (326.716 - 64.163 n_l + 1.594 n_l^2 - 0.0343 n_l^3) a_s^4(\mu_h) \right]. \tag{13}
\end{aligned}$$

The last expression should be modified if one want to use the pole quark mass M_h instead of the running $\overline{\text{MS}}$ one. Using the corresponding conversion formula from [26–28] we arrive at

$$\begin{aligned} C_1 \approx -\frac{1}{12} a_s(M_h) & \left[1 + 2.75 a_s(M_h) + (9.351 - 0.698 n_l) a_s^2(M_h) \right. \\ & + (43.609 - 6.538 n_l - 0.221 n_l^2) a_s^3(M_h) \\ & \left. + (240.383 - 30.231 n_l + 0.8001 n_l^2 - 0.0343 n_l^3) a_s^4(M_h) \right]. \end{aligned} \quad (14)$$

For the case of the SM Higgs we obtain:

$$C_1 = -\frac{1}{12} a_s(\mu_t) \left[1 + 2.750 a_s(\mu_t) + 6.306 a_s^2(\mu_t) + 4.794 a_s^3(\mu_t) + 41.447 a_s^4(\mu_t) \right]. \quad (15)$$

Note that the contribution due to β_4 to the last coefficient (boxed below) is significant, namely,

$$41.447 = -47.611 + \boxed{89.058}. \quad (16)$$

After transformation to the pole top quark mass eq. (15) reads

$$C_1 = -\frac{1}{12} a_s(M_t) \left[(1 + 2.750 a_s(M_t) + 5.861 a_s^2(M_t) + 5.3996 a_s^3(M_t) + 54.951 a_s^4(M_t)) \right], \quad (17)$$

with the last term decomposition assuming the form:

$$54.951 = -34.107 + \boxed{89.058}. \quad (18)$$

Let us close this paper with a brief discussion of its technical details³. The total number of five-loop diagrams contributing to the β -function (as generated by QGRAF [30]) amounts to about one and a half million. Every power of n_f in (2) was computed separately with the help of the FORM [31, 32] program BAICER, implementing the algorithm of works [33–35].

Our analytical results are also available in computer readable form under the URL
<http://www.ttp.kit.edu/Progdata/ttp16/ttp16-032>

We thank Johannes Blümlein for his interest to our work and encouragement.

The work by K. G. Chetyrkin and J. H. Kühn was supported by the Deutsche Forschungsgemeinschaft through CH1479/1-1. The work of P. A. Baikov is supported in part by grant NSh-7989.2016.2 of the President of Russian Federation.

References

- [1] D. J. Gross and F. Wilczek, *Ultraviolet Behavior of Nonabelian Gauge Theories*, *Phys.Rev.Lett.* **30** (1973) 1343–1346.
- [2] H. D. Politzer, *Reliable Perturbative Results for Strong Interactions?*, *Phys.Rev.Lett.* **30** (1973) 1346–1349.
- [3] W. E. Caswell, *Asymptotic Behavior of Nonabelian Gauge Theories to Two Loop Order*, *Phys.Rev.Lett.* **33** (1974) 244.
- [4] D. Jones, *Two Loop Diagrams in Yang-Mills Theory*, *Nucl.Phys.* **B75** (1974) 531.

³More information about our calculation and its implications can be found in Ref. [29].

- [5] O. Tarasov, A. Vladimirov and A. Y. Zharkov, *The Gell-Mann-Low Function of QCD in the Three Loop Approximation*, *Phys.Lett.* **B93** (1980) 429–432.
- [6] S. A. Larin and J. A. M. Vermaasen, *The three loop qcd beta function and anomalous dimensions*, *Phys. Lett.* **B303** (1993) 334–336, [[hep-ph/9302208](#)].
- [7] T. van Ritbergen, J. Vermaseren and S. Larin, *The Four loop beta function in quantum chromodynamics*, *Phys.Lett.* **B400** (1997) 379–384, [[hep-ph/9701390](#)].
- [8] M. Czakon, *The Four-loop QCD beta-function and anomalous dimensions*, *Nucl.Phys.* **B710** (2005) 485–498, [[hep-ph/0411261](#)].
- [9] J. Gracey, *The QCD Beta function at $\mathcal{O}(1/N_f)$* , *Phys.Lett.* **B373** (1996) 178–184, [[hep-ph/9602214](#)].
- [10] T. Luthe, A. Maier, P. Marquard and Y. Schroder, *Towards the five-loop Beta function for a general gauge group*, [1606.08662](#).
- [11] PARTICLE DATA GROUP collaboration, K. Olive et al., *Review of Particle Physics*, *Chin.Phys.* **C38** (2014) 090001.
- [12] F. Wilczek, *Decays of heavy vector mesons into higgs particles*, *Phys. Rev. Lett.* **39** (1977) 1304.
- [13] M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, *Low-Energy Theorems for Higgs Boson Couplings to Photons*, *Sov. J. Nucl. Phys.* **30** (1979) 711–716.
- [14] T. Inami, T. Kubota and Y. Okada, *Effective gauge theory and the effect of heavy quarks in higgs boson decays*, *Z. Phys.* **C18** (1983) 69.
- [15] B. A. Kniehl and M. Spira, *Low-energy theorems in higgs physics*, *Z. Phys.* **C69** (1995) 77–88, [[hep-ph/9505225](#)].
- [16] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, *Decoupling relations to $O(\alpha(s)^{**3})$ and their connection to low-energy theorems*, *Nucl. Phys.* **B510** (1998) 61–87, [[hep-ph/9708255](#)].
- [17] M. Kramer, E. Laenen and M. Spira, *Soft gluon radiation in higgs boson production at the lhc*, *Nucl. Phys.* **B511** (1998) 523–549, [[hep-ph/9611272](#)].
- [18] M. Spira, *Effective Multi-Higgs Couplings to Gluons*, [1607.05548](#).
- [19] K. G. Chetyrkin, J. H. Kühn and C. Sturm, *QCD decoupling at four loops*, *Nucl. Phys.* **B744** (2006) 121–135, [[hep-ph/0512060](#)].
- [20] Y. Schroder and M. Steinhauser, *Four-loop decoupling relations for the strong coupling*, *JHEP* **01** (2006) 051, [[hep-ph/0512058](#)].
- [21] B. A. Kniehl and A. V. Kotikov, *Heavy-quark QCD vacuum polarisation function: Analytical results at four loops*, *Phys. Lett.* **B642** (2006) 68–71, [[hep-ph/0607201](#)].
- [22] R. N. Lee and I. S. Terekhov, *Application of the DRA method to the calculation of the four-loop QED-type tadpoles*, *JHEP* **01** (2011) 068, [[1010.6117](#)].
- [23] B. A. Kniehl, A. V. Kotikov, A. I. Onishchenko and O. L. Veretin, *Strong-coupling constant with flavor thresholds at five loops in the anti-MS scheme*, *Phys. Rev. Lett.* **97** (2006) 042001, [[hep-ph/0607202](#)].
- [24] K. G. Chetyrkin, *Quark mass anomalous dimension to $O(\alpha(s)^{**4})$* , *Phys. Lett.* **B404** (1997) 161–165, [[hep-ph/9703278](#)].

- [25] J. A. M. Vermaseren, S. A. Larin and T. van Ritbergen, *The 4-loop quark mass anomalous dimension and the invariant quark mass*, *Phys. Lett.* **B405** (1997) 327–333, [[hep-ph/9703284](#)].
- [26] N. Gray, D. J. Broadhurst, W. Grafe and K. Schilcher, *Three loop relation of quark (modified) ms and pole masses*, *Z. Phys.* **C48** (1990) 673–680.
- [27] K. G. Chetyrkin and M. Steinhauser, *Short distance mass of a heavy quark at order alpha(s)**3*, *Phys. Rev. Lett.* **83** (1999) 4001–4004, [[hep-ph/9907509](#)].
- [28] K. Melnikov and T. v. Ritbergen, *The three-loop relation between the ms-bar and the pole quark masses*, *Phys. Lett.* **B482** (2000) 99–108, [[hep-ph/9912391](#)].
- [29] P. A. Baikov, K. G. Chetyrkin and J. H. Kühn, *Five-Loop Running of the QCD coupling constant*, [1606.08659](#).
- [30] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279–289.
- [31] J. A. M. Vermaseren, *New features of FORM*, [math-ph/0010025](#).
- [32] M. Steinhauser, T. Ueda and J. A. M. Vermaseren, *Parallel versions of FORM and more*, *Nucl. Part. Phys. Proc.* **261-262** (2015) 45–57, [[1501.07119](#)].
- [33] P. A. Baikov, *A practical criterion of irreducibility of multi-loop feynman integrals*, *Phys. Lett.* **B634** (2006) 325–329, [[hep-ph/0507053](#)].
- [34] P. A. Baikov, *Explicit solutions of the 3-loop vacuum integral recurrence relations*, *Phys. Lett.* **B385** (1996) 404–410, [[hep-ph/9603267](#)].
- [35] P. A. Baikov, *Recurrence relations in the large space-time dimension limit*, *PoS RADCOR2007* (2007) 022.