

New Results on Massive 3-Loop Wilson Coefficients in Deep-Inelastic scattering^{*†‡}

J. Ablinger

*Research Institute for Symbolic Computation (RISC), Johannes Kepler University,
Altenbergerstraße 69, A-4040, Linz, Austria*

A. Behring, J. Blümlein, G. Falcioni, A. De Freitas

Deutsches Elektronen-Synchrotron, DESY, Platanenallee 6, D-15738 Zeuthen, Germany

A. Hasselhuhn

*Institut für Theoretische Teilchenphysik Campus Süd, Karlsruher Institut für Technologie (KIT),
D-76128 Karlsruhe, Germany*

A. von Manteuffel

*Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA
PRISMA Cluster of Excellence, Johannes Gutenberg University, 55099 Mainz, Germany*

M. Round, C. Schneider

*Research Institute for Symbolic Computation (RISC), Johannes Kepler University,
Altenbergerstraße 69, A-4040, Linz, Austria*

F. Wißbrock

IHES, 35 Route de Chartres, F-91440 Bures-sur-Yvette, France

We present recent results on newly calculated 2- and 3-loop contributions to the heavy quark parts of the structure functions in deep-inelastic scattering due to charm and bottom.

Loops and Legs in Quantum Field Theory

24-29 April 2016

Leipzig, Germany

^{*}This work was supported in part by the Austrian Science Fund (FWF) grant SFB F50 (F5009-N15), the European Commission through contract PITN-GA-2012-316704 (HIGGSTOOLS) and by FP7 ERC Starting Grant 257638 PAGAP.

[†]Based on the talks presented by A. Behring, A. De Freitas, and G. Falcioni.

[‡]Also contribution to the Proceedings of DIS 2016.

1. Introduction

At present the 3-loop heavy flavor corrections form the missing link to analyze the World deep-inelastic scattering data at next-to-next-to leading order with respect to the determination of the unpolarized parton distribution functions (PDFs) [1], the strong coupling constant [2], and the masses of the charm and bottom quarks [3]. These distributions and parameters form an essential input for all precision measurements at the LHC, notably the measurement of the Higgs boson [4] and $t\bar{t}$ [5] production cross sections. Furthermore, their precise knowledge provides new unprecedented tests of the Standard Model and may help in this way to reveal potential deviations pointing to new physics.

The present project is devoted to calculate the 3-loop heavy flavor Wilson coefficients at large scales $Q^2 \gg m^2$ in analytic form. In case of the structure function $F_2(x, Q^2)$ this approximation suffices at the 1% level for scales $Q^2/m^2 \gtrsim 10$ [6], a region to be selected anyway in case of charm to stay off higher twist terms [7] at this precision.

In 2009 a series of Mellin moments has been calculated for the corresponding massive operator matrix elements (OMEs) ranging from $N = 2$ to $N = 10 \dots 14$, depending on the Wilson coefficient, in Refs. [8] by mapping the operator matrix elements for the different moments onto massive tadpoles, which were computed using the code MATAD [9]. These moments serve now for comparison in the computation of the general- N results. In the time since, four out of five neutral current massive 3-loop Wilson coefficients have been calculated [10–14], along with seven out of eight massive OMEs [10–13, 15–19]. For all processes the logarithmic 3-loop contributions are known [11]. To perform these computations, several technical, computer-algebraic and mathematical innovations were necessary, which are described in Refs. [20–36]. The different physics results, including also necessary 2-loop calculations, for the neutral and charged current reactions were published in Refs. [37–41] and recent surveys were given in [18, 42].

In this note we describe recent developments of the project. The paper is organized as follows. The basic formalism and a series of technical details of the calculation are summarized in Section 2. In Section 3 we review the status of the calculation of the neutral current structure function $F_2(x, Q^2)$. A survey on the calculation of the 3-loop corrections to non-singlet charged and neutral current structure functions is given in Section 4. Results on the calculation of the heavy flavor corrections in the full region of Q^2 for various polarized and unpolarized non-singlet structure functions to $O(\alpha_s^2)$ and associated sum rules are given in Section 5. Recent results of 3-loop two-mass corrections are reported in Section 6, and Section 7 contains the conclusions.

2. Basic Formalism and Technical Aspects of the Calculation

As it has been outlined in Refs. [8, 11] the massive 3-loop Wilson coefficients for deep-inelastic scattering can be represented in terms of the massless Wilson coefficients [43, 44] and massive operator matrix elements in the asymptotic region $Q^2 \gg m^2$, if the single heavy quark case is considered. Here Q^2 denotes the virtuality of the process and m the heavy quark mass. In the present project the massive OMEs are newly computed and used in the calculation of the corresponding massive Wilson coefficients. Furthermore, the massive OMEs describe the transition of a single heavy flavor becoming light in the variable flavor number scheme (VFNS), cf. [11, 45]. Up to

2-loop order the Wilson coefficients are known (semi)analytically [46]¹. In the asymptotic region analytic expressions have been derived in Refs. [6, 37].

The whole 3-loop project constitutes a calculation of 2864 Feynman diagrams mapping to some dozens of 10^4 scalar integrals. They are generated using `QGraf` [48], calculating their Dirac structure using `FORM` [49] and the color structure using `Color` [50]. Except for simpler topologies, we use the integration by part (IBP)-reduction [51] implemented in the package `Reduze 2` [52]² to map the major part of the problem to the calculation of 687 master integrals, out of which 116 remain to be calculated. 571 master integrals can be solved by sum-structures using difference-field theory [55–63] implemented in the package `Sigma` [64, 65], or other techniques, like the use of hypergeometric functions [66], Mellin-Barnes representations [67], the method of hyperlogarithms [33, 68], the solution of differential equations [69], and the Almkvist-Zeilberger algorithm [70], implemented within the package `MultiIntegrate` [71].

They can be expressed by iterative sum-structures in N -space or iterative integrals in x -space, i.e. in terms of harmonic polylogarithms [72], generalized harmonic polylogarithms of the Kummer-type [28, 29], cyclotomic harmonic polylogarithms [31], or root-valued iterated integrals [30], which all correspond to difference or differential equations that can be factorized to first order. This is not the case for all the remaining 116 master integrals, which depend on structures obeying 2nd order equations. Here we expect complete elliptic integrals of rational argument and related functions to emerge, see [73]³, over which first order structures are iterated again. We currently work on the solution of these systems.

In all these methods the solution of recurrences and the various properties of special functions emerging in this context play a central role and have to be used algorithmically. This is made possible by the packages `Sigma` [64, 65], `EvaluateMultiSums` and `SumProduction`, [77], `RhoSum` [78], decoupling formalisms [79], `HarmonicSums` [29–31, 71, 80] and `MultiIntegrate` [71].

3. Status of the 3-Loop Neutral Current Corrections

The heavy flavor contributions to the structure functions $F_{2,L}(x, Q^2)$ are given by

$$F_{2,L}^{\text{heavy}}(x, Q^2) = F_{2,L}(x, Q^2) - F_{2,L}^{\text{massless}}(x, Q^2), \quad (3.1)$$

for a single heavy quark Q and N_F massless quarks

$$\begin{aligned} F_{(2,L)}^{\text{heavy}}(x, N_F + 1, Q^2, m^2) = & \\ & \sum_{k=1}^{N_F} e_k^2 \left\{ L_{q,(2,L)}^{\text{NS}} \left(x, N_F + 1, \frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) \otimes \left[f_k(x, \mu^2, N_F) + f_{\bar{k}}(x, \mu^2, N_F) \right] \right. \\ & \left. + \frac{1}{N_F} L_{q,(2,L)}^{\text{PS}} \left(x, N_F + 1, \frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) \otimes \Sigma(x, \mu^2, N_F) \right\} \end{aligned}$$

¹For a precise implementation in Mellin space, see [47].

²The package uses the codes `GiNaC` [53] and `FerMat` [54].

³Similar structures were observed also in Refs. [74–76].

$$\begin{aligned}
 & \left. + \frac{1}{N_F} L_{g,(2,L)}^S \left(x, N_F + 1, \frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) \otimes G(x, \mu^2, N_F) \right\} \\
 & + e_Q^2 \left[H_{q,(2,L)}^{\text{PS}} \left(x, N_F + 1, \frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) \otimes \Sigma(x, \mu^2, N_F) \right. \\
 & \left. + H_{g,(2,L)}^S \left(x, N_F + 1, \frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) \otimes G(x, \mu^2, N_F) \right]. \quad (3.2)
 \end{aligned}$$

Here e_k and e_Q denote the light and heavy quark charges, f_k and G are the quarkonic and gluon parton densities and $\Sigma = \sum_{k=1}^{N_F} (f_k + \bar{f}_k)$ the singlet density, and μ denotes the factorization scale. The five heavy flavor Wilson coefficients are given by $L_q^{\text{NS}}, L_q^{\text{PS}}, L_g^S$ and H_q^{PS}, H_g^S , cf. Refs. [10–13].

In Figure 1 the present status of the 3-loop results on $F_2^{\text{charm}}(x, Q^2)$ is illustrated for $Q^2 = 100 \text{ GeV}^2$ referring to the parton densities [81] and $m_c^{\text{pole}} = 1.59 \text{ GeV}$. We show the different contributions in $O(\alpha_s), O(\alpha_s^2)$ and $O(\alpha_s^3)$ using the asymptotic representation, highlighting their behaviour at large values of x in the inset.




Figure 1: The different known contributions to $F_2^{\text{charm}}(x, Q^2)$ by the five heavy flavor Wilson coefficients from $O(\alpha_s)$ (dotted lines), $O(\alpha_s^2)$ (dashed lines), $O(\alpha_s^3)$ (full lines), for $m_c^{\text{pole}} = 1.59 \text{ GeV}$ and the PDFs [81]. The contributions due to H_g^S have been scaled down by a factor of 20 for better visibility.

The intended accuracy is $O(1\%)$ and requires all the contributions mentioned above, at least in some part of the kinematic range. The largest contribution is due to H_g^S driven by the gluon distribution starting at $O(\alpha_s)$. Here the calculation of the 3-loop term is underway, while all other contributions are known already. The next largest term in the small x region is H_q^{PS} , yielding negative corrections there. It is followed by L_g^S with smaller positive corrections at 3-loop order, which are larger than those in 2-loop order by the same Wilson coefficient. The fourth largest contribution is due to L_q^{PS} and the smallest contributions are due to L_q^{NS} , while it is largest in

the valence region. The full knowledge of the 3-loop heavy flavor corrections to $F_2(x, Q^2)$ will improve on presently remaining theory errors both on $\alpha_s(M_Z^2)$ and m_c [3] due to a yet approximate treatment [82] of these corrections, based on the previous work [8, 10, 37–39].

4. 3-Loop Non-Singlet Polarized and Charged Current Corrections

The flavor non-singlet OME [12] appears in various unpolarized and polarized neutral and charged current structure functions, combined with the different massless Wilson coefficients and helicity dependent anomalous dimensions [44, 83–85]. This allows to calculate the polarized 3-loop massive non-singlet Wilson coefficient for twist-2 part of the structure functions $g_{1,2}(x, Q^2)$ and the charged current structure functions $xF_3^{W^+-W^-}(x, Q^2)$ and $F_{1,2}^{W^+-W^-}(x, Q^2)$, cf. Refs. [86–88]; for the lower order corrections see [40, 89]. Here, g_2 is obtained by the Wandzura-Wilczek relation [90–93].

In Figure 2 we show the relative 3-loop corrections to $g_1^{\text{NS}}(x, Q^2)$ and $xF_3^{W^++W^-}(x, Q^2)$ for typical scales Q^2 using the polarized PDFs [94] and the unpolarized PDFs [81], normalizing to the massless contributions.




Figure 2: The relative 3-loop charm quark corrections to the polarized neutral current structure function $g_1^{\text{NS}}(x, Q^2)$ using the PDFs [94] and the unpolarized charged current structure function $xF_3^{W^++W^-}(x, Q^2)$ using the PDFs [81] for $m_c^{\text{pole}} = 1.59 \text{ GeV}$ as functions of x and Q^2 ; from [86] and [87].

The 3-loop heavy flavor corrections to g_1^{NS} vary between +2% and –5% and those for the charged current structure function $xF_3^{W^++W^-}$ between +3% and –3%, compared to the light flavor contributions. These corrections cannot be resolved in present measurements but will play a role in high luminosity measurements at planned future colliders [95].

The asymptotic 3-loop heavy flavor corrections to the charged current structure functions $F_{1(2)}^{W^+-W^-}$ is shown in Figure 3. The relative contribution ranges from –1(–2)% to –8%, with the largest corrections around $x \sim 0.03$. The higher order terms yield positive corrections.

5. Power Corrections in the Non-Singlet Case at $O(\alpha_s^2)$

In the foregoing sections we have presented results calculating the massive Wilson coefficients in the asymptotic region $Q^2 \gg m^2$. It is, however, also interesting to study the Wilson coefficients at lower scales Q^2 for data in this region and to estimate the scales at which the asymptotic representation is valid. We investigate the non-singlet massive Wilson coefficients to $O(\alpha_s^2)$ for the inclusive structure functions [41].




Figure 3: The relative 3-loop charm quark corrections to the charged current structure functions $F_{1(2)}^{W^+ - W^-}(x, Q^2)$ at $Q^2 = 100 \text{ GeV}^2$ for $m_c^{\text{pole}} = 1.59 \text{ GeV}$ using the PDFs of Ref. [81]; form [88].




Figure 4: Left panel: The charm and bottom quark contributions to the polarized Bjorken sum rule as a function of $\xi_c = Q^2/m_c^2$, for $m_c^{\text{pole}} = 1.59 \text{ GeV}$ and $m_b^{\text{pole}} = 4.78 \text{ GeV}$ [96]. The function C_{pBj} describes the flavor excitation, with $C_{\text{pBj}} = 1$ one heavy quark being effectively massless. Right panel: The 2-loop charm quark contribution to the unpolarized Bjorken sum rule; from [41].

In the tagged flavor case a similar analysis was performed in Ref. [97]⁴. It turns out that the tagged flavor case does not lead to a stable description at large scales Q^2 , containing logarithmic terms, which are absent in the inclusive case due to heavy flavor loop effects in diagrams with massless final states. In the inclusive case the decoupling of heavy flavors is uniquely described. Going to very low scales Q^2 for deep-inelastic scattering logarithmic terms remain. However, the deep-inelastic description is limited to virtualities Q^2 of a certain size, say $Q^2 \gtrsim Q_0^2 = 5 \text{ GeV}^2$ and cannot be applied below.

We show the effect of these contributions on the structure functions $F_{1(2)}$ in Figure 3 (dash-dotted lines) replacing the contributions up to 2-loops by the complete results. At $Q^2 = 100 \text{ GeV}^2$ the corrections do widely agree with the asymptotic result but a small difference in the large x range still remains.

One also obtains interesting results on different sum rules for the structure functions, such as the Adler- [98], unpolarized Bjorken- [99], polarized Bjorken- [100], and Gross-Lewellyn Smith

⁴For the $O(\alpha_s^2)$ results on the tagged contributions to the structure functions see [6,37].

sum rule [101]. While due to fermion number conservation the contribution to these sum rules is free of mass effects in the asymptotic range, the last three sum rules receive heavy quark mass effects. We have shown that the Adler sum rule does not receive any correction, also no target mass corrections.

Figure 4 demonstrates the transition of the massive quark contributions to massless ones for the massive 2-loop corrections of the polarized and unpolarized Bjorken sum-rule as a function of Q^2 . One should note that the respective corrections enter the N_F -terms of the sum rule only, cf. [41]. The transition of this term to 1 only proceeds slowly both for charm and bottom quarks as a function of Q^2 . At lower scales Q^2 the corresponding contributions are even negative in case of the polarized Bjorken sum rule due to virtual corrections, which is of importance in experimental analyses.

6. 3-Loop Two-Mass Corrections in the Asymptotic Region

Starting at 3-loop order the massive Wilson coefficients contain Feynman diagrams in which two internal massive fermion lines are present leading to contributions of charm and bottom in individual terms, which cannot be separated. Due to this the variable flavor number scheme presented in [11, 45] is no longer applicable in the strict sense, but needs to be generalized. As $m_b^2/m_c^2 \sim 10$, one possibility for large scales $Q^2 \gg m_c^2, m_b^2$ consists in decoupling both heavy quarks together. The corresponding variable flavor number scheme has been worked out in Ref. [102] and requires to obtain as well the two-mass contributions to the different OMEs at 3-loop order. Their moments $N = 2, 4, 6$ have already been calculated [102, 103] projecting them to massive tadpoles and using the package Q2e [104]. In the two-mass case the renormalization of the OMEs, given in the single mass case in [8], has to be extended. Furthermore, the 2-mass OMEs $A_{qq,Q}^{\text{NS}}$, $A_{qq,Q}^{\text{NS,TR}}$ and $A_{gq,Q}$ have been calculated for general mass assignment analytically and the calculation of A_{Qq}^{PS} is underway. In case of the OME $A_{gg,Q}$ all scalar topologies have been calculated. Here new classes of iterative integrals do emerge, which can be handled with the package `HarmonicSums.m` by now. While an expansion in the mass ratio m_c^2/m_b^2 is possible for fixed moments, it turns out not to be possible for a general Mellin variable N , thus requiring the complete calculation.

7. Conclusions

After having calculated a set of 3-loop moments for all massive OMEs contributing to the massive 3-loop Wilson coefficients in the asymptotic region $Q^2 \gg m^2$ in 2009 [8], progress has been made in the calculation of the general N results of these Wilson coefficients. All logarithmic contributions are known [11] and the Wilson coefficients $L_q^{\text{NS}}, L_q^{\text{PS}}, L_g^{\text{S}}$ and H_q^{PS} have been calculated along with the OMEs appearing as the transition matrix elements in the variable flavor number scheme at 3-loop order, cf. [10–13]. In case of the charged current structure functions and the non-singlet contribution to the polarized structure functions $g_{1(2)}(x, Q^2)$ a series of 3-loop Wilson coefficients have been calculated [86–88], completing the 2-loop charged current programme [40, 89] before. Recently, the non-singlet $O(\alpha_s^2)$ corrections have been completed for the most important inclusive polarized and unpolarized neutral and charged current structure functions and the sum rules associated to them in the whole range Q^2 relevant for deep-inelastic scattering [41]. It has been shown

that the process of a heavy quark becoming effectively massless at high scales proceeds slowly, which should be considered in the matching in the different schemes being used in data analysis currently. Clearly, the matching at $m_Q^2 = Q^2$ is unfortunate since a heavy quark's velocity is not ultra-relativistic there.

At 3-loop order two-heavy-mass Feynman diagrams contribute to the OMEs. This requires a different renormalization compared to that of the single-heavy-mass case and leads to a change of the associated variable flavor number scheme since these terms are neither charm nor bottom contributions. One may, however, design a variable flavor number scheme in decoupling both contributions together. Beyond a series of moments also the general N contributions have been computed in some cases [102].

We have devised an algorithm to calculate massive Feynman diagrams containing local operator insertions, mapping the differential equations obtained from the IBP-relations into systems of difference equations, which are solved automatically [34, 35]. This algorithm factorizes the corresponding problem to first order structures as far as possible and into potential remaining terms, which cannot be factorized neither in N - nor x -space. If the latter terms are not present, we receive an iterative sum or integral solution of the corresponding physical problem starting from differential equations in whatever basis and obtaining the emerging letter-representation of the contributing alphabet in an automatic manner together with a proof certificate. In this way we automatically detect non-factorizing contributions uniquely. They need a further separate treatment to obtain the solution.

In the present project various new mathematical structures have been found for Feynman-integrals in general. While at the time of 1998 the harmonic sums yielded a sufficient systematic representation for massless 2-loop problems [20, 21], the present massive 3-loop problems require Kummer-iterated integrals [28, 29], cyclotomic iterated integrals [31], root-valued iterated integrals [30], and further generalizations to elliptic and ${}_2F_1$ -valued non-iterative letters in otherwise iterative integrals [73]. This list of structures will be growing further addressing processes at even higher loops and with more scales. The packages `Sigma`, `EvaluateMultiSums`, `SumProduction`, `RhoSum`, `HarmonicSums` and `MultiIntegrate` [29–31, 64, 65, 71, 77–80] were significantly extended or newly created in the project and are already in partly use in a series of other projects.

Presently we work on the completion of $A_{Qg}^{(3)}$, in which new mathematical structures appear, the solution of which has to be automated.

References

- [1] A. Accardi *et al.*, *A Critical Appraisal and Evaluation of Modern PDFs*, Eur. Phys. J. C **76** (2016) no.8, 471 [arXiv:1603.08906 [hep-ph]].
- [2] S. Bethke *et al.*, *Workshop on Precision Measurements of α_s* , arXiv:1110.0016 [hep-ph];
S. Moch, S. Weinzierl *et al.*, *High precision fundamental constants at the TeV scale*, arXiv:1405.4781 [hep-ph];
S. Alekhin, J. Blümlein and S.O. Moch, *α_s from global fits of parton distribution functions*, Mod. Phys. Lett. A **31** (2016) no.25, 1630023.
- [3] S. Alekhin, J. Blümlein, K. Daum, K. Lipka and S. Moch, *Precise charm-quark mass from deep-inelastic scattering*, Phys. Lett. B **720** (2013) 172 [arXiv:1212.2355 [hep-ph]];

- S. Alekhin, J. Blümlein, S.-O. Moch, R. Plačakytė, *The new ABMP16 PDFs*, DESY 16-121; DESY 16-179.
- [4] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog, A. Lazopoulos and B. Mistlberger, *High precision determination of the gluon fusion Higgs boson cross-section at the LHC*, JHEP **1605** (2016) 058 [arXiv:1602.00695 [hep-ph]].
- [5] M. Czakon, P. Fiedler and A. Mitov, *Total Top-Quark Pair-Production Cross Section at Hadron Colliders Through $O(\alpha_s^4)$* , Phys. Rev. Lett. **110** (2013) 252004 [arXiv:1303.6254 [hep-ph]].
- [6] M. Buza, Y. Matiounine, J. Smith, R. Migneron and W.L. van Neerven, *Heavy quark coefficient functions at asymptotic values $Q^2 \gg m^2$* , Nucl. Phys. B **472** (1996) 611 [hep-ph/9601302].
- [7] S. Alekhin, J. Blümlein and S. Moch, *Parton Distribution Functions and Benchmark Cross Sections at NNLO*, Phys. Rev. D **86** (2012) 054009 [arXiv:1202.2281 [hep-ph]].
- [8] I. Bierenbaum, J. Blümlein and S. Klein, *Mellin Moments of the $O(\alpha_s^3)$ Heavy Flavor Contributions to unpolarized Deep-Inelastic Scattering at $Q^2 \gg m^2$ and Anomalous Dimensions*, Nucl. Phys. B **820** (2009) 417 [arXiv:0904.3563 [hep-ph]];
J. Blümlein, S. Klein and B. Tödtli, *$O(\alpha_s^2)$ and $O(\alpha_s^3)$ Heavy Flavor Contributions to Transversity at $Q^2 \gg m^2$* , Phys. Rev. D **80** (2009) 094010 [arXiv:0909.1547 [hep-ph]].
- [9] M. Steinhauser, *MATAD: A Program package for the computation of MASSive TADpoles*, Comput. Phys. Commun. **134** (2001) 335 [hep-ph/0009029].
- [10] J. Ablinger, J. Blümlein, S. Klein, C. Schneider and F. Wißbrock, *The $O(\alpha_s^3)$ Massive Operator Matrix Elements of $O(N_F)$ for the Structure Function $F_2(x, Q^2)$ and Transversity*, Nucl. Phys. B **844** (2011) 26 [arXiv:1008.3347 [hep-ph]].
- [11] A. Behring, I. Bierenbaum, J. Blümlein, A. De Freitas, S. Klein and F. Wißbrock, *The logarithmic contributions to the $O(\alpha_s^3)$ asymptotic massive Wilson coefficients and operator matrix elements in deeply inelastic scattering*, Eur. Phys. J. C **74** (2014) 9, 3033 [arXiv:1403.6356 [hep-ph]].
- [12] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider, and F. Wißbrock, *The 3-Loop Non-Singlet Heavy Flavor Contributions and Anomalous Dimensions for the Structure Function $F_2(x, Q^2)$ and Transversity*, Nucl. Phys. B **886** (2014) 733 [arXiv:1406.4654 [hep-ph]].
- [13] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, *The 3-loop pure singlet heavy flavor contributions to the structure function $F_2(x, Q^2)$ and the anomalous dimension*, Nucl. Phys. B **890** (2014) 48 [arXiv:1409.1135 [hep-ph]].
- [14] J. Blümlein, A. De Freitas, W.L. van Neerven and S. Klein, *The Longitudinal Heavy Quark Structure Function $F_L^{Q\bar{Q}}$ in the Region $Q^2 \gg m^2$ at $O(\alpha_s^3)$* , Nucl. Phys. B **755** (2006) 272 [hep-ph/0608024].
- [15] J. Blümlein, A. Hasselhuhn, S. Klein and C. Schneider, *The $O(\alpha_s^3 N_F T_F^2 C_{A,F})$ Contributions to the Gluonic Massive Operator Matrix Elements*, Nucl. Phys. B **866** (2013) 196 [arXiv:1205.4184 [hep-ph]].
- [16] J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round, C. Schneider and F. Wißbrock, *The Transition Matrix Element $A_{gq}(N)$ of the Variable Flavor Number Scheme at $O(\alpha_s^3)$* , Nucl. Phys. B **882** (2014) 263 [arXiv:1402.0359 [hep-ph]].
- [17] J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, M. Round and C. Schneider, *The $O(\alpha_s^3 T_F^2)$ Contributions to the Gluonic Operator Matrix Element*, [arXiv:1405.4259 [hep-ph]].

- [18] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel, C.G. Raab, M. Round, C. Schneider and F. Wißbrock, *3-Loop Corrections to the Heavy Flavor Wilson Coefficients in Deep-Inelastic Scattering*, PoS (EPS-HEP2015) (2015) 504 [arXiv:1602.00583 [hep-ph]].
- [19] J. Ablinger et al., DESY 15–112.
- [20] J. Blümlein and S. Kurth, *Harmonic sums and Mellin transforms up to two-loop order*, Phys. Rev. D **60** (1999) 014018 [arXiv:hep-ph/9810241].
- [21] J.A.M. Vermaseren, *Harmonic sums, Mellin transforms and integrals*, Int. J. Mod. Phys. A **14** (1999) 2037 [arXiv:hep-ph/9806280].
- [22] J. Blümlein, *Algebraic relations between harmonic sums and associated quantities*, Comput. Phys. Commun. **159** (2004) 19 [hep-ph/0311046].
- [23] J. Blümlein, *Structural Relations of Harmonic Sums and Mellin Transforms up to Weight $w = 5$* , Comput. Phys. Commun. **180** (2009) 2218 [arXiv:0901.3106 [hep-ph]].
- [24] J. Blümlein, *Structural Relations of Harmonic Sums and Mellin Transforms at Weight $w = 6$* , in: Proceedings of the Workshop *Motives, Quantum Field Theory, and Pseudodifferential Operators*, Clay Mathematics Institute, Boston University, June 2–13, 2008, Clay Mathematics Proceedings Vol. **12** (2010) 167, eds. A. Carey, D. Ellwood, S. Paycha, S. Rosenberg, [arXiv:0901.0837 [math-ph]].
- [25] J. Blümlein, D.J. Broadhurst and J.A.M. Vermaseren, *The Multiple Zeta Value Data Mine*, Comput. Phys. Commun. **181** (2010) 582 [arXiv:0907.2557 [math-ph]].
- [26] J. Blümlein, M. Kauers, S. Klein and C. Schneider, *Determining the closed forms of the $O(a_s^3)$ anomalous dimensions and Wilson coefficients from Mellin moments by means of computer algebra*, Comput. Phys. Commun. **180** (2009) 2143 [arXiv:0902.4091 [hep-ph]].
- [27] J. Blümlein, S. Klein, C. Schneider and F. Stan, *A Symbolic Summation Approach to Feynman Integral Calculus*, J. Symbolic Comput. **47** (2012) 1267–1289 [arXiv:1011.2656 [cs.SC]].
- [28] S. Moch, P. Uwer and S. Weinzierl, *Nested sums, expansion of transcendental functions and multiscale multiloop integrals*, J. Math. Phys. **43** (2002) 3363 [hep-ph/0110083].
- [29] J. Ablinger, J. Blümlein and C. Schneider, *Analytic and Algorithmic Aspects of Generalized Harmonic Sums and Polylogarithms*, J. Math. Phys. **54** (2013) 082301 [arXiv:1302.0378 [math-ph]].
- [30] J. Ablinger, J. Blümlein, C.G. Raab and C. Schneider, *Iterated Binomial Sums and their Associated Iterated Integrals*, J. Math. Phys. **55** (2014) 112301 [arXiv:1407.1822 [hep-th]].
- [31] J. Ablinger, J. Blümlein and C. Schneider, *Harmonic Sums and Polylogarithms Generated by Cyclotomic Polynomials*, J. Math. Phys. **52** (2011) 102301 [arXiv:1105.6063 [math-ph]].
- [32] J. Ablinger, J. Blümlein, A. Hasselhuhn, S. Klein, C. Schneider and F. Wißbrock, *Massive 3-loop Ladder Diagrams for Quarkonic Local Operator Matrix Elements*, Nucl. Phys. B **864** (2012) 52 [arXiv:1206.2252 [hep-ph]].
- [33] J. Ablinger, J. Blümlein, C. Raab, C. Schneider and F. Wißbrock, *Calculating Massive 3-loop Graphs for Operator Matrix Elements by the Method of Hyperlogarithms*, Nucl. Phys. B **885** (2014) 409 [arXiv:1403.1137 [hep-ph]].
- [34] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, *Calculating Three Loop Ladder and V-Topologies for Massive Operator Matrix Elements by Computer Algebra*, Comput. Phys. Commun. **202** (2016) 33 [arXiv:1509.08324 [hep-ph]].

- [35] J. Ablinger, A. Behring, J. Blümlein, A. de Freitas and C. Schneider, *Algorithms to solve coupled systems of differential equations in terms of power series*, arXiv:1608.05376 [cs.SC];
 J. Ablinger, J. Blümlein, A. de Freitas, C. Schneider, *A toolbox to solve coupled systems of differential and difference equations*, In: Proc. of 12th International Symposium on Radiative Corrections (Radcor 2015) and LoopFest XIV (Radiative Corrections for the LHC and Future Colliders), PoS (RADCOR2015) 060, pp. 1-13 [arXiv:1601.01856].
- [36] J. Ablinger, J. Blümlein and C. Schneider, *Generalized Harmonic, Cyclotomic, and Binomial Sums, their Polylogarithms and Special Numbers*, J. Phys. Conf. Ser. **523** (2014) 012060 [arXiv:1310.5645 [math-ph]];
 J. Ablinger and J. Blümlein, in: *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions*, C. Schneider, J. Blümlein, Eds., p. 1, (Springer, Wien, 2013) *Harmonic Sums, Polylogarithms, Special Numbers, and their Generalizations*, [arXiv:1304.7071 [math-ph]].
- [37] I. Bierenbaum, J. Blümlein and S. Klein, *Two-Loop Massive Operator Matrix Elements and Unpolarized Heavy Flavor Production at Asymptotic Values $Q^2 \gg m^2$* , Nucl. Phys. B **780** (2007) 40 [hep-ph/0703285].
- [38] I. Bierenbaum, J. Blümlein, S. Klein and C. Schneider, *Two-Loop Massive Operator Matrix Elements for Unpolarized Heavy Flavor Production to $O(\epsilon)$* , Nucl. Phys. B **803** (2008) 1 [arXiv:0803.0273 [hep-ph]].
- [39] I. Bierenbaum, J. Blümlein and S. Klein, *The Gluonic Operator Matrix Elements at $O(\alpha_s^2)$ for DIS Heavy Flavor Production*, Phys. Lett. B **672** (2009) 401 [arXiv:0901.0669 [hep-ph]].
- [40] J. Blümlein, A. Hasselhuhn and T. Pfoh, *The $O(\alpha_s^2)$ heavy quark corrections to charged current deep-inelastic scattering at large virtualities*, Nucl. Phys. B **881** (2014) 1 [arXiv:1401.4352 [hep-ph]].
- [41] J. Blümlein, G. Falcioni and A. De Freitas, *The Complete $O(\alpha_s^2)$ Non-Singlet Heavy Flavor Corrections to the Structure Functions $g_{1,2}^{ep}(x, Q^2)$, $F_{1,2,L}^{ep}(x, Q^2)$, $F_{1,2,3}^{v(\bar{v})}(x, Q^2)$ and the Associated Sum Rules*, Nucl. Phys. B **910** (2016) 568 [arXiv:1605.05541 [hep-ph]].
- [42] J. Blümlein, A. De Freitas and C. Schneider, *Higher Order Heavy Quark Corrections to Deep-Inelastic Scattering*, Nucl. Part. Phys. Proc. **261-262** (2015) 185 [arXiv:1411.5669 [hep-ph]].
- [43] W.L. van Neerven and E.B. Zijlstra, *$O(\alpha_s^2)$ contributions to the deep-inelastic Wilson coefficient*, Phys. Lett. B **272** (1991) 127; *Contribution of the second order gluonic Wilson coefficient to the deep-inelastic structure function*, Phys. Lett. B **273** (1991) 476; *$O(\alpha_s^2)$ QCD corrections to the deep-inelastic proton structure functions F_2 and F_L* , Nucl. Phys. B **383** (1992) 525;
 S. Moch and J.A.M. Vermaseren, *Deep-inelastic structure functions at two loops*, Nucl. Phys. B **573** (2000) 853 [hep-ph/9912355].
- [44] J.A.M. Vermaseren, A. Vogt and S. Moch, *The Third-order QCD corrections to deep-inelastic scattering by photon exchange*, Nucl. Phys. B **724** (2005) 3 [hep-ph/0504242].
- [45] M. Buza, Y. Matiounine, J. Smith and W. L. van Neerven, *Charm electroproduction viewed in the variable flavor number scheme versus fixed order perturbation theory*, Eur. Phys. J. C **1** (1998) 301 [hep-ph/9612398].
- [46] E. Laenen, S. Riemersma, J. Smith and W.L. van Neerven, *Complete $O(\alpha_s)$ corrections to heavy flavor structure functions in electroproduction*, Nucl. Phys. B **392** (1993) 162; *$O(\alpha_s)$ corrections to heavy flavor inclusive distributions in electroproduction*, 229;
 S. Riemersma, J. Smith and W. L. van Neerven, *Rates for inclusive deep-inelastic electroproduction of charm quarks at HERA*, Phys. Lett. B **347** (1995) 143 [hep-ph/9411431].

- [47] S.I. Alekhin and J. Blümlein, *Mellin representation for the heavy flavor contributions to deep-inelastic structure functions*, Phys. Lett. B **594** (2004) 299 [hep-ph/0404034].
- [48] P. Nogueira, *Automatic Feynman graph generation*, J. Comput. Phys. **105** (1993) 279.
- [49] J.A.M. Vermaseren, *New features of FORM*, math-ph/0010025.
- [50] T. van Ritbergen, A.N. Schellekens and J.A.M. Vermaseren, Int. J. Mod. Phys. A **14** (1999) 41 [hep-ph/9802376].
- [51] J. Lagrange, *Nouvelles recherches sur la nature et la propagation du son*, Miscellanea Taurinensis, t. II, 1760-61; Oeuvres t. I, p. 263;
 C.F. Gauß, *Theoria attractionis corporum sphaeroidicorum ellipticorum homogeneorum methodo novo tractate*, Commentationes societatis scientiarum Gottingensis recentiores, Vol III, 1813, Werke Bd. V pp. 5-7;
 G. Green, *Essay on the Mathematical Theory of Electricity and Magnetism*, Nottingham, 1828 [Green Papers, pp. 1-115];
 M. Ostrogradski, Mem. Ac. Sci. St. Peters., **6**, (1831) 39;
 K.G. Chetyrkin, A.L. Kataev and F.V. Tkachov, *New Approach To Evaluation Of Multiloop Feynman Integrals: The Gegenbauer Polynomial x -Space Technique*, Nucl. Phys. B **174** (1980) 345;
 S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, Int. J. Mod. Phys. A **15** (2000) 5087 [hep-ph/0102033].
- [52] C. Studerus, *Reduze-Feynman Integral Reduction in C++*, Comput. Phys. Commun. **181** (2010) 1293 [arXiv:0912.2546 [physics.comp-ph]];
 A. von Manteuffel and C. Studerus, *Reduze 2 - Distributed Feynman Integral Reduction*, arXiv:1201.4330 [hep-ph].
- [53] C. W. Bauer, A. Frink and R. Kreckel, *Introduction to the GiNaC framework for symbolic computation within the C++ programming language*, Symbolic Computation **33** (2002) 1, cs/0004015 [cs-sc].
- [54] R.H. Lewis, *Computer Algebra System Fermat*, <http://home.bway.net/lewis>.
- [55] M. Karr, *Summation in Finite Terms*, J. ACM **28** (1981) 305.
- [56] C. Schneider, *Symbolic Summation in Difference Fields* Ph.D. Thesis RISC, Johannes Kepler University, Linz technical report 01-17 (2001).
- [57] C. Schneider, *A Collection of Denominator Bounds to Solve Parameterized Linear Difference Equations in $\Pi\Sigma$ -Extensions*, An. Univ. Timisoara Ser. Mat.-Inform. **42** (2004) 163;
J. Differ. Equations Appl. **11** (2005) 799;
Degree Bounds To Find Polynomial Solutions of Parameterized Linear Difference Equations in $\Pi\Sigma$ -Fields, Appl. Algebra Engrg. Comm. Comput. **16** (2005) 1.
- [58] C. Schneider, *Simplifying Sums in $\Pi\Sigma$ Extensions*, J. Algebra Appl. **6** (2007) 415.
- [59] C. Schneider, *Motives, Quantum Field Theory, and Pseudodifferential Operators (Clay Mathematics Proceedings Vol. 12* ed. A. Carey, D. Ellwood, S. Paycha and S. Rosenberg, (Amer. Math. Soc) (2010), 285 [arXiv:0904.2323].
- [60] C. Schneider, *Parameterized Telescoping Proves Algebraic Independence of Sums*, Ann. Comb. **14** (2010) 533 [arXiv:0808.2596].
- [61] C. Schneider, *Fast Algorithms for Refined Parameterized Telescoping in Difference Fields*, in: Computer Algebra and Polynomials, Applications of Algebra and Number Theory, J. Gutierrez, J. Schicho, M. Weimann (ed.), Lecture Notes in Computer Science (LNCS) 8942 (2015), 157[arXiv:13077887 [cs.SC]].

- [62] C. Schneider, *A Refined Difference Field Theory for Symbolic Summation*, J. Symbolic Comput. **43** (2008) 611 [arXiv:0808.2543v1]; *A Difference Ring Theory for Symbolic Summation*, J. Symb. Comput. **72** (2016) 82 [arXiv:1408.2776 [cs.SC]]; *Summation Theory II: Characterizations of $\Pi\Sigma$ -extensions and algorithmic aspects*, J. Symb. Comput. (2016), in press, [arXiv:1603.04285 [cs.SC]].
- [63] C. Schneider, *Product representations in $\Pi\Sigma$ -fields*, Ann. Comb. **9**(1) (2005) 75;
 S.A. Abramov and M. Petkovšek, *Polynomial ring automorphisms, rational (w, σ) -canonical forms, and the assignment problem*, J. Symbolic Comput., **45**(6) (2010) 684;
 C. Schneider, *Structural theorems for symbolic summation*, Appl. Algebra Engrg. Comm. Comput., **21**(1) (2010) 1;
 C. Schneider, *A streamlined difference ring theory: Indefinite nested sums, the alternating sign and the parameterized telescoping problem*, In: Symbolic and Numeric Algorithms for Scientific Computing (SYNASC), 2014, 15th International Symposium, F. Winkler, V. Negru, T. Ida, T. Jebelean, D. Petcu, S. Watt, D. Zaharie (ed.), (2015) pp. 26; IEEE Computer Society, arXiv:1412.2782v1 [cs.SC].
- [64] C. Schneider, *Symbolic summation assists combinatorics*, Sém. Lothar. Combin. **56** (2007) 1, article B56b.
- [65] C. Schneider, *Simplifying Multiple Sums in Difference Fields*, in: *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions* Texts and Monographs in Symbolic Computation eds. C. Schneider and J. Blümlein (Springer, Wien, 2013) 325 [arXiv:1304.4134 [cs.SC]].
- [66] W.N. Bailey, *Generalized Hypergeometric Series*, (Cambridge University Press, Cambridge, 1935);
 L.J. Slater, *Generalized Hypergeometric Functions*, (Cambridge University Press, Cambridge, 1966);
 P. Appell and J. Kampé de Fériet, *Fonctions Hypergéométriques et Hypersphériques, Polynômes d' Hermite*, (Gauthier-Villars, Paris, 1926);
 P. Appell, *Les Fonctions Hypergéométriques de Plusieurs Variables*, (Gauthier-Villars, Paris, 1925);
 J. Kampé de Fériet, *La fonction hypergéométrique*, (Gauthier-Villars, Paris, 1937);
 H. Exton, *Multiple Hypergeometric Functions and Applications*, (Ellis Horwood, Chichester, 1976);
 H. Exton, *Handbook of Hypergeometric Integrals*, (Ellis Horwood, Chichester, 1978);
 H.M. Srivastava and P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, (Ellis Horwood, Chichester, 1985);
 M.J. Schlosser, in: *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions*, C. Schneider, J. Blümlein, Eds., p. 305, (Springer, Wien, 2013) [arXiv:1305.1966 [math.CA]].
- [67] E.W. Barnes, *A new development of the theory of the hypergeometric functions*, Proc. Lond. Math. Soc. (2) **6** (1908) 141; *A transformation of generalised hypergeometric series*, Quart. Journ. Math. **41** (1910) 136;
 H. Mellin, *Abriß einer einheitlichen Theorie der Gamma- und der hypergeometrischen Funktionen*, Math. Ann. **68** (1910) 305;
 M. Czakon, *Automatized analytic continuation of Mellin-Barnes integrals*, Comput. Phys. Commun. **175** (2006) 559 [hep-ph/0511200];
 A.V. Smirnov and V.A. Smirnov, *On the Resolution of Singularities of Multiple Mellin-Barnes Integrals*, Eur. Phys. J. C **62** (2009) 445 [arXiv:0901.0386 [hep-ph]].
- [68] F.C.S. Brown, *The Massless higher-loop two-point function*, Commun. Math. Phys. **287** (2009) 925 [arXiv:0804.1660 [math.AG]].

- F. Wißbrock, $O(\alpha_s^3)$ Contributions to the Heavy Flavor Wilson Coefficients of the Structure Function $F_2(x, Q^2)$ at $Q^2 \gg m^2$, PhD Thesis, TU Dortmund, 2015;
- E. Panzer, *On hyperlogarithms and Feynman integrals with divergences and many scales*, JHEP **1403** (2014) 071 [arXiv:1401.4361 [hep-th]].
- [69] A.V. Kotikov, *Differential equations method: New technique for massive Feynman diagrams calculation*, Phys. Lett. B **254** (1991) 158;
- E. Remiddi, *Differential equations for Feynman graph amplitudes*, Nuovo Cim. A **110** (1997) 1435 [hep-th/9711188];
- M. Caffo, H. Czyz, S. Laporta and E. Remiddi, *Master equations for master amplitudes*, Acta Phys. Polon. B **29** (1998) 2627 [hep-th/9807119]; *The Master differential equations for the two loop sunrise selfmass amplitudes*, Nuovo Cim. A **111** (1998) 365 [hep-th/9805118];
- T. Gehrmann and E. Remiddi, *Differential equations for two loop four point functions*, Nucl. Phys. B **580** (2000) 485 [hep-ph/9912329].
- [70] G. Almkvist and D. Zeilberger, *The Method of Differentiating under the Integral Sign*, J. Symb. Comp. **10** (1990) 571.
- M. Apagodu and D. Zeilberger, *Multi-variable Zeilberger and Almkvist-Zeilberger algorithms and the sharpening of Wilf-Zeilberger theory*, Adv. Appl. Math. (Special Regev Issue), **37** (2006) 139.
- [71] J. Ablinger, *Computer Algebra Algorithms for Special Functions in Particle Physics*, Ph.D. Thesis, J. Kepler University Linz, 2012, arXiv:1305.0687 [math-ph];
- [72] E. Remiddi and J.A.M. Vermaseren, *Harmonic polylogarithms*, Int. J. Mod. Phys. A **15** (2000) 725 [hep-ph/9905237].
- [73] J. Ablinger, A. Behring, J. Blümlein, A. De Freitas, M. van Hoeij, C.G. Raab, and C. Schneider, DESY 16–147.
- [74] S. Laporta and E. Remiddi, *Analytic treatment of the two loop equal mass sunrise graph*, Nucl. Phys. B **704** (2005) 349 [hep-ph/0406160];
- S. Bloch and P. Vanhove, *The elliptic dilogarithm for the sunset graph*, J. Number Theor. **148** (2015) 328 [arXiv:1309.5865 [hep-th]];
- E. Remiddi and L. Tancredi, *Differential equations and dispersion relations for Feynman amplitudes. The two-loop massive sunrise and the kite integral*. Nucl. Phys. B **907** (2016) 400 [arXiv:1602.01481 [hep-ph]].
- [75] L. Adams, C. Bogner, A. Schweitzer and S. Weinzierl, *The kite integral to all orders in terms of elliptic polylogarithms*, arXiv:1607.01571 [hep-ph].
- [76] L. Adams, C. Bogner and S. Weinzierl, *The iterated structure of the all-order result for the two-loop sunrise integral*, J. Math. Phys. **57** (2016) no.3, 032304 [arXiv:1512.05630 [hep-ph]] and references therein.
- [77] J. Ablinger, J. Blümlein, S. Klein and C. Schneider, *Modern Summation Methods and the Computation of 2- and 3-loop Feynman Diagrams*, Nucl. Phys. Proc. Suppl. **205-206** (2010) 110 [arXiv:1006.4797 [math-ph]];
- J. Blümlein, A. Hasselhuhn and C. Schneider, *Evaluation of Multi-Sums for Large Scale Problems*, PoS (RADCOR 2011) 032 [arXiv:1202.4303 [math-ph]];
- C. Schneider, *Symbolic Summation in Difference Fields and Its Application in Particle Physics*, Computer Algebra Rundbrief **53** (2013), 8;
- C. Schneider, *Modern Summation Methods for Loop Integrals in Quantum Field Theory: The Packages Sigma, EvaluateMultiSums and SumProduction*, J. Phys. Conf. Ser. **523** (2014) 012037 [arXiv:1310.0160 [cs.SC]].

- [78] C. Schneider, *A new Sigma approach to multi-summation*, Advances in Applied Math. **34**(4) (2005) 740;
 J. Ablinger, J. Blümlein, M. Round and C. Schneider, *Advanced Computer Algebra Algorithms for the Expansion of Feynman Integrals*, PoS(LL2012)050, (2012) 14 p. [arXiv:1210.1685 [cs.SC]];
 M. Round et al., in preparation.
- [79] B. Zürcher, *Rationale Normalformen von pseudo-linearen Abbildungen*, Master's thesis, Mathematik, ETH Zürich (1994);
 S. Gerhold, *Uncoupling systems of linear Ore operator equations*, Master's thesis, RISC, J. Kepler University, Linz, 2002;
 C. Schneider, A. De Freitas and J. Blümlein, *Recent Symbolic Summation Methods to Solve Coupled Systems of Differential and Difference Equations*, PoS (LL2014) 017, (2014) [arXiv:1407.2537 [cs.SC]].
- [80] J. Ablinger, *The package HarmonicSums: Computer Algebra and Analytic Aspects of Nested Sums*, PoS (LL2014) 019, (2014); *Inverse Mellin Transforms of Holonomic Sequences*, PoS (LL2016) 067;
A Computer Algebra Toolbox for Harmonic Sums Related to Particle Physics, Diploma Thesis, J. Kepler University Linz, 2009, arXiv:1011.1176 [math-ph].
- [81] S. Alekhin, J. Blümlein and S. Moch, *The ABM parton distributions tuned to LHC data*, Phys. Rev. D **89** (2014) no.5, 054028 [arXiv:1310.3059 [hep-ph]].
- [82] H. Kawamura, N.A. Lo Presti, S. Moch and A. Vogt, *On the next-to-next-to-leading order QCD corrections to heavy-quark production in deep-inelastic scattering*, Nucl. Phys. B **864** (2012) 399 [arXiv:1205.5727 [hep-ph]].
- [83] S. Moch, J.A.M. Vermaseren and A. Vogt, *Third-order QCD corrections to the charged-current structure function F_3* , Nucl. Phys. B **813** (2009) 220 [arXiv:0812.4168 [hep-ph]].
- [84] J. Davies, A. Vogt, S. Moch and J.A.M. Vermaseren, *Non-singlet coefficient functions for charged-current deep-inelastic scattering to the third order in QCD*, arXiv:1606.08907 [hep-ph].
- [85] S. Moch, J.A.M. Vermaseren and A. Vogt, *The Three-Loop Splitting Functions in QCD: The Helicity-Dependent Case*, Nucl. Phys. B **889** (2014) 351 [arXiv:1409.5131 [hep-ph]].
- [86] A. Behring, J. Blümlein, A. De Freitas, A. von Manteuffel and C. Schneider, *The 3-Loop Non-Singlet Heavy Flavor Contributions to the Structure Function $g_1(x, Q^2)$ at Large Momentum Transfer*, Nucl. Phys. B **897** (2015) 612 [arXiv:1504.08217 [hep-ph]].
- [87] A. Behring, J. Blümlein, A. De Freitas, A. Hasselhuhn, A. von Manteuffel and C. Schneider, *$O(\alpha_s^3)$ heavy flavor contributions to the charged current structure function $xF_3(x, Q^2)$ at large momentum transfer*, Phys. Rev. D **92** (2015) 11, 114005 [arXiv:1508.01449 [hep-ph]].
- [88] A. Behring, J. Blümlein, G. Falcioni, A. De Freitas, A. von Manteuffel and C. Schneider, DESY 16-148.
- [89] M. Glück, S. Kretzer and E. Reya, *The Strange sea density and charm production in deep-inelastic charged current processes*, Phys. Lett. B **380** (1996) 171 [Erratum: Phys. Lett. B **405** (1997) 391] [hep-ph/9603304];
 J. Blümlein, A. Hasselhuhn, P. Kovacikova and S. Moch, *$O(\alpha_s)$ Heavy Flavor Corrections to Charged Current Deep-Inelastic Scattering in Mellin Space*, Phys. Lett. B **700** (2011) 294 [arXiv:1104.3449 [hep-ph]];
 M. Buza and W.L. van Neerven, *$O(\alpha_s^2)$ contributions to charm production in charged current deep-inelastic lepton - hadron scattering*, Nucl. Phys. B **500** (1997) 301 [hep-ph/9702242].

- [90] S. Wandzura and F. Wilczek, *Sum Rules for Spin Dependent Electroproduction: Test of Relativistic Constituent Quarks*, Phys. Lett. B **72** (1977) 195.
- [91] J.D. Jackson, G.G. Ross and R.G. Roberts, *Polarized Structure Functions in the Parton Model*, Phys. Lett. B **226** (1989) 159.
- [92] J. Blümlein and N. Kochelev, *On the twist two and twist three contributions to the spin dependent electroweak structure functions*, Nucl. Phys. B **498** (1997) 285 [hep-ph/9612318]; *On the twist-2 contributions to polarized structure functions and new sum rules*, Phys. Lett. B **381** (1996) 296 [hep-ph/9603397].
- [93] J. Blümlein and A. Tkabladze, *Target mass corrections for polarized structure functions and new sum rules*, Nucl. Phys. B **553** (1999) 427 [hep-ph/9812478].
- [94] J. Blümlein and H. Böttcher, *QCD Analysis of Polarized Deep-inelastic Scattering Data*, Nucl. Phys. B **841** (2010) 205 [arXiv:1005.3113 [hep-ph]].
- [95] D. Boer *et al.*, *Gluons and the quark sea at high energies: Distributions, polarization, tomography*, arXiv:1108.1713 [nucl-th];
J. L. Abelleira Fernandez *et al.* [LHeC Study Group Collaboration], *A Large Hadron Electron Collider at CERN: Report on the Physics and Design Concepts for Machine and Detector*, J. Phys. G **39** (2012) 075001 [arXiv:1206.2913 [physics.acc-ph]].
- [96] K.A. Olive *et al.* [Particle Data Group Collaboration], *Review of Particle Physics*, Chin. Phys. C **38** (2014) 090001.
- [97] J. Blümlein and W. L. van Neerven, *Heavy flavor contributions to the deep-inelastic scattering sum rules*, Phys. Lett. B **450** (1999) 417 [hep-ph/9811351].
- [98] S.L. Adler, *Sum rules giving tests of local current commutation relations in high-energy neutrino reactions*, Phys. Rev. **143** (1966) 1144.
- [99] J.D. Bjorken, *Inequality for Backward electron-Nucleon and Muon-Nucleon Scattering at High Momentum Transfer*, Phys. Rev. **163** (1967) 1767.
- [100] J.D. Bjorken, *Inelastic Scattering of Polarized Leptons from Polarized Nucleons*, Phys. Rev. D **1** (1970) 1376.
- [101] D.J. Gross and C.H. Llewellyn Smith, *High-energy neutrino - nucleon scattering, current algebra and partons*, Nucl. Phys. B **14** (1969) 337.
- [102] J. Ablinger *et al.*, DESY 14–019.
- [103] J. Ablinger, J. Blümlein, S. Klein, C. Schneider and F. Wißbrock, *3-Loop Heavy Flavor Corrections to DIS with two Massive Fermion Lines*, arXiv:1106.5937 [hep-ph];
J. Ablinger, J. Blümlein, A. Hasselhuhn, S. Klein, C. Schneider and F. Wißbrock, *New Heavy Flavor Contributions to the DIS Structure Function $F_2(x, Q^2)$ at $O(\alpha_s^3)$* , PoS (RADCOR2011) 031 [arXiv:1202.2700 [hep-ph]].
- [104] R. Harlander, T. Seidensticker and M. Steinhauser, *Complete corrections of $O(\alpha_s)$ to the decay of the Z boson into bottom quarks*, Phys. Lett. B **426** (1998) 125 [hep-ph/9712228];
T. Seidensticker, *Automatic application of successive asymptotic expansions of Feynman diagrams*, hep-ph/9905298.