

Leptogenesis: Improving predictions for experimental searches

Marco Drewes*

Physik Department T70, Technische Universität München, James Franck Straße 1, D-85748 Garching, Germany

E-mail: marco.drewes@tum.de

Björn Garbrecht

Physik Department T70, Technische Universität München, James Franck Straße 1, D-85748 Garching, Germany

Dario Gueter

Physik Department T70, Technische Universität München, James Franck Straße 1, D-85748 Garching, Germany

Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany

Excellence Cluster Universe, Boltzmannstraße 2, Technische Universität München, 85748 Garching, Germany

Juraj Klarić

Physik Department T70, Technische Universität München, James Franck Straße 1, D-85748 Garching, Germany

Heavy right handed neutrinos could not only explain the observed neutrino masses via the see-saw mechanism, but also generate the baryon asymmetry of the universe via leptogenesis due to their CP-violating interactions in the early universe. We review recent progress in the theoretical description of this nonequilibrium process. Improved calculations are particularly important for a comparison with experimental data in testable scenarios with Majorana masses below the TeV scale, in which the heavy neutrinos can be found at the LHC, in the NA62 experiment, at T2K or in future experiments, including SHiP, DUNE and experiments at the FCC, ILC or CEPC. In addition, the relevant source of CP-violation may be experimentally accessible, and the heavy neutrinos can give a sizable contribution to neutrinoless double β decay. In these low scale leptogenesis scenarios, the matter-antimatter asymmetry is generated at temperatures when the heavy neutrinos are relativistic, and thermal corrections to the transport equations in the early universe are large.

38th International Conference on High Energy Physics

3-10 August 2016

Chicago, USA

*Speaker.

Introduction - There is compelling evidence that the observable universe does not contain significant amounts of antimatter, and that the (baryonic) matter is the remnant of a tiny matter-antimatter asymmetry in the primordial plasma that survived after mutual annihilation of all particles and antiparticles, see e.g. Ref. [1] for a review. This *baryon asymmetry of the universe* (BAU) is commonly expressed in terms of the net baryon-to-photon ratio $\eta_B = n_B/n_\gamma \simeq 6 \times 10^{-10}$. It cannot be explained within the Standard Model (SM) of particle physics and has to be generated dynamically via *baryogenesis* if the radiation dominated epoch started with $\eta_B = 0$, as suggested by many models of cosmic inflation. A particularly economic solution to the baryogenesis problem that relates η_B to the observed neutrino masses and mixings is given by *leptogenesis* [2]. Consider the minimal extension of the SM Lagrangian \mathcal{L}_{SM} by n heavy Majorana neutrinos N_i ,

$$\mathcal{L} = \mathcal{L}_{SM} + \frac{1}{2} \bar{N}_i (i \not{\partial} - M_{ij}) N_j - Y_{ia}^* \bar{\ell}_a \varepsilon \phi P_R N_i - Y_{ia} \bar{N}_i P_L \phi^\dagger \varepsilon^\dagger \ell_a, \quad (1)$$

with $N_i = N_i^c$, where the superscript c denotes a charge conjugation. The N_i interact with the SM only via their Yukawa interactions Y_{ia} with the SM lepton doublets ℓ_a ($a = e, \mu, \tau$) and the Higgs field ϕ . Here ε is the antisymmetric SU(2)-invariant tensor with $\varepsilon^{12} = 1$. In the (type I) *seesaw model* (1), the same particles N_i that generate the light neutrino mass matrix

$$m_\nu = -v^2 Y^* M^{-1} Y^\dagger = -\theta M \theta^T \quad (2)$$

via the seesaw mechanism [3] can produce a lepton asymmetry via their CP violating interactions in the early universe, which is transferred into a BAU via electroweak sphaleron processes [4]. Here $\theta_{ai} = v Y_{ai}^\dagger M_i^{-1}$ (with $a = e, \mu, \tau$) are the active-sterile neutrino mixing angles at low energies, and $v = 174$ GeV is the Higgs vev. Various different authors have studied different implementations of this idea, which are e.g. summarised in the reviews [5–7] and references therein. Here we only summarise recent developments in the field and give references for further reading.

One can qualitatively distinguish two ways to generate the BAU in the minimal model (1), either in the decay of the right handed neutrinos (*freeze out scenario* or “thermal leptogenesis”) [2], or in CP violating oscillations during their production (*freeze in scenario* or “baryogenesis from neutrino oscillations”) [8, 9]. Which one of them is realised depends on the masses M_i of the N_i . For $M_i \gg v$, the right handed neutrinos come into equilibrium, freeze out and decay long before electroweak sphalerons freeze out at $T \simeq 130$ GeV. In this case the final asymmetry η_B that can be observed today is usually created in the decay of the lightest right handed neutrino. This scenario generically requires $M_i > 10^9$ GeV [10] unless the N_i mass spectrum is highly degenerate [13], leading to *resonant leptogenesis* [14]. For $M_i < v$, the seesaw relation (2) implies that at least some of the Y_{ai} are much smaller than the electron Yukawa coupling. In this case a $\eta_B \neq 0$ freezes in because the heavy neutrinos do not reach thermal equilibrium before sphaleron freezeout.

Leptogenesis is usually described in terms of momentum averaged semiclassical Boltzmann equations of the form

$$xH \frac{dY_N}{dx} = -\Gamma_N (Y_N - Y_N^{\text{eq}}) \quad (3)$$

$$xH \frac{dY_{B-L}}{dx} = (\varepsilon \Gamma_N^D + \varepsilon' \Gamma_N^S) (Y_N - Y_N^{\text{eq}}) - c_W \Gamma_N Y_{B-L}. \quad (4)$$

Here Y_N is the abundance of heavy neutrinos and Y_{B-L} the total $B-L$ charge (both normalised to the cosmic entropy density), H is the Hubble rate, $\Gamma_N = \Gamma_N^D + \Gamma_N^S$ the thermal production rate of heavy neutrinos, including contributions from decays and inverse decays (Γ_N^D) as well as scatterings (Γ_N^S), and $c_W \Gamma_N$ is the washout rate. The parameters ε and ε' measure the difference between the rates for processes involving leptons and antileptons, normalised to their sum, and c_W is a numerical factor that is governed by the ratios of the number densities of heavy neutrinos and SM leptons, which is of order unity if all particles are near thermal equilibrium. We use $x = M/T$ as time variable, where T is the temperature and M an appropriately chosen mass scale (usually the mass of the lightest heavy neutrino). This description has been refined in various ways in recent years.

First principles derivation of kinetic equations - The generation of a matter-antimatter asymmetry in leptogenesis is a pure quantum effect, as it relies on a CP violating quantum interference. It is not obvious that semiclassical Boltzmann equations, which are commonly used in cosmology, are suitable to describe this process in the dense primordial plasma quantitatively [15]. This has motivated efforts to study leptogenesis from first principles of quantum field theory [16–19], in particular in the resonant case [20–26] and in the context of flavour effects [26, 27] and in the freeze in scenario [28, 29]. When all coupling constants are perturbative, then the first principles treatment at leading order reproduces the density matrix formalism [9, 30] commonly used in neutrino physics, assuming that the necessary resummations of all thermal corrections to the dispersion relations and rates are taken into account. The first principles approach provides a systematic way to calculate the coefficients in these equations that is inherently free of double counting problems, and in principle it allows to compute corrections to the leading order result (though this is practically challenging).

Momentum averaging - The effect of the momentum averaging has been studied systematically in Refs. [31, 32]. It was found that the averaged equations are accurate up to corrections between $\sim 10\%$ (for $M > T$) and order one (for $M < T$). In the non-relativistic regime $M > T$, it has been shown that it is convenient to use the moments of the N_i distribution function as dynamical variables [33] (instead of the occupation numbers for individual momentum modes). Compared to tracking individual momentum modes, one can achieve the same accuracy with much less computational effort.

Quasiparticle dispersion relations - The dispersion relations of (quasi)particles in a hot plasma can considerably differ from those in vacuum. In the context of leptogenesis, this issue has been investigated in Ref. [34] and, in more detail, in Refs. [18, 24, 35–39]. In the freeze out scenario with strong washout, thermal corrections are small. In the weak washout regime with $T > M$ and in the freeze in scenario, where the BAU is generated at $T \gg M$, they have to be taken into account [9, 28, 29, 40–45]. At tree level, the larger “thermal mass” of the Higgs boson implies that there exist three qualitatively different temperature regimes: a low temperature regime where the decay $N \rightarrow \phi \ell$ is kinematically allowed, a high temperature regime where the decay $\phi \rightarrow N \ell$ is allowed, and an intermediate regime in which no decay is allowed [34, 46] and Γ_N vanishes. However, a consistent resummation of all thermal contributions shows that the inclusion of scatterings lead to a smooth dependence of Γ_N on T , and there is no regime with $\Gamma_N = 0$ [47]¹, see next paragraph.

¹This effect may also play an important role in the context of perturbative cosmic reheating [48].

Flavour effects - If the couplings Y_{ia} are of order unity, the seesaw relation (2) suggests $M_i > 10^{14}$ GeV, and the BAU is generated in the decay of N_i at temperatures $T > 10^{12}$ GeV (freezeout scenario), at which the SM flavours are indistinguishable ("vanilla leptogenesis"). Then η_B is independent of the phases in the light neutrino mixing matrix. For smaller values of M_i and $T < 10^{12}$ GeV, the charged lepton Yukawa couplings affect the evolution of leptonic asymmetries, and flavour effects are relevant [49–54]. In this case (3,4) should be replaced by matrix valued generalisations [30], which have e.g. been derived in Refs. [9, 26–28, 39, 43, 55, 56]. Flavour effects introduce a dependence of η_B on the phases in the light neutrino mixing matrix and reduce the lower bound on non-degenerate M_i for leptogenesis in the freezeout scenario to $M_i > 10^6$ GeV [11, 12]. The intermediate regime between the vanilla scenario and the fully flavoured scenario has been studied in detail in Ref. [18]. If one wants to achieve successful leptogenesis with even smaller M_i , this requires a mass degeneracy to ensure a resonant enhancement of η_B . Moreover, sizeable Y_{ia} can only be made consistent with small neutrino masses if there are cancellations in (2). Both can be achieved naturally in models with an approximate conservation of $B - L$, where leptogenesis relies mostly on flavour effects [12, 57]. In the freeze in scenario, the asymmetry is always generated at $T \gg M_i$, where the total lepton number is negligible [8, 9], and baryogenesis relies entirely on flavour effects.

Thermal production and damping rates - The rate Γ_N at $T < M$ is in good approximation given by the N_i vacuum decay rate [47, 58–62]. For $T > M$, on the other hand, thermal effects can dominate. These include quantum statistical factors, modifications of the (quasi)particle dispersion relations in the plasma and the fact that the N_i can be produced in scatterings in the dense plasma. For $T \gg M$, there is a collinear enhancement of multiple scatterings, and the Landau-Pomeranchuk-Migdal effect should be taken into account [47, 62–64]. The dominant contribution to Γ_N in this regime comes from logarithmically enhanced t-channel scatterings [61, 65].² In Ref. [66, 67], this calculation has been extended to temperatures across and below the electroweak scale. Thermal corrections do not only affect Γ_N , but also the washout rate $c_W \Gamma_N$ [62, 69] and its flavoured generalisations [61, 67, 70]. In the freeze in scenario, a consistent analysis also requires to include the temperature dependence of the sphaleron rate across the electroweak crossover [71, 72].

Spectator effects - The effect of spectator processes [73] that do not change the total asymmetry, but reshuffle it between different degrees of freedom, has recently been revisited in both, the freeze out [74] and freeze in [28, 29, 45, 75] scenarios. In the latter the spectators can have a significant effect on the viable parameter space.

CP violating parameter - Thermal corrections to the CP violating parameter ε have been studied by several authors [21, 24, 26, 27, 39, 76–78]. A question of particular interest is the dependence of η_B on the N_i mass spectrum in the regime where the splitting ΔM of two M_i is much smaller than their average mass \bar{M} , leading to a resonant enhancement that allows to generate the observed η_B with \bar{M} below the TeV scale [14]. In the resonant limit $\Delta M/\bar{M} \rightarrow 0$, η_B is regularised by the finite thermal width Γ_N . The precise form of the "regulator" that determines the maximal η_B has been the subject of a long standing dispute [14, 79–83]. The issue has been studied from first

²These calculations have been performed in the minimal model (1). In theories with an extended Higgs sector, there can be additional contributions from decays of the heavy scalars, see e.g. [68] and references therein.

principles in Refs. [21, 24, 26, 26, 39, 84], but disagreement on how to count (and not double count) contributions from "mixing" and "oscillations" of the N_i remains, leading to factors ~ 2 uncertainty in these contributions. Note that, while parameter relevant for the enhancement of the source for the total lepton asymmetry is $\bar{M}^2/\Delta M^2$, the flavoured source is enhanced by $T^2/\Delta M^2$, which allows to generate the observed η_B with M_i below the electroweak scale without a mass degeneracy [85]. The CP violating phases in the Y_{ia} can be experimentally constrained from light neutrino oscillations at DUNE and NOvA, measurements of the mixings $U_{ai}^2 = |\theta_{ai}|^2$ in direct search experiments [29, 40, 45, 86] and in the CP violation in N_i decays [87].³

Low scale leptogenesis - The experimental perspectives for a discovery of the N_i are most promising if they have masses M_i below that of the W boson. A discussion of the experimental constraints and perspectives can e.g. be found in the reviews [6, 7, 29, 89, 90, 90–95] and references therein. In this mass range, leptogenesis in the minimal model (1) can only be achieved in the freeze in scenario.⁴ Since the generation of the BAU happens at $T \gg M_i$, relies on flavour effects and may continue across the electroweak transition, all effects discussed above are potentially relevant in this scenario. The viable parameter space has been studied in the models with two [28, 29, 41–45, 75, 97, 98] and three [44, 75, 85, 98, 99] heavy neutrinos. A convenient way to illustrate the results is given by the projection in the $\bar{M} - U_a^2$ plane, where $U_a^2 = \sum_i |\theta_{ai}|^2$. As an example, we show the $\bar{M} - U_\mu^2$ in Fig. 1. If any heavy neutral leptons are discovered in the future, an independent measurement of all U_{ai}^2 , which could be done at the SHiP experiment or a future lepton collider, would provide a powerful test to assess whether these particles are the origin of neutrino masses and matter in the universe in the minimal seesaw model, see Fig. 2. Together with an observation of neutrinoless double β decay and CP violation in light neutrino oscillations, this would impose significant constraints on all model parameters [29, 44, 45, 109].

References

- [1] L. Canetti, M. Drewes and M. Shaposhnikov, *New J. Phys.* **14** (2012) 095012.
- [2] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174** (1986) 45.
- [3] P. Minkowski, *Phys. Lett.* **67B** (1977) 421. M. Gell-Mann, P. Ramond and R. Slansky, *Conf. Proc. C* **790927** (1979) 315. R. N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44** (1980) 912. T. Yanagida, *Prog. Theor. Phys.* **64** (1980) 1103. J. Schechter and J. W. F. Valle, *Phys. Rev. D* **22** (1980) 2227. J. Schechter and J. W. F. Valle, *Phys. Rev. D* **25** (1982) 774.
- [4] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, *Phys. Lett.* **155B** (1985) 36. doi:10.1016/0370-2693(85)91028-7
- [5] W. Buchmuller, R. D. Peccei and T. Yanagida, *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 311. S. Davidson, E. Nardi and Y. Nir, *Phys. Rept.* **466** (2008) 105. S. Blanchet and P. Di Bari, *New J. Phys.* **14** (2012) 125012. C. S. Fong, E. Nardi and A. Riotto, *Adv. High Energy Phys.* **2012** (2012) 158303.
- [6] A. Boyarskiy, O. Ruchayskiy and M. Shaposhnikov, *Ann. Rev. Nucl. Part. Sci.* **59** (2009) 191.

³See e.g. Ref. [88] and references therein for a recent discussion of the relation between η_B and observable CP violation in the lepton sector.

⁴Note that this scenario may also work for larger masses [96], so that there is an overlap between the M_i ranges where the freeze out and freeze in scenario are feasible.




Figure 1: Viable leptogenesis parameter space for inverted hierarchy and $n = 2$ (between the blue lines) and combined constraints from past experiments (grey area), both as found in Ref. [29]. Here $U^2 = \sum_a U_a^2$. We compare this to the expected sensitivity of future searches. The purple lines show the 90% c.l. upper limits for SHiP assuming 0.1 background events and with 2×10^{20} proton target collisions for a ratio of $U_e^2 : U_\mu^2 : U_\tau^2 \sim 52 : 1 : 1$ [100, 101]. The light blue line marks the sensitivity of a detector length of 30m placed near the DUNE/LBNE with an exposure of 5×10^{21} protons on target [102]. The solid olive green line corresponds to the expected reach of FCC-ee for 10^{12} Z bosons with a displaced vertex at 10 – 100 cm [103]. The expected sensitivity of FCC-ee at 2σ for heavy neutrino searches via displaced vertices is presented as dashed olive green line. The brown and dark orange lines show the sensitivity of CEPC and ILC, each at the Z pole run for a center of mass energy $m_{\text{cms}} = m_Z$ [104]. It is important to point out that the ILC can potentially do better at higher centre of mass energies [104]. The turquoise line is the expected limit of the NA62 experiment on U_μ^2 with 2×10^{18} 400 GeV protons on target [105]. The red line is the T2K sensitivity, as estimated in Ref. [106] for 10^{21} protons on target at 90% c.l. with full volume for both the $K^+ \rightarrow \mu^+ N \rightarrow \mu^+ \mu^- \pi^+$ two-body decays (red, solid) and the $K^+ \rightarrow \mu^+ N \rightarrow \mu^+ \mu^- e^+ \nu_e$ three-body decays (red, dashed) [106]. Limits on U_μ^2 can also be obtained from LNV decays of 5×10^{10} B^+ mesons at Belle II (blue line) and 2×10^8 W bosons (yellow line) at the FCC-ee [107]. The expected sensitivity at the FCC-ee from B meson decay is given by the light orange line [107]. The violet lines represent the limits on U_μ^2 from displaced lepton jet (solid) and prompt trilepton (dashed) searches for $\sqrt{s} = 13 \text{ TeV}$ and 300 fb^{-1} at the LHC [108].




Figure 2: Constraints on U_a^2/U^2 from neutrino oscillation data, where $U_a^2 = \sum_i U_{ai}^2$ and $U^2 = \sum_a U_a^2$ [29]. Values inside the black line are consistent with neutrino oscillation data for normal hierarchy (left) and inverted hierarchy (right) of light neutrino masses. The dashed lines correspond to constant U_τ^2 , the light red region is unphysical, as it would require $U_\tau^2 < 0$. The coloured regions indicate the maximally allowed U^2 for given U_a^2/U^2 if one requires that the observed η_B can be generated by leptogenesis with $\bar{M} = 1 \text{ GeV}$.

- [7] M. Drewes, *Int. J. Mod. Phys. E* **22** (2013) 1330019 doi:10.1142/S0218301313300191.
- [8] E. K. Akhmedov, V. A. Rubakov and A. Y. Smirnov, *Phys. Rev. Lett.* **81** (1998) 1359.
- [9] T. Asaka and M. Shaposhnikov, *Phys. Lett. B* **620** (2005) 17.
- [10] S. Davidson and A. Ibarra, *Phys. Lett. B* **535** (2002) 25.
- [11] S. Antusch, S. Blanchet, M. Blennow and E. Fernandez-Martinez, *JHEP* **1001** (2010) 017.
- [12] J. Racker, M. Pena and N. Rius, *JCAP* **1207** (2012) 030.
- [13] L. Covi, E. Roulet and F. Vissani, *Phys. Lett. B* **384** (1996) 169.
- [14] A. Pilaftsis and T. E. J. Underwood, *Nucl. Phys. B* **692** (2004) 303.
- [15] W. Buchmuller and S. Fredenhagen, *Phys. Lett. B* **483** (2000) 217.
- [16] A. De Simone and A. Riotto, *JCAP* **0708** (2007) 002.
- [17] A. Anisimov, W. Buchmuller, M. Drewes and S. Mendizabal, *Phys. Rev. Lett.* **104** (2010) 121102. A. Anisimov, W. Buchmüller, M. Drewes and S. Mendizabal, *Annals Phys.* **326** (2011) 1998.
- [18] M. Beneke, B. Garbrecht, C. Fidler, M. Herranen and P. Schwaller, *Nucl. Phys. B* **843** (2011) 177.
- [19] T. Frossard, M. Garny, A. Hohenegger, A. Kartavtsev and D. Mitrouskas, *Phys. Rev. D* **87** (2013) no.8, 085009.
- [20] A. De Simone and A. Riotto, *JCAP* **0708** (2007) 013.
- [21] B. Garbrecht and M. Herranen, *Nucl. Phys. B* **861** (2012) 17.
- [22] M. Garny, A. Hohenegger, A. Kartavtsev and M. Lindner, *Phys. Rev. D* **80** (2009) 125027.
- [23] M. Garny, A. Hohenegger, A. Kartavtsev and M. Lindner, *Phys. Rev. D* **81** (2010) 085027.
- [24] M. Garny, A. Kartavtsev and A. Hohenegger, *Annals Phys.* **328** (2013) 26.
- [25] S. Iso, K. Shimada and M. Yamanaka, *JHEP* **1404** (2014) 062.
- [26] P. S. Bhupal Dev, P. Millington, A. Pilaftsis and D. Teresi, *Nucl. Phys. B* **891** (2015) 128.
- [27] M. Beneke, B. Garbrecht, M. Herranen and P. Schwaller, *Nucl. Phys. B* **838** (2010) 1.
- [28] M. Drewes, B. Garbrecht, D. Gueter and J. Klaric, arXiv:1606.06690 [hep-ph].
- [29] M. Drewes, B. Garbrecht, D. Gueter and J. Klaric, arXiv:1609.09069 [hep-ph].
- [30] G. Sigl and G. Raffelt, *Nucl. Phys. B* **406** (1993) 423.
- [31] F. Hahn-Woernle, M. Plumacher and Y. Y. Y. Wong, *JCAP* **0908** (2009) 028.
- [32] T. Asaka, S. Eijima and H. Ishida, *JCAP* **1202** (2012) 021.
- [33] D. Bödeker and M. Wörmann, *JCAP* **1402** (2014) 016.
- [34] G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, *Nucl. Phys. B* **685** (2004) 89.
- [35] C. Kiessig and M. Plumacher, *JCAP* **1207** (2012) 014.
- [36] C. Kiessig and M. Plumacher, *JCAP* **1209** (2012) 012.
- [37] A. Hohenegger and A. Kartavtsev, *JHEP* **1407** (2014) 130.
- [38] K. Miura, Y. Hidaka, D. Satow and T. Kunihiro, *Phys. Rev. D* **88** (2013) no.6, 065024.
- [39] P. S. Bhupal Dev, P. Millington, A. Pilaftsis and D. Teresi, *Nucl. Phys. B* **886** (2014) 569.
- [40] M. Shaposhnikov, *JHEP* **0808** (2008) 008.
- [41] L. Canetti and M. Shaposhnikov, *JCAP* **1009** (2010) 001.
- [42] L. Canetti, M. Drewes and M. Shaposhnikov, *Phys. Rev. Lett.* **110** (2013) no.6, 061801.
- [43] L. Canetti, M. Drewes, T. Frossard and M. Shaposhnikov, *Phys. Rev. D* **87** (2013) 093006.
- [44] M. Drewes and S. Eijima, *Phys. Lett. B* **763** (2016) 72.
- [45] P. Hernández, M. Kekic, J. López-Pavón, J. Racker and J. Salvado, *JHEP* **1608** (2016) 157.

- [46] T. Hambye and D. Teresi, Phys. Rev. Lett. **117** (2016) no.9, 091801.
- [47] A. Anisimov, D. Besak and D. Bodeker, JCAP **1103** (2011) 042.
- [48] M. Drewes and J. U. Kang, Nucl. Phys. B **875** (2013) 315 Erratum: [Nucl. Phys. B **888** (2014) 284].
- [49] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, Nucl. Phys. B **575** (2000) 61.
- [50] S. Blanchet and P. Di Bari, JCAP **0703** (2007) 018.
- [51] A. Abada, S. Davidson, A. Ibarra, F.-X. Josse-Michaux, M. Losada and A. Riotto, JHEP **0609** (2006) 010.
- [52] A. Abada, S. Davidson, F. X. Josse-Michaux, M. Losada and A. Riotto, JCAP **0604** (2006) 004.
- [53] E. Nardi, Y. Nir, E. Roulet and J. Racker, JHEP **0601** (2006) 164.
- [54] F. X. Josse-Michaux and A. Abada, JCAP **0710** (2007) 009.
- [55] V. Cirigliano, C. Lee, M. J. Ramsey-Musolf and S. Tulin, Phys. Rev. D **81** (2010) 103503.
- [56] C. Fidler, M. Herranen, K. Kainulainen and P. M. Rakhila, JHEP **1202** (2012) 065.
- [57] D. Aristizabal Sierra, L. A. Munoz and E. Nardi, Phys. Rev. D **80** (2009) 016007.
- [58] A. Salvio, P. Lodone and A. Strumia, JHEP **1108** (2011) 116.
- [59] M. Laine and Y. Schroder, JHEP **1202** (2012) 068.
- [60] S. Biondini, N. Brambilla, M. A. Escobedo and A. Vairo, JHEP **1312** (2013) 028.
- [61] B. Garbrecht, F. Glowna and P. Schwaller, Nucl. Phys. B **877** (2013) 1.
- [62] I. Ghisoiu and M. Laine, JCAP **1412** (2014) no.12, 032.
- [63] D. Besak and D. Bodeker, JHEP **1005** (2010) 007.
- [64] D. Besak and D. Bodeker, JCAP **1203** (2012) 029.
- [65] B. Garbrecht, F. Glowna and M. Herranen, JHEP **1304** (2013) 099.
- [66] J. Ghiglieri and M. Laine, JHEP **1511** (2015) 171.
- [67] J. Ghiglieri and M. Laine, JCAP **1607** (2016) no.07, 015.
- [68] M. Drewes and J. U. Kang, JHEP **1605** (2016) 051.
- [69] D. Bodeker and M. Laine, JCAP **1405** (2014) 041.
- [70] B. Garbrecht, F. Gautier and J. Klaric, JCAP **1409** (2014) no.09, 033.
- [71] M. D'Onofrio, K. Rummukainen and A. Tranberg, JHEP **1208** (2012) 123.
- [72] M. D'Onofrio, K. Rummukainen and A. Tranberg, Phys. Rev. Lett. **113** (2014) no.14, 141602.
- [73] W. Buchmuller and M. Plumacher, Phys. Lett. B **511** (2001) 74.
- [74] B. Garbrecht and P. Schwaller, JCAP **1410** (2014) no.10, 012.
- [75] B. Shuve and I. Yavin, Phys. Rev. D **89** (2014) no.7, 075014.
- [76] M. Garny, A. Hohenegger and A. Kartavtsev, Phys. Rev. D **81** (2010) 085028.
- [77] S. Biondini, N. Brambilla, M. A. Escobedo and A. Vairo, JHEP **1603** (2016) 191, Erratum: [JHEP **1608** (2016) 072].
- [78] S. Biondini, N. Brambilla and A. Vairo, JHEP **1609** (2016) 126.
- [79] J. Liu and G. Segre, Phys. Rev. D **49** (1994) 1342.
- [80] L. Covi and E. Roulet, Phys. Lett. B **399** (1997) 113.
- [81] M. Flanz, E. A. Paschos, U. Sarkar and J. Weiss, Phys. Lett. B **389** (1996) 693.
- [82] A. Pilaftsis, Nucl. Phys. B **504** (1997) 61.
- [83] W. Buchmuller and M. Plumacher, Phys. Lett. B **431** (1998) 354.
- [84] A. Kartavtsev, P. Millington and H. Vogel, JHEP **1606** (2016) 066.

- [85] M. Drewes and B. Garbrecht, *JHEP* **1303** (2013) 096.
- [86] T. Asaka, S. Eijima and H. Ishida, *JHEP* **1104** (2011) 011. O. Ruchayskiy and A. Ivashko, *JHEP* **1206** (2012) 100.
- [87] G. Cvetič, C. S. Kim and J. Zamora-Saa, *Phys. Rev. D* **89** (2014) no.9, 093012.
- [88] M. Drewes, B. Garbrecht, D. Gueter and J. Klaric, arXiv:1611.04769 [hep-ph].
- [89] A. Atre, T. Han, S. Pascoli and B. Zhang, *JHEP* **0905** (2009) 030.
- [90] O. Ruchayskiy and A. Ivashko, *JHEP* **1206** (2012) 100.
- [91] A. Kusenko, *Phys. Rept.* **481** (2009) 1.
- [92] S. Alekhin *et al.*, *Rept. Prog. Phys.* **79** (2016) no.12, 124201.
- [93] M. Drewes and B. Garbrecht, arXiv:1502.00477 [hep-ph].
- [94] F. F. Deppisch, P. S. Bhupal Dev and A. Pilaftsis, *New J. Phys.* **17** (2015) no.7, 075019.
- [95] R. Adhikari *et al.*, [arXiv:1602.04816 [hep-ph]].
- [96] B. Garbrecht, *Phys. Rev. D* **90** (2014) no.6, 063522.
- [97] A. Abada, G. Arcadi, V. Domcke and M. Lucente, *JCAP* **1511** (2015) no.11, 041.
- [98] P. Hernandez, M. Kekic, J. Lopez-Pavon, J. Racker and N. Rius, *JHEP* **1510** (2015) 067.
- [99] L. Canetti, M. Drewes and B. Garbrecht, *Phys. Rev. D* **90** (2014) no.12, 125005.
- [100] M. Anelli *et al.* [SHiP Collaboration], arXiv:1504.04956 [physics.ins-det].
- [101] E. Graverini [SHiP Collaboration], CERN-SHiP-NOTE-2016-003.
- [102] C. Adams *et al.* [LBNE Collaboration], arXiv:1307.7335 [hep-ex].
- [103] A. Blondel *et al.* [FCC-ee study Team], arXiv:1411.5230 [hep-ex].
- [104] S. Antusch, E. Cazzato and O. Fischer, *JHEP* **1612** (2016) 007.
- [105] T. Spadaro, Perspectives from the NA62 experiment, Talk given at the PBC Kickoff Meeting - CERN (September 2016).
- [106] T. Asaka, S. Eijima and A. Watanabe, *JHEP* **1303** (2013) 125.
- [107] T. Asaka and H. Ishida, *Phys. Lett. B* **763** (2016) 393.
- [108] E. Izaguirre and B. Shuve, *Phys. Rev. D* **91** (2015) no.9, 093010.
- [109] T. Asaka, S. Eijima and H. Ishida, *Phys. Lett. B* **762** (2016) 371.