

Four-jet production in single- and double-parton scattering within high-energy factorization

Krzysztof Kutak

*Institute of Nuclear Physics, Polish Academy of Sciences,
Radzikowskiego 152, 31-342 Kraków, Poland
E-mail: Krzysztof.Kutak@ifj.edu.pl*

We report on a first study of 4-jet production in a complete high-energy factorization (HEF) framework [1, 2]. We include and discuss contributions from both single-parton scattering (SPS) and double-parton scattering (DPS) and compare to the measured data. The DPS HEF result is considerably smaller than the one obtained with collinear factorization. The mechanism leading to this difference is of kinematical nature. In contrast to the collinear approach, the HEF approach nicely describes the distribution of the ΔS variable, which involves all four jets and their angular correlations.

*38th International Conference on High Energy Physics
3-10 August 2016
Chicago, USA*

1. Introduction

So far, complete ($n \geq 4$)-jet production via single-parton scattering (SPS) was discussed only within collinear factorization. Results up to next-to-leading (NLO) precision can be found in [4, 5]. Here we report on recent study of production of four jets including SPS processes and Double-parton scattering (DPS) within high-energy (k_T -)factorization (HEF) [3]. Double-parton scattering (DPS) was claimed to have been observed for the first time at the Tevatron [6]. In the LHC era, with much higher collision energies available, the field has received a new impulse and several experimental and theoretical studies address the problem of pinning down DPS effects. Even just from purely theoretical point of view, the problem is quite subtle. As for the non perturbative side, it is in principle necessary, when considering a double-parton scattering, to take into account the correlations between the two partons coming from the same protons and involved in the scattering processes. Such an information should be encoded in a set of double parton distribution functions (DPDFs), generalising usual parton distribution functions (PDFs). Some successful attempts to generalize the usual evolution and to have relevance for phenomenology are becoming to appear only recently [8, 9, 10, 11].

In the meanwhile, phenomenological and experimental studies of double-parton scattering rely on factorized Ansatz for the DPDFs, which amount to neglecting longitudinal momentum correlations between partons and treating transversal ones by introducing an effective cross section, σ_{eff} . The latter quantity is usually extracted from experimental data. In the present approach we will use the factorized Ansatz and concentrate on the difference between leading-order collinear and high-energy-factorization results. The latter includes effectively higher-order corrections. For most of high-energy reactions the single-parton scattering dominates over the double-parton scattering. The extraordinary example is double production of $c\bar{c}$ pairs [7]. For four-jet production, disentangling the ordinary SPS contributions from the DPS corrections can be quite challenging for several reasons: first of all, it is necessary to define sufficiently sensitive, process-dependent observables, w.r.t. which the DPS differential cross section manifestly dominates at least in some corners of phase space.

2. Single-parton scattering production of four jets

The HEF factorization formula for the calculation of the inclusive partonic 4-jet cross section at the Born level reads

$$\begin{aligned} \sigma_{4-jets}^B &= \sum_{i,j} \int \frac{dx_1}{x_1} \frac{dx_2}{x_2} d^2k_{T1} d^2k_{T2} \mathcal{F}_i(x_1, k_{T1}, \mu_F) \mathcal{F}_j(x_2, k_{T2}, \mu_F) \\ &\times \frac{1}{2\hat{s}} \prod_{l=1}^4 \frac{d^3k_l}{(2\pi)^3 2E_l} \Theta_{4-jet} (2\pi)^4 \delta \left(x_1 P_1 + x_2 P_2 + \vec{k}_{T1} + \vec{k}_{T2} - \sum_{l=1}^4 k_l \right) \overline{|\mathcal{M}(i^*, j^* \rightarrow 4 \text{ part.})|^2}. \end{aligned} \quad (2.1)$$

Here $\mathcal{F}_i(x_k, k_{Tk}, \mu_F)$ is a transverse momentum dependent (TMD) distribution function for a given type of parton carrying $x_{1,2}$ momentum fractions of the proton and evaluated at the factorization scale μ_F . The index l runs over the four partons in the final state, the partonic center of mass energy squared is $\hat{s} = 2x_1x_2 P_i \cdot P_j$; the function Θ_{4-jet} takes into account the kinematic cuts applied

and \mathcal{M} is the partonic on-shell matrix element, which includes symmetrization effects due to identity of particles in the final state. The new degrees of freedom are introduced via \vec{k}_{Tk} , which are the parton's transverse momenta, i.e. the momenta perpendicular to the collision axis. The formula is valid when the x 's are not too large and not too small when complications from nonlinearities may eventually arise [12]. $\mathcal{M}(i^*, j^* \rightarrow 4 \text{ part.})$ is the gauge invariant matrix element for $2 \rightarrow 4$ particle scattering with two initial off-shell legs calculated numerically with a numerical package [13].

3. Double-parton scattering production of four jets

Single-parton scattering contributions are expected to be dominant for high momentum transfer, as it is highly unlikely that two partons from one proton and two from the other one are energetic enough for two hard scatterings to take place, as the behaviour of the PDFs for large x suggests. However, as the cuts on the transverse momenta of the final state are softened, a window opens to possibly observe significant double parton scattering effects, as often stated in the literature on the subject. First of all, let us recall the formula usually employed for the computation of DPS cross sections, adjusting it to the 4-parton final state,

$$\frac{d\sigma_{4\text{-jet,DPS}}^B}{d\xi_1 d\xi_2} = \frac{m}{\sigma_{eff}} \sum_{i_1, j_1, k_1, l_1; i_2, j_2, k_2, l_2} \frac{d\sigma^B(i_1 j_1 \rightarrow k_1 l_1)}{d\xi_1} \frac{d\sigma^B(i_2 j_2 \rightarrow k_2 l_2)}{d\xi_2}, \quad (3.1)$$

where the $\sigma(ab \rightarrow cd)$ cross sections are obtained by restricting formula (2.1) to a single channel and the symmetry factor m is 1 unless the two hard scatterings are identical, in which case it is 1/2, so as to avoid double counting them. Above ξ_1 and ξ_2 stand for generic kinematical variables for the first and second scattering, respectively. The effective cross section σ_{eff} can be interpreted as a measure of the transverse correlation of the two partons inside the hadrons, whereas the possible longitudinal correlations are usually neglected. In this paper we use σ_{eff} provided by the CDF, D0 collaborations and recently confirmed by the LHCb collaboration $\sigma_{eff} = 15 \text{ mb}$, when all SPS mechanisms of double charm production are included.

3.1 Comparison to CMS data

As discussed in Ref. [15], so far the only experimental analysis of four-jet production relevant for the DPS studies was realized by the CMS collaboration [14]. The cuts used in this analysis are $p_T > 50 \text{ GeV}$ for the first and second jets, $p_T > 20 \text{ GeV}$ for the third and fourth jets, $|\eta| < 4.7$ and the jet cone radius parameter $\Delta R > 0.5$.

It goes without saying that the LO result with soft cuts applied needs refinements from NLO contributions. For this reason, in the following we will always perform comparisons only to data (re)normalised to the total (SPS+DPS) cross sections. What is interesting in the HEF result, compared to collinear factorization, is the dramatic damping of the DPS contribution. In Figs. 1 and 2 we compare the predictions in HEF to the CMS data. Here both the SPS and DPS contributions are normalized to the total cross section, i.e. the sum of the SPS and DPS contributions. In all cases the renormalized transverse momentum distributions agree with the CMS data. However, the absolute cross sections obtained in this case within the HEF approach are too large.

Not only transverse momentum dependence is interesting. The CMS collaboration extracted also a more complicated observables [14]. One of them, which involves all four jets in the final




Figure 1: Comparison of the LO collinear and HEF predictions to the CMS data for the 1st and 2nd leading jets. In addition we show the ratio of the SPS HEF result to the CMS data.




Figure 2: Comparison of the LO collinear and HEF predictions to the CMS data for the 3rd and 4th leading jets. In addition we show the ratio of the SPS HEF result to the CMS data.

state, is the ΔS variable, defined in Ref. [14] as the angle between pairs of the harder and the softer jets,

$$\Delta S = \arccos \left(\frac{\vec{p}_T(j_1^{hard}, j_2^{hard}) \cdot \vec{p}_T(j_1^{soft}, j_2^{soft})}{|\vec{p}_T(j_1^{hard}, j_2^{hard})| \cdot |\vec{p}_T(j_1^{soft}, j_2^{soft})|} \right), \quad (3.2)$$

where $\vec{p}_T(j_i, j_k)$ stands for the sum of the transverse momenta of the two jets in arguments.

In Fig. 3 we present our HEF prediction for the normalized to unity distribution in the ΔS variable. Our HEF result approximately agrees with the experimental ΔS distribution. In contrast the LO collinear approach leads to $\Delta S = 0$, i.e. a Dirac-delta peak at $\Delta S = 0$ for the distribution in ΔS . For the DPS case this is rather trivial. The two hard and two soft jets come in this case from the same scatterings and are back-to-back (LO), so each term in the argument of \arccos is zero (jets are balanced in transverse momenta). For the SPS case the transverse momenta of the two jet pairs (with hard jets and soft jets) are identical and have opposite direction (the total transverse momentum of all four jets must be zero from the momentum conservation). Then it is easy to see that the argument of \arccos is just -1. This means $\Delta S = 0$. The above relations are not fulfilled in

the HEF approach. The SPS contribution clearly dominates and approximately gives the shape of the ΔS distribution. It is anyway interesting that we describe the data via pQCD effects within our HEF approach which are in Ref. [14] described by parton-showers and soft MPIs.




Figure 3: Comparison of the HEF predictions to the CMS data for ΔS spectrum. In addition we show the ratio of the (SPS+DPS) HEF result to the CMS data.

Acknowledgments

The work of K.K. have been supported by Narodowe Centrum Nauki with Sonata Bis grant DEC-2013/10/E/ST2/00656. The paper is based on common publication with Rafal Maciula, Mirko Serino, Andreas van Hameren, Antoni Szczurek.

References

- [1] K. Kutak, R. Maciula, M. Serino, A. Szczurek and A. van Hameren, *JHEP* **1604** (2016) 175
doi:10.1007/JHEP04(2016)175.
- [2] K. Kutak, R. Maciula, M. Serino, A. Szczurek and A. van Hameren, *Phys. Rev. D* **94** (2016) no.1, 014019 doi:10.1103/PhysRevD.94.014019 [arXiv:1605.08240 [hep-ph]].
- [3] S. Catani, M. Ciafaloni and F. Hautmann, *Phys. Lett. B* **242** (1990) 97.
doi:10.1016/0370-2693(90)91601-7
- [4] Z. Bern, *et al.*, Four-Jet Production at the Large Hadron Collider at Next-to-Leading Order in QCD, *Phys. Rev. Lett.*, **109**(2012), 042001, 1112.3940.
- [5] Simon Badger, Benedikt Biedermann, Peter Uwer, and Valery Yundin, NLO QCD corrections to multi-jet production at the LHC with a centre-of-mass energy of $\sqrt{s} = 8$ TeV, *Phys. Lett.*, **B718**(2013), 965, 1209.0098.
- [6] F. Abe *et al.* (CDF collaboration), Study of four jet events and evidence for double parton interactions in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV, *Phys. Rev.*, **D47**(1993), 4857.
- [7] Rafal Maciula and Antoni Szczurek, Production of $c\bar{c}c\bar{c}$ in double-parton scattering within k_t -factorization approach – meson-meson correlations, (2013), 1301.4469.

- [8] Matteo Rinaldi, Sergio Scopetta, Marco Traini, and Vicente Vento, Double parton correlations and constituent quark models: a Light Front approach to the valence sector, *JHEP*, **12**(2014), 028, 1409.1500.
- [9] Wojciech Broniowski, Enrique Ruiz Arriola, and Krzysztof Golec-Biernat, Generalized Valon Model for Double Parton Distributions, in Theory and Experiment for Hadrons on the Light-Front (Light Cone 2015) Frascati , Italy, September 21-25, 2015, 2016, 1602.00254, <http://inspirehep.net/record/1418836/files/arXiv:1602.00254.pdf>.
- [10] Krzysztof Golec-Biernat, Emilia Lewandowska, Mirko Serino, Zachary Snyder, and Anna M. Stasto, Constraining the double gluon distribution by the single gluon distribution, *Phys. Lett.*, **B750**(2015), 559, 1507.08583.
- [11] K. Golec-Biernat and A. M. Stasto, arXiv:1611.02033 [hep-ph].
- [12] P. Kotko, *et al.*, Improved TMD factorization for forward dijet production in dilute-dense hadronic collisions, *JHEP*, **09**(2015), 106, 1503.03421.
- [13] A. van Hameren, arXiv:1611.00680 [hep-ph].
- [14] Serguei Chatrchyan *et al.* (CMS), Measurement of four-jet production in proton-proton collisions at $\sqrt{s} = 7\text{v}2009\text{v}2009\text{TeV}$, *Phys. Rev.*, **D89**(2014) (9), 092010, 1312.6440.
- [15] Rafal Maciula and Antoni Szczurek, Searching for and exploring double-parton scattering effects in four-jet production at the LHC, *Phys. Lett.*, **B749**(2015), 57, 1503.08022.