

Excessive double strange baryon production due to strangeness oscillation in p+A, A+A collisions

Peter Filip^{*†}

Institute of Physics, Slovak Academy of Sciences, Dúbravská cesta 9, Bratislava 845 11, Slovakia

E-mail: fyziflip@savba.sk

Production of double strange Ξ^- hyperons at sub-threshold energies has been observed by HADES experiment [1] to be unexpectedly enhanced in comparison to theoretical estimates. We suggest, that $K^0 \leftrightarrow \bar{K}^0$ oscillation of neutral kaons can be affected in very dense baryonic matter in a specific way, which may result in the oscillation length 5-10 fm. This allows for the strangeness violation process $(\bar{s}d) \rightarrow (s\bar{d})$ to occur in a very short time, within the volume of dense hadronic medium, and excessive double strange hyperons can be created via rescattering $\bar{K}^0 + (\Sigma^0, \Lambda) \rightarrow \Xi + \pi$ interactions. The significance of such processes is underestimated, if global strangeness conservation is assumed in $p + A$ and $A + A$ collisions at low energies.

*38th International Conference on High Energy Physics
3-10 August 2016
Chicago, USA*

^{*}Speaker.

[†]Support from ICHEP organizers and from Slovak Grant Agency VEGA (project 2/197/14) is acknowledged.

1. Introduction

The oscillation of neutral mesons ($K^0 \leftrightarrow \bar{K}^0$) is a beautiful quantum mechanical phenomenon, which allows [2] us to study also the fundamental properties of nature (CP symmetry). $B_d^0 \leftrightarrow \bar{B}_d^0$ and $B_s^0 \leftrightarrow \bar{B}_s^0$ oscillations have been clearly observed as well [3], while D^0, \bar{D}^0 mesons containing only up -type quarks have been proved to oscillate only recently [4].


The period of $K^0 \leftrightarrow \bar{K}^0$ transitions (oscillation length L_{osc}) is determined by the mass difference $\Delta m_{K^0} = m(K_2^0) - m(K_1^0)$ of eigenstates K_2^0 and K_1^0 of the weak \mathbb{H}_w hamiltonian. Measured values of $\Delta m_{K^0}, \Delta m_{D^0}, \Delta m_{B_d^0}$, and $\Delta m_{B_s^0}$ mass differences *in vacuum* are (in $10^{10}\hbar/s$ units): 0.529 ± 0.001 , 0.95 ± 0.44 , 51.0 ± 0.3 , and 1776 ± 2 , which gives [3] oscillation lengths $L_{osc} = c\hbar/\Delta m$: 35cm, ≈ 20 cm, 3.7mm, and 0.11mm. Standard Model explains these oscillations successfully by 2^{nd} order flavour-changing transitions $(s\bar{d}) \leftrightarrow (\bar{s}d)$ and $(c\bar{u}) \leftrightarrow (\bar{c}u)$ and $(b\bar{d}, \bar{s}) \leftrightarrow (\bar{b}d, s)$ taking place in the vacuum due to short-distance (box diagrams) and long-distance effects [3].

In the material medium (a regenerator), or in a dense nuclear matter, the value of Δm_{K^0} mass difference of K_2^0, K_1^0 eigenstates may become modified due to meson-baryon (repulsive) and antimeson-baryon (attractive) potentials [5]. Linear approximation (see Eq. 66 and 67 in [5])

$$m_{K^0}(\rho) = (1 + \alpha_K \frac{\rho}{\rho_0}) m_{K^0}^{\rho=0} \quad ; \quad m_{\bar{K}^0}(\rho) = (1 - \tilde{\alpha}_{\bar{K}} \frac{\rho}{\rho_0}) m_{\bar{K}^0}^{\rho=0}. \quad (1.1)$$

gives $m(K^0) - m(\bar{K}^0) \approx 80 \text{ MeV}$ at $\rho = \rho_0$ density, if values $\alpha_K \approx 0.05$ and $\tilde{\alpha}_{\bar{K}} \approx 0.12$ are used.

In this contribution we suggest Δm_{K^0} mass difference in dense baryonic medium may become so large, that $(\bar{s} \rightarrow s)$ transition length $L^{\bar{s} \rightarrow s} = c\hbar/2\Delta m_{K^0}$ can be very short: 2 – 10 fm. This may allow for a *single* $(s\bar{s})$ pair (created in the low-energy $p + A$ and $A + A$ collisions) to be sufficient for the production of double strange $\Xi^-(ssd)$ hyperons via process:



At high baryonic densities, \bar{s} quarks preferentially hadronize into $K^0(d\bar{s})$ or $K^+(u\bar{s})$ mesons, while s quarks are trapped into hyperons. When multiplicities of neutral $K^0(d\bar{s})$ and $\bar{K}^0(s\bar{d})$ mesons are very asymmetric (e.g. $N[K]/N[\bar{K}] \geq 100$), mesons $K^0(d\bar{s})$ may enhance s quark population via fast $K^0 \rightarrow \bar{K}^0$ transition. Excessive $\Xi^-(ssd)$ or $\Xi^0(ssu)$ hyperons may thus be created in $p + A$ and $A + A$ collisions at *subthreshold* energy [1] via rescattering process: $\bar{K}^0 + (\Sigma^0, \Lambda) \rightarrow \Xi + \pi$.

2. Neutral kaons in dense baryonic medium

In the vacuum (if CP violation effects $|\varepsilon| \approx 2 \cdot 10^{-3}$ are neglected) the eigenstates $K_{1,2}^0$ of weak Hamiltonian \mathbb{H}_w are: $K_{1,2}^0 = (K^0 \pm \bar{K}^0)/\sqrt{2}$. In a medium, Hamiltonian $\mathbb{H}'_w = \mathbb{M}' - \frac{i}{2}\mathbb{G}'$ is

$$\mathbb{H}'_w = \left[\begin{array}{cc} M_{11} + V_{K^0}(\rho) & M_{12} \\ M_{21} & M_{22} - \bar{V}_{\bar{K}^0}(\rho) \end{array} \right] - \frac{i}{2} \left(\begin{array}{cc} \Gamma_{11} + A_{K^0}(\rho) & \Gamma_{12} \\ \Gamma_{21} & \Gamma_{22} + \bar{A}_{\bar{K}^0}(\rho) \end{array} \right) \quad (2.1)$$

and linear approximation (1.1) gives potentials $V_{K^0} = m_{K^0} \alpha_K(\rho/\rho_0)$ and $\bar{V}_{\bar{K}^0} = m_{\bar{K}^0} \tilde{\alpha}_{\bar{K}}(\rho/\rho_0)$. This means $V_{K^0} \approx 20 \text{ MeV}$ and $\bar{V}_{\bar{K}^0} \approx 60 \text{ MeV}$ at nuclear density $\rho \approx \rho_0 = 2 \cdot 10^{17} \text{ kg/m}^3$, for (momentum

averaged [5]) parameters $\alpha_K \approx 0.04$ and $\tilde{\alpha}_{\bar{K}} \approx 0.12$. Absorption coefficients $A_{K^0}, \bar{A}_{\bar{K}^0}$ in (2.1) are related to forward scattering amplitude difference $f_K(0) - \bar{f}_{\bar{K}}(0)$ of K^0, \bar{K}^0 mesons in medium [6].

Diagonalization of 2×2 non-hermitian Hamiltonian (2.1) with $M_{11} = M_{22} = 497 \text{ MeV}$ and $\Gamma_{11} = \Gamma_{22} = 3.7 \cdot 10^{-12} \text{ MeV}$ (using $|\Gamma_{12}| = |\Gamma_{21}| \approx 3.48 \cdot 10^{-12}$ and $|M_{12}| = |M_{21}| \approx 1.74 \cdot 10^{-12}$) allows to obtain difference of $K_{1,2}^0$ eigenstate masses and decay widths in medium [7] as

$$\Delta\mu = \tilde{m}(K_2^0) - \tilde{m}(K_1^0) - \frac{i}{2}(\tilde{\Gamma}_{K_2^0} - \tilde{\Gamma}_{K_1^0}) = \sqrt{4H_{12}H_{21} + (H_{22} - H_{11})^2} = \Delta\tilde{m}_K - \frac{i}{2}\Delta\tilde{\Gamma}_K. \quad (2.2)$$

Probabilities of $K^0 \leftrightarrow \bar{K}^0$ transitions are (using Eq. 9.7 and 9.8 from Ref. [7])

$$P[K^0 \rightsquigarrow \bar{K}^0] = \left| \frac{q_H}{p_H} \right|^2 |g_-(\tau)|^2 |(1 - \theta)|^2, \quad P[\bar{K}^0 \rightsquigarrow K^0] = \left| \frac{p_L}{q_L} \right|^2 |g_-(\tau)|^2 |(1 - \theta)|^2, \quad (2.3)$$

where the interference term

$$|g_-(\tau)|^2 = \frac{1}{4} \left[e^{-\tau\tilde{\Gamma}_{K_2^0}(\rho)} + e^{-\tau\tilde{\Gamma}_{K_1^0}(\rho)} - 2 \cos[\Delta\tilde{m}_K(\rho)\tau] \cdot e^{-\tau[\tilde{\Gamma}_{K_2^0}(\rho) + \tilde{\Gamma}_{K_1^0}(\rho)]/2} \right] \quad (2.4)$$

in (2.3) is multiplied by $|q_H/p_H| = 2|H_{21}|/|\Delta\mu(1 - \theta)|$ quantity [7]. Conservation of CP symmetry gives $|q_H/p_H| = |p_L/q_L|$, and weak hamiltonian \mathbb{H}'_w eigenvectors are: $K_2^0 = p_H|K^0\rangle + q_H|\bar{K}^0\rangle$ and $K_1^0 = p_L|K^0\rangle - q_L|\bar{K}^0\rangle$. Consequently, one obtains for $K^0 \rightsquigarrow \bar{K}^0$ transition probability

$$P[K^0 \rightsquigarrow \bar{K}^0] = \frac{|2H_{21}|^2}{|\Delta\mu|^2} |g_-(\tau)|^2 \approx \frac{4|M_{21} - \frac{i}{2}\Gamma_{21}|^2}{|H_{22} - H_{11}|^2} |g_-(\tau)|^2 = S_\rho |g_-(\tau)|^2 \quad (2.5)$$

where S_ρ is the suppression factor of $|\Delta S| = 2$ process $K^0(d\bar{s}) \rightsquigarrow \bar{K}^0(s\bar{d})$ in the medium. For $\bar{K}^0, K^0(497)$ pseudoscalar mesons at nuclear density $\rho \approx \rho^0$, one may expect $|H_{22} - H_{11}| \approx 100 \text{ MeV}$. Values $|\Gamma_{12}| = |\Gamma_{21}| = 3.48 \cdot 10^{-12}$ and $|M_{12}| = |M_{21}| = 1.74 \cdot 10^{-12} \text{ MeV}$ [7] then give enormous suppression factor $S_\rho \leq 10^{-26}$ for $K^0 \leftrightarrow \bar{K}^0$ oscillations in nuclear medium, in agreement with [8].




Figure 1: $K^{0*} \rightarrow \bar{K}^{0*}$ transition probability as a function of time τ obtained for density $\rho/\rho_0 = 1.2$.




Figure 2: Probability of $K^{0*} \rightarrow \bar{K}^{0*}$ oscillation as a function of time and baryonic density $\rho < 1.5\rho_0$.

However, for $K^{0*}(896)$ and $\bar{K}^{0*}(896)$ mesons, which may also form weak eigenstates [9] $K_{L,S}^{0*} = (K^{0*} \pm \bar{K}^{0*})/\sqrt{2}$, one obtains $S_\rho \approx 10^{-2}$ (see Figures 1 and 2), assuming K^{0*}, \bar{K}^{0*} mesons share 33% of their decay products ($K^{0*} \rightarrow K_{S,L}^0 + \pi^0$ and $\bar{K}^{0*} \rightarrow K_{S,L}^0 + \pi^0$). Indeed, one has [7]

$$\Gamma_{12} = \rho_c \langle K^{0*} | H'_w | K_S^0 \pi^0 \rangle \langle K_S^0 \pi^0 | H'_w | \bar{K}^{0*} \rangle + \rho_c \langle K^{0*} | H'_w | K_L^0 \pi^0 \rangle \langle K_L^0 \pi^0 | H'_w | \bar{K}^{0*} \rangle \quad (2.6)$$

which gives $\Gamma_{12} \approx 16 \text{ MeV}$, for widths $\Gamma(K^{0*} \rightarrow K_{L,S}^0 + \pi^0) = \Gamma(\bar{K}^{0*} \rightarrow K_{L,S}^0 + \pi^0) = 8 + 8 \text{ MeV}$. Using $\Gamma_{12} = 16 \text{ MeV}$ and $\Gamma_{11} = \Gamma_{22} = \Gamma_{K^*} = 48 \text{ MeV}$ in the Hamiltonian for K^{0*}, \bar{K}^{0*} mesons

$$\mathbb{H}'_{K^{0*}} = \begin{bmatrix} 896 + V_{K^0}(\rho) & 1.7 \cdot 10^{-12} \\ 1.7 \cdot 10^{-12} & 896 - \bar{V}_{\bar{K}^0}(\rho) \end{bmatrix} - \frac{i}{2} \begin{pmatrix} 48 + A_{K^0} & 16 \cdot e^{i\zeta} \\ 16 \cdot e^{-i\zeta} & 48 + \bar{A}_{\bar{K}^0} \end{pmatrix} \quad (2.7)$$

suppression factor $S(\rho_B) \approx 10^{-2}$ is obtained in Eq.(2.5) for $\Delta V_{K^*} = 80 \text{ MeV}$ at density $\rho = \rho^0$. In Figures 3 and 4 we show $\Delta m_{K^*} = m(K_2^{0*}) - m(K_1^{0*})$ mass difference and $K^{0*} \rightarrow \bar{K}^{0*}$ transition length $L^{\bar{s} \rightarrow s} = c \cdot \tau_{osc}/2$ evaluated for K^{0*}, \bar{K}^{0*} mesons in baryonic matter using Hamiltonian (2.7).




Figure 3: Mass difference $\Delta m_{K^*} = m(K_2^{0*}) - m(K_1^{0*})$ of weak eigenstates K_2^{0*}, K_1^{0*} in the nuclear medium.




Figure 4: Dependence of $K^{0*} \rightarrow \bar{K}^{0*}$ transition length $L^{\bar{s} \rightarrow s} = \hbar c / 2\Delta m_{K^*}$ on baryonic density ρ/ρ_0 .

3. Summary and conclusions

We have considered $(d\bar{s}) \leftrightarrow (\bar{d}s)$ oscillations in dense nuclear matter. We suggest $K^{0*} \rightarrow \bar{K}^{0*}$ process may happen in $p + A$ or $A + A$ collisions [1] within time scale (3–10 fm/c) with probability $\approx 1\%$. This may allow for the excessive $\Xi^-(ssd)$ hyperon production via $\bar{K}^0 + (\Sigma^0, \Lambda) \rightarrow \Xi + \pi$ reaction at sub-threshold energies, when single $(s\bar{s})$ pair is produced. If $N(K^{0*})/N(\bar{K}^{0*}) \geq 100$ condition is valid in $A + A$ collisions, (\bar{s}/s) ratios may be modified due to $K^{0*} \rightarrow \bar{K}^{0*}$ processes. In agreement with Ref. [8] we find $K^0 \rightarrow \bar{K}^0$ transitions in dense baryonic matter to be negligible.

Although fast oscillations of K^0, B_s^0, B^0 mesons in the nuclear medium are unlikely, a modification of $\Delta\tilde{m}_K$ and $\Delta\tilde{m}_B$ parameters in dense regenerators might be experimentally observable.

References

- [1] G. Agakishiev, et al., *Phys. Rev. Lett.* **103** (2009) 132301; *Phys. Rev. Lett.* **114** (2015) 212301.
- [2] A. Angelopoulos, et al., *Physics Reports* **374** (2003) 165.
- [3] C. Patrignani et al., (Particle Data Group) *Chin. Phys.* **C40** (2016) 100001.
- [4] A. Aaij, et al., (LHCb Collaboration) *Phys. Rev. Lett.* **110** (2013) 101802.
- [5] C. Hartnack, et al., *Physics Reports* **510** (2012) 119.
- [6] P.H. Eberhard, F. Uchiyama, *Nucl. Inst. Meth. Phys. Res.* **A350** (1994) 144.
- [7] C.G. Branco, L. Lavoura, J.P. Silva, *CP Violation*, Clarendon Press, Oxford (1999).
- [8] G. Amelino-Camelia and J. Kapusta, *Phys. Lett.* **B465** (1999) 291.
- [9] L.S. Littenberg, *Phys. Rev.* **D21** (1980) 2027.