

The Giant Radio Array for Neutrino Detection (GRAND): Present and Perspectives

Ke Fang^{*1,†}, Jaime Álvarez-Muñiz², Rafael Alves Batista³, Mauricio Bustamante⁴, Washington Carvalho⁵, Didier Charrier⁶, Ismaël Cognard^{7,8}, Sijbrand De Jong⁹, Krijn D. de Vries¹⁰, Chad Finley¹¹, Quanbu Gou¹², Junhua Gu¹³, Claire Guépin¹⁴, Jordan Hanson⁴, Hongbo Hu¹², Kumiko Kotera¹⁴, Sandra Le Coz¹³, Yi Mao¹⁵, Olivier Martineau-Huynh¹⁶, Clementina Medina^{16,17}, Miguel Mostafa¹⁸, Fabrice Mottez¹⁹, Kohta Murase¹⁸, Valentin Niess¹⁹, Foteini Oikonomou¹⁸, Frank Schröder²⁰, Cyril Tasse²¹, Charles Timmermans⁹, Nicolas Renault-Tinacci¹⁴, Matías Tueros¹⁷, Xiang-Ping Wu¹³, Philippe Zarka²², Andreas Zech¹⁹, Yi Zhang¹², Qian Zheng¹³, Anne Zilles¹⁴

¹Joint Space-Science Institute, University of Maryland, College Park, MD, 20742, USA

²Departamento de Física de Partículas, Universidade de Santiago de Compostela, 15782, Santiago de Compostela, Spain

³Department of Physics - Astrophysics, University of Oxford, DWB, Keble Road, OX1 3RH, Oxford, UK

⁴Center for Cosmology and AstroParticle Physics, Department of Physics, Ohio State University, Columbus, OH 43210, USA

⁵Physics institute, University of São Paulo, Rua do Matão, trav. R, Cid. Universitária, São Paulo, Brazil

⁶SUBATECH, CNRS-IN2P3, Université de Nantes, Ecole des Mines de Nantes, Nantes, France

⁷Laboratoire de Physique et Chimie de l'Environnement et de l'Espace LPC2E CNRS-Université d'Orléans, F-45071 Orléans, France

⁸Station de radioastronomie de Nançay, Observatoire de Paris, CNRS/INSU F-18330 Nançay, France

⁹Nikhef/Radboud University, Nijmegen, the Netherlands

¹⁰Vrije Universiteit Brussel, Dienst ELEM, B-1050 Brussels, Belgium

¹¹Oskar Klein Centre and Dept. of Physics, Stockholm University, SE-10691 Stockholm, Sweden

¹²Key Lab of Particle Astrophysics, IHEP, Chinese Academy of Sciences, Beijing 100049, China

¹³National Astronomical Observatory, Chinese Academy of Sciences, Beijing 100012, China

¹⁴Sorbonne Univ., Paris 6, CNRS UMR 7095, Institut d'Astrophysique de Paris, 98 bis bd Arago, 75014 Paris, France

¹⁵Department of Physics, Tsinghua Center for Astrophysics, Tsinghua University, Beijing 100084, China

¹⁶LPNHE, CNRS-IN2P3 and Universités Paris 6 & 7, BP200, 4 place Jussieu, 75252 Paris, France

¹⁷Instituto de Física La Plata, CONICET CCT-La Plata, Calle 49 esquina 115 (1900), La Plata, Argentina

¹⁸Dept. of Physics, Dept. of Astronomy & Astrophysics, Penn State University, University Park, PA, USA

¹⁹Clermont Université, Université Blaise Pascal, CNRS/IN2P3, Laboratoire de Physique Corpusculaire, BP. 10448, 63000 Clermont-Ferrand, France

²⁰Institut für Kernphysik, Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

²¹LUTH, Observatoire de Paris, CNRS, Université Paris Diderot, PSL Research University, 5 place Jules Janssen, 92190, Meudon, France

²²LESIA & USN, Observatoire de Paris, CNRS, PSL, UPMC-SU/UPD/UO, 92195 Meudon, France

The Giant Radio Array for Neutrino Detection (GRAND) aims at detecting ultra-high energy extraterrestrial neutrinos via the extensive air showers induced by the decay of tau leptons created in the interaction of neutrinos under the Earth's surface. Consisting of an array of $\sim 10^5$ radio antennas deployed over $\sim 2 \times 10^5 \text{ km}^2$, GRAND plans to reach, for the first time, an all-flavor sensitivity of $\sim 1.5 \times 10^{-10} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ above $5 \times 10^{17} \text{ eV}$ and a sub-degree angular resolution, beyond the reach of other planned detectors. We describe here preliminary designs and simulation results, plans for the ongoing, staged approach to the construction of GRAND, and the rich research program made possible by GRAND's design sensitivity and angular resolution.

*35th International Cosmic Ray Conference — ICRC2017
10–20 July, 2017
Bexco, Busan, Korea*

*Speaker.

†E-mail: kefang@umd.edu

1. Introduction

The workings of the nature's highest-energy accelerators are a half-century-old mystery [1]. It is difficult to use ultra-high energy cosmic rays (UHECR, with $E > 10^{18}$ eV) to pinpoint a source as these charged particles can be deflected by up to tens of degrees by extragalactic and Galactic magnetic fields. On the contrary, secondary neutrinos, produced when cosmic rays interact with matter inside a source or with the cosmic microwave background (CMB) during intergalactic propagation, serve as direct probes, as neutrinos are not deflected and barely interact over cosmological distances. Two steps need to be achieved toward finding sources of UHE neutrinos:

1. Detecting an UHE neutrino. Depending on the chemical composition and maximum energy of UHECRs, the flux of cosmogenic neutrino background is expected at a level of $\sim 10^{-9} - 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ per flavor at EeV energies. A small number of diffuse neutrinos found in this step, however, will not reveal an UHE neutrino point source.
2. Identifying individual sources. This step requires the detection of a sizable number of neutrinos to allow point sources to stand out of a diffuse background [2, 3]. For instance, if sources follow a uniform number density of 10^{-7} Mpc^{-3} , a 5σ detection of point sources requires measuring ~ 200 neutrino events with a sub-degree angular resolution [3].

Astrophysical tau neutrinos (ν_τ) can be detected through extensive air showers (EAS) induced by tau (τ) decays in the atmosphere [4, 5]. Specifically, a cosmic ν_τ can produce tau leptons under the Earth surface via charge-current interactions. The short-lived τ decays in the atmosphere after traveling tens of kilometers and generates an EAS that emits coherent electromagnetic emissions up to frequencies of hundreds of MHz. Such inclined showers have been measured and confirmed by the Auger Engineering Radio Array to have a radio footprint covering several km^2 [6]. With practical advantages such as low unit cost, easiness of deployment and maintenance, radio antennas present appealing characteristics for the detection of very inclined showers [7].

Consisting of $\sim 10^5$ radio antennas covering $200,000 \text{ km}^2$ in a mountainous area, the Giant Radio Array for Neutrino Detection (GRAND) will detect air showers generated by Earth-skimming UHE tau neutrinos with unprecedented sensitivity. Mountain ranges with low electromagnetic background noise are preferable sites, as mountains provide both additional interaction targets and a slanted surface that is more favorable to detect inclined showers compared to flat ground. The antennas are foreseen to operate in the 60-200 MHz band to avoid the short-wave background at lower frequencies. The highest frequencies enable detection of the Cherenkov ring associated with the air shower [8], which can be used as an unambiguous signature to reject background.

2. Detector performance

2.1 Sensitivity

We present a preliminary evaluation of the GRAND sensitivity based on a simulation of the response of 90,000 antennas deployed on a square layout of $60,000 \text{ km}^2$ in a remote mountainous area (Tianshan mountains, XinJiang province, China). The simulations are composed of three




Figure 1: Three-year differential sensitivity of a 10,000 km² simulated setup (red thin curve) and of the projected GRAND array (red thick curve). The grey shaded region brackets fluxes of cosmogenic neutrinos from models with different cosmic ray compositions, spectra, and source evolution histories that provide reasonable fits to the Auger UHECR spectrum [9]. The grey dashed and dotted lines indicate the predicted cosmogenic neutrino flux in a pessimistic scenario where the bulk of UHECRs are iron nuclei. High-energy neutrinos measured by IceCube [10], UHE neutrino sensitivities of IceCube [11], Auger [12], ARA [13] and ARIANA [14] are also shown for reference.

parts: I) ν_τ s interact with the rock, producing τ leptons, II) τ s propagate and decay, and III) The τ decay products initiate an EAS that triggers the antennas to be read-out.

In part I, standard rock with a density of 2.65 g cm^{-3} is assumed down to the sea level, and the Earth core is modeled following the Preliminary Reference Earth Model [15]. We simulate the deep inelastic scattering of neutrinos with Pythia6.4, using the CTEQ5d probability distribution functions (PDF) combined with the cross sections from [16].

In part II, we simulate the propagation of the produced τ s with GEANT4.9, using parameterizations of PDFs of the τ path length and proper time. We simulate τ decays with the TAUOLA package [17]. For the above two parts, 1D tracking of a primary ν_τ is performed down to the converted τ decay.

We have refined our previous calculation of part III [18] with numerical simulations. We first simulate a set of ν_τ showers with ZHaires [19] at different energies, and determine a conical volume inside which the electric field is above the expected detection threshold of the GRAND antennas. Based on a parametrization of the shape (including angle and height) of the detection cones as a function of energy, we compute the expected shape and position of cones for each neutrino-initiated EAS in our simulation (so far in the 30-80 MHz band). We select the antennas located inside the cones, taking into account signal shadowing by mountains. When the selection




Figure 2: Three-year integrated exposure (in units of $\text{cm}^2 \text{s}$) of GRAND10k at 3 EeV in Galactic coordinates.

consists of 8 or more neighboring antennas in direct view of the τ decay point, we count the primary ν_τ as detected.

Our analysis finds that about 50% of detectable neutrino interactions take place inside mountains rather than in the Earth crust. Moreover, some parts of the array provide 2-3 times higher detection rates compared to the average rate of the entire simulated area. These “hotspots” are typically located in a low electromagnetic noise background regions and have desirable geometric features (for e.g., large mountains, slopes facing another mountain range at distances of 30-80 km). An envisioned GRAND setup is composed of smaller sub-arrays of a few 10,000 km^2 each, deployed solely on such favorable sites.

Figure 1 presents the differential 90%-CL sensitivity of a GRAND simulation for a 10,000 km^2 hotspot and the projected full GRAND array limit after a 3-year observation. An all-flavor 90% CL integral limit of $2 \times 10^{-10} \text{GeVcm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ is derived for an unbroken E^{-2} power law spectrum in an aggressive scenario (corresponding to a detection threshold of $30 \mu\text{V m}^{-1}$; $3.9 \times 10^{-9} \text{GeVcm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ in a conservative scenario with a detection threshold of $100 \mu\text{V m}^{-1}$). For comparison, a prediction of the cosmogenic neutrino flux is shown as the grey shaded region. The flux mostly depends on three properties of UHECRs: spectral index, chemical composition, and maximum energy E_{max} . The shaded area brackets a wide range of parameters that reflect different compositions, source evolution and Galactic to extragalactic transition models, all required to fit the UHECR spectrum measured by Auger [20]. Specifically, the upper bound corresponds to an UHECR dip model with a pure proton composition and source evolution following the star formation rate (SFR). The lower bound corresponds to a mixed composition with 30% iron nuclei, SFR evolution and $E_{\text{max}} = 10^{19} Z \text{ eV}$. The grey dotted and dashed lines show the most pessimistic scenarios, where UHECRs are mostly composed of iron nuclei which barely produce neutrinos in a photo-disintegration process. We note that the best-fit models, found by a combined fit to the Auger spectrum, X_{max} and the standard derivation of X_{max} measurements [21], correspond to a cosmogenic neutrino flux level of $\sim 10^{-9} \text{GeVcm}^{-2}\text{s}^{-1}\text{sr}^{-1}$. For comparison, we show UHE neutrino upper limits of current experiments IceCube [11] and Auger [12] (the sensitivity curve of ANITA-II [22] is not shown as it would be too high for the plot). GRAND10k will be as sensitive as the proposed ARA [13] and ARIANA [14] experiments at 1 EeV. The full GRAND array will nominally detect even the lowest expected fluxes of cosmogenic neutrinos.

Following [23], the angular resolution of the radio array is estimated by the resolution of the time difference between pulses in different antennas. Using a 3 ns precision on the antenna trigger timing, GRAND can reach an angular resolution as accurate as 0.1° .

The instantaneous field of view covers a band with zenith angles between 86° and 93° in the simulation (the zenith angle-dependent effective area is shown in Figure 1 of [18]). The integrated exposure over 3 years of GRAND operation will cover a large fraction of the sky as shown in Figure 2. With stacking searches, GRAND will be sensitive to transient sources and flaring events with a duration longer than ~ 1 day. Several interesting UHECR features, such as the Telescope Array hotspot and the Auger hotspot, are located close to GRAND's best sensitivity region.

2.2 Background rejection

Radio background from high-energy particles other than cosmic neutrinos should be manageable. The atmospheric neutrino background is mostly negligible above 10^{16} eV. EAS generated by muon decay are expected at a rate of a few per century as muons have a significantly longer life time than taus. Rejecting trajectories far from the horizon should effectively suppress the EAS generated by cosmic rays.

Background event rates associated to terrestrial sources such as human activities and thunderstorms are difficult to evaluate. The background measurement of the Tianshan Radio Experiment for Neutrino Detection (TREND) [23] provides a conservative estimate. TREND consisted of 50 self-triggered antennas deployed in a populated valley of the Tianshan mountains, with an antenna design and sensitivity similar to what is foreseen for GRAND. The observed rate of events triggering six selected TREND antennas separated by ~ 800 m over a sample period of 120 live days was found to be around 1 per day, among which two-thirds arrived as bursts of events mostly due to air plane activities. An extrapolation from this measurement leads to an expected background rate of 1 Hz for GRAND for a trigger algorithm based on coincident pulses on neighboring antennas and a rejection of event bursts.

Specific amplitude patterns on the ground (such as beamed emission along the shower axis and signal enhancement along the Cherenkov ring [8]), and polarization of signals [24] are strong indicators of neutrino-initiated EAS. We are investigating the possibilities of using these signatures as efficient background discrimination tools through both simulations and experiments.

3. Science case

In addition to a guaranteed detection of cosmogenic neutrinos, GRAND will allow a differentiation of the source types by measuring neutrinos produced inside sources. In Figure 3, we show fluxes of neutrinos from four typical classes of astrophysical sources that can explain UHECR measurements, including galaxy clusters with central sources [25, 26], fast-spinning newborn pulsars [27], active galactic nuclei [28], and afterglows of gamma-ray bursts [29]. The GRAND sensitivity is sufficient to observe neutrinos from sources in all cases.

A 5σ identification of individual point sources out of a diffuse background requires $\sim 100 - 1000$ events with a sub-degree angular resolution for sources that have a local density of $10^{-9} - 10^{-7} \text{ Mpc}^{-3}$ [3]. Assuming a neutrino flux of $10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$, the GRAND neutrino sen-




Figure 3: Three-year differential sensitivity of the GRAND10k and GRAND full array comparing to fluxes of neutrinos from interactions inside astrophysical sources. Four source classes that can explain UHECR measurements are shown: galaxy clusters with central sources [25, 26], fast-spinning newborn pulsars [27], active galactic nuclei [28], and afterglows of gamma-ray bursts [29]. High-energy neutrino observed by IceCube [10], UHE neutrino sensitivities of IceCube [11] and Auger [12] are shown for reference.

sensitivity corresponds to a detection of ~ 100 events after three years of observation. The unprecedented sensitivity of GRAND is crucial for a direct identification of UHE neutrino point sources.

GRAND will observe UHECRs with an effective area that is at least an order of magnitude larger than Auger. The high statistics will resolve any small-scale anisotropies and features near the end of the cosmic ray spectrum. GRAND will also reach an UHE photon sensitivity exceeding that of current experiments. Future simulation studies will be dedicated to performance of energy and X_{\max} reconstructions.

GRAND will contribute in a unique way to the measurement of fast radio bursts (FRB) and giant radio pulses by collecting unprecedented statistics at low frequencies [30]. Simulations are being performed to quantify the detection power with incoherent sum of FRB spectra computed at each antenna. The possibility of GRAND's contribution to additional science topics, such as the study of the epoch of reionization, is also being investigated.

4. Development plan

The GRAND project aims at building a next-generation neutrino telescope composed of a radio antenna array deployed over $200,000 \text{ km}^2$. Preliminary simulations indicate that 5% of the array deployed in favorable sites will improve the sensitivity over that of current-generation telescopes by an order of magnitude. The full array will offer a sensitivity that ensures a detection of cosmogenic neutrinos in the most pessimistic scenario, and an identification of individual point sources

in an optimistic scenario. GRAND will also be a powerful instrument for UHECR observations with high statistics. Simulation and experimental work is ongoing on technological development and background rejection strategies.

The GRAND development plan consists of several steps. Presently, deployment of GRANDproto [31], which is a hybrid detector composed of 35 3-arm antennas and 24 scintillators, is underway at the site of the TREND experiment, in the Tianshan mountains. The GRANDproto project will demonstrate that EAS can be detected with an autonomous radio array with both high efficiency and high purity. Following this step, a dedicated setup with a size of 300 km² will be deployed. This array will establish the autonomous radio detection of very inclined EAS with excellent background rejection, as a validation for the future GRAND layout.

Acknowledgments

The GRAND and GRANDproto project are supported by the France China Particle Physics Laboratory, the Institut Lagrange de Paris, the APACHE grant (ANR-16-CE31- 0001) of the French Agence Nationale de la Recherche, the Natural Science Foundation of China (Nos.11135010, 11105156, 11375209 and 11405180), the Chinese Ministry of Science and Technology, the Youth Innovation Promotion Association of Chinese Academy of Sciences, and the São Paulo Research Foundation FAPESP (grant 2015/15735-1).

References

- [1] K. Kotera and A. V. Olinto, “The Astrophysics of Ultrahigh Energy Cosmic Rays,” *Ann.Rev.Astron.Astrophys.*, vol. 49, pp. 119–153, 2011.
- [2] M. Ahlers and F. Halzen, “Pinpointing Extragalactic Neutrino Sources in Light of Recent IceCube Observations,” *Phys. Rev.*, vol. D90, no. 4, p. 043005, 2014.
- [3] K. Fang, K. Kotera, M. C. Miller, K. Murase, and F. Oikonomou, “Identifying ultrahigh-energy cosmic-ray accelerators with future ultrahigh-energy neutrino detectors,” *JCAP*, vol. 12, p. 017, Dec. 2016.
- [4] D. Fargion, “Horizontal Tau air showers from mountains in deep vally :Traces of Ultrahigh neutrino tau,” *International Cosmic Ray Conference*, vol. 2, p. 396, Aug. 1999.
- [5] X. Bertou, P. Billoir, O. Deligny, C. Lachaud, and A. Letessier-Selvon, “Tau neutrinos in the Auger Observatory: a new window to UHECR sources,” *Astroparticle Physics*, vol. 17, pp. 183–193, May 2002.
- [6] Kambeitz, Olga, “Measurement of horizontal air showers with the auger engineering radio array,” *EPJ Web Conf.*, vol. 135, p. 01015, 2017.
- [7] F. G. Schröder, “Radio detection of cosmic-ray air showers and high-energy neutrinos,” *Progress in Particle and Nuclear Physics*, vol. 93, pp. 1–68, Mar. 2017.
- [8] J. Alvarez-Muñiz, W. R. Carvalho, Jr., A. Romero-Wolf, M. Tueros, and E. Zas, “Coherent radiation from extensive air showers in the ultrahigh frequency band,” *PRD*, vol. 86, p. 123007, Dec. 2012.
- [9] K. Kotera, D. Allard, and A. V. Olinto, “Cosmogenic neutrinos: parameter space and detectability from PeV to ZeV,” *JCAP*, vol. 10, p. 013, Oct. 2010.
- [10] The IceCube Collaboration, “The IceCube Neutrino Observatory - Contributions to ICRC 2015 Part II: Atmospheric and Astrophysical Diffuse Neutrino Searches of All Flavors,” *ArXiv 1510.05223*.

- [11] M. G. Aartsen, K. Abraham, M. Ackermann, and et al., “Constraints on Ultrahigh-Energy Cosmic-Ray Sources from a Search for Neutrinos above 10 PeV with IceCube,” *Physical Review Letters*, vol. 117, p. 241101, Dec. 2016.
- [12] A. Aab, P. Abreu, M. Aglietta, et al., “Improved limit to the diffuse flux of ultrahigh energy neutrinos from the pierre auger observatory,” *Phys. Rev. D*, vol. 91, p. 092008, May 2015.
- [13] Ara Collaboration, “Design and initial performance of the Askaryan Radio Array prototype EeV neutrino detector at the South Pole,” *Astroparticle Physics*, vol. 35, pp. 457–477, Feb. 2012.
- [14] S. W. Barwick, E. C. Berg, D. Z. Besson, et al., “A first search for cosmogenic neutrinos with the ARIANNA Hexagonal Radio Array,” *Astroparticle Physics*, vol. 70, pp. 12–26, Oct. 2015.
- [15] A. M. Dziewonski and D. L. Anderson, “Preliminary reference Earth model,” *Physics of the Earth and Planetary Interiors*, vol. 25, pp. 297–356, June 1981.
- [16] R. Gandhi, C. Quigg, M. H. Reno, and I. Sarcevic, “Neutrino interactions at ultrahigh energies,” *PRD*, vol. 58, p. 093009, Nov. 1998.
- [17] S. Jadach, J. H. KÄijhn, and Z. WałgŁs, “Tauola - a library of monte carlo programs to simulate decays of polarized ĪD leptons,” *Computer Physics Communications*, vol. 64, no. 2, pp. 275 – 299, 1991.
- [18] GRAND Collaboration, Martineau-Huynh, Kotera, Charrier, et al., “Contributions to the 34th International Cosmic Ray Conference (ICRC 2015),” 2015.
- [19] J. Alvarez-Muñiz, W. R. Carvalho, and E. Zas, “Monte Carlo simulations of radio pulses in atmospheric showers using ZHAireS,” *Astroparticle Physics*, vol. 35, pp. 325–341, Jan. 2012.
- [20] A. Aab et al., “Contributions to the 34th International Cosmic Ray Conference (ICRC 2015),” 2015.
- [21] The Pierre Auger Collaboration, A. Aab, P. Abreu, M. Aglietta, and et al., “Combined fit of spectrum and composition data as measured by the Pierre Auger Observatory,” *ArXiv e-prints*, Dec. 2016.
- [22] P. W. ANITA Collaboration, Gorham, P. Allison, B. M. Baughman, et al., “Observational constraints on the ultrahigh energy cosmic neutrino flux from the second flight of the ANITA experiment,” *PRD*, vol. 82, p. 022004, July 2010.
- [23] D. Ardouin, C. Cârloganu, D. Charrier, et al., “First detection of extensive air showers by the TREND self-triggering radio experiment,” *Astroparticle Physics*, vol. 34, pp. 717–731, Apr. 2011.
- [24] A. Aab, P. Abreu, M. Aglietta, and et al., “Probing the radio emission from air showers with polarization measurements,” *PRD*, vol. 89, p. 052002, Mar. 2014.
- [25] K. Murase, S. Inoue, and S. Nagataki, “Cosmic Rays above the Second Knee from Clusters of Galaxies and Associated High-Energy Neutrino Emission,” *ApJ*, vol. 689, p. L105, Dec. 2008.
- [26] K. Fang and K. Murase, “Linking High-Energy Cosmic Particles by Black Hole Jets Embedded in Large-Scale Structures,” *ArXiv e-prints*, Mar. 2017.
- [27] K. Fang, K. Kotera, K. Murase, and A. V. Olinto, “Testing the newborn pulsar origin of ultrahigh energy cosmic rays with EeV neutrinos,” *PRD*, vol. 90, p. 103005, Nov. 2014.
- [28] K. Murase, “Active Galactic Nuclei as High-Energy Neutrino Sources,” *ArXiv e-prints*, Nov. 2015.
- [29] K. Murase, “High energy neutrino early afterglows from gamma-ray bursts revisited,” *Phys. Rev. D*, vol. 76, p. 123001, Dec 2007.
- [30] P. Zarka and F. Mottez, “Observations of Fast Radio Bursts and perspectives at low frequencies,” in *SF2A-2016: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics*.
- [31] Gou, Martineau-Huynh, David, et al., “The GRANDproto35 experiment (ICRC 2017 proceedings),”