

Pgg TMD splitting function in k_T factorization

A. Kusina*

*The H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences,
ul. Radzikowskiego 152, 31-342, Cracow, Poland
E-mail: Aleksander.Kusina@ifj.edu.pl*

M. Hentschinski

*Departamento de Actuaría, Física y Matemáticas, Universidad de las Américas Puebla,
Santa Catarina Martir, 72820 Puebla, Mexico*

K. Kutak

*The H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences,
ul. Radzikowskiego 152, 31-342, Cracow, Poland*

M. Serino

*Department of Physics, Ben Gurion University of the Negev,
Beer Sheva 8410501, Israel*

We calculate the transverse momentum dependent gluon-to-gluon splitting function within k_T -factorization [1], generalizing the framework of [2, 3] and extending our previous works [4, 5] for the quark splittings. While existing versions of k_T factorized evolution equations contain already a gluon-to-gluon splitting function i.e. the leading order BFKL kernel or the CCFM kernel, the obtained splitting function has the important property that it reduces to the leading order BFKL kernel in the low z limit, to the DGLAP gluon-to-gluon splitting function in the collinear limit as well as to the CCFM kernel in the angular ordered region.

*XXVI International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS2018)
16-20 April 2018
Kobe, Japan*

*Speaker.

1. Introduction

Parton distributions functions (PDFs) are crucial elements of collider phenomenology. In presence of a hard scale M with $M \gg \Lambda_{\text{QCD}}$, factorization theorems allow to express cross-sections as convolutions of parton densities (PDFs) and hard matrix elements, where the latter are calculated within perturbative QCD [6]. This was first achieved within the framework of collinear factorization [7–9], where the incoming partons are taken to be collinear with the respective mother hadron. Calculating hard matrix elements to higher orders in the strong coupling constant, one can systematically improve the precision of the theoretical prediction, by incorporating more loops and more emissions of real partons. These extra emissions allow to improve the kinematic approximation inherent to the leading order (LO) description. As an alternative to improving the kinematic description through the calculation of higher order corrections, one may attempt to account for the bulk of kinematic effects already at leading order. An important example of such kinematic effects is the transverse momentum k_T of the initial state partons, which is set to zero within collinear factorization. Schemes which provide an improved kinematic description already at the leading order involve in general Transverse-Momentum-Dependent (TMD) or ‘unintegrated’ PDFs [10]. TMD PDFs arise naturally in regions of phase space characterized by a hierarchy of scales. A particularly interesting example is provided by the so called low x region, where x is the ratio of the hard scale M^2 of the process and the center-of-mass energy squared s . The low x region corresponds therefore to the hierarchy $s \gg M^2 \gg \Lambda_{\text{QCD}}^2$. In such a kinematical setup, large logarithms $\ln 1/x$ can compensate for the smallness of the perturbative strong coupling α_s and it is necessary to resum terms $(\alpha_s \ln 1/x)^n$ to all orders to maintain the predictive power of the perturbative expansion. Such a resummation is achieved by the Balitsky-Fadin-Kuraev-Lipatov (BFKL) [11–14] evolution equation. Its formulation is based on the so called k_T (or high-energy) factorization [15] which is strictly speaking valid in the high energy limit, $s \gg M^2$. In this approach one obtains QCD cross-sections as convolutions (in transverse momentum) of unintegrated gluon density and k_T -dependent perturbative coefficients.

While high energy factorization provides a well defined calculational framework the applicability of the results is naturally limited to the low x limit. If the ensuing formalism is naively extrapolated to intermediate or large x , the framework is naturally confronted with a series of problems and short-comings, e.g. contributions of quarks to the evolution arise as a pure next-to-leading order (NLO) effect and elementary vertices violate energy conservation *i.e.* conservation of the longitudinal momentum fraction. One can account for such effects by including a resummation of terms which restore subleading, but numerically relevant, pieces of the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) [16–18] splitting functions [19–26]. Even though these resummations have been successful in stabilizing low x evolution into the region of intermediate $x \sim 10^{-2}$, extrapolations to larger values of x are still prohibited. Moreover, by merely resumming and calculating higher order corrections within the BFKL formalism, one essentially repeats the program initially outlined for collinear factorization: higher order corrections are calculated to account for kinematic effects which are beyond the regarding factorization scheme. To arrive at a framework which avoids the need to account for kinematic effects through the calculation of higher order corrections, it is therefore necessary to devise a scheme which accounts for both DGLAP (conservation of longitudinal momentum) and BFKL (conservation of transverse momentum) kinematics. Note

that the mere definition of such a scheme is difficult: neither the hard scale of the process (as in DGLAP evolution) nor x (BFKL evolution) provides at first a suitable expansion parameter, if one desires to keep exact kinematics in both variables. To overcome these difficulties, we follow here a proposal initially outlined in [3]. There, the low x resummed DGLAP splitting functions have been constructed following the definition of DGLAP splittings by Curci-Furmanski-Petronzio (CFP) [2]. The authors of [3] were able to define a TMD gluon-to-quark splitting function \tilde{P}_{qg} , both exact in transverse momentum and longitudinal momentum fraction.¹ Following observation of [27] two of us generalized this scheme to calculate the remaining splittings which involve quarks, \tilde{P}_{gq} , \tilde{P}_{qg} and \tilde{P}_{qq} [4]. The computation of the gluon-to-gluon splitting \tilde{P}_{gg} required a further modification of the formalism used in [3, 4] which we did recently in [1]. In this contribution we summarize the most relevant results obtained in this work.

2. Calculation of gluon-to-gluon splitting

The calculation of the \tilde{P}_{gg} splitting function follows the prescription used in our papers [1, 4]. We are working in the high energy kinematics where the momentum of the incoming parton is off-shell (see Fig. 1) and given by: $k^\mu = yp^\mu + k_\perp^\mu$, where p is a light-like momentum defining the direction of a hadron beam and n defines the axial gauge with $n^2 = p^2 = 0$ (that is necessary when using CFP inspired formalism). Additionally, the outgoing momenta is parametrized as: $q^\mu = xp^\mu + q_\perp^\mu + \frac{q_\perp^2 + q^2}{2xp \cdot n} n^\mu$ and we also use: $\tilde{\mathbf{q}} = \mathbf{q} - z\mathbf{k}$ with $z = x/y$.

The TMD splitting function, \tilde{P}_{gg} , is defined as

$$\hat{K}_{gg} \left(z, \frac{\mathbf{k}^2}{\mu^2}, \varepsilon \right) = z \int \frac{d^{2+2\varepsilon} \mathbf{q}}{2(2\pi)^{4+2\varepsilon}} \underbrace{\int dq^2 \mathbb{P}_{g, \text{in}} \otimes \hat{K}_{gg}^{(0)}(q, k) \otimes \mathbb{P}_{g, \text{out}} \Theta(\mu_F^2 + q^2)}_{\tilde{P}_{gg}^{(0)}(z, \mathbf{k}, \tilde{\mathbf{q}}, \varepsilon)}, \quad (2.1)$$

with \mathbb{P}_g being appropriate projection operators and \hat{K}_{gg} the matrix element contributing to the kernel, which at LO is given by the diagram of Fig. 1.

In order to calculate \tilde{P}_{gg} splitting we needed to extend formalism of [3, 4] to the gluon case. This was achieved in [1] by generalizing definition of projector operators and defining appropriate generalized 3-gluon vertex that is gauge invariant in the presence of the off-shell momentum k . Definition of the generalized vertex follows from application of the spinor helicity methods to the high-energy factorization [28–33] and can be obtained by summing the diagrams of Fig. 2, giving:

$$\Gamma_{g^*g^*g}^{\mu_1\mu_2\mu_3}(q, k, p') = \gamma^{\lambda\kappa\mu_3}(-q, k, -p') d^{\mu_1\lambda}(q) d^{\mu_2\kappa}(k) + d^{\mu_1\mu_2}(k) \frac{q^2 n^{\mu_3}}{n \cdot p'} - d^{\mu_1\mu_2}(q) \frac{k^2 p^{\mu_3}}{p \cdot p'}, \quad (2.2)$$

¹Hereafter, we will use the symbol \tilde{P} to indicate a transverse momentum dependent splitting function.




Figure 1: Diagram contributing to the real \tilde{P}_{gg} splitting function at leading order.

with $\mathcal{V}^{\lambda\kappa\mu_3}(-q, k, -p')$ being the ordinary 3-gluon QCD vertex, and $d^{\mu_1\mu_2}(q) = -g^{\mu_1\mu_2} + \frac{q^{\mu_1}n^{\mu_2} + q^{\mu_2}n^{\mu_1}}{q \cdot n}$ is the numerator of the gluon propagator in the light-cone gauge. More details on the exact procedure can be found in [1].




Figure 2: Feynman diagrams contributing to the amplitude used to define the generalized 3-gluon vertex.

The definition of the projectors is more involved. We need to ensure that in the collinear limit the new projectors reduce to the ones introduced by the CFP [2] and that the appropriate high-energy limit for the gluon splitting is also obtained. A natural approach is to modify only the incoming projector $\mathbb{P}_{g,\text{in}}$, as the kinematics of the incoming momentum is more general, and keep the collinear outgoing projector $\mathbb{P}_{g,\text{out}}$ unchanged (as the kinematics of the outgoing momentum is the same). This is what was done by Catani and Hautmann [3, 15] and later by us [4] when defining the projector for the calculation of the quark splitting functions. In that case we used a transverse projector: $k_{\perp}^{\mu}k_{\perp}^{\nu}/k_{\perp}^2$ which, however, due to the more complicated structure of the 3-gluon vertex $\Gamma_{g^*g^*g}^{\mu_1\mu_2\mu_3}$ can not be used any more. Instead we use, a more natural (for high-energy factorization), longitudinal projector given by: $\mathbb{P}_{g,\text{in}}^{\mu\nu} = y^2 p^{\mu} p^{\nu} / k_{\perp}^2$. However, in order to satisfy the requirements of $\mathbb{P}_g^s = \mathbb{P}_{g,\text{in}} \mathbb{P}_{g,\text{out}}$ being a projector operator ($\mathbb{P}_g^s \otimes \mathbb{P}_g^s = \mathbb{P}_g^s$) we also need to modify the outgoing projector. In the end we end up with the following gluon projectors:

$$\mathbb{P}_{g,\text{in}}^{\mu\nu} = -y^2 \frac{p^{\mu} p^{\nu}}{k_{\perp}^2}, \quad \mathbb{P}_{g,\text{out}}^{\mu\nu} = -g^{\mu\nu} + \frac{k^{\mu} n^{\nu} + k^{\nu} n^{\mu}}{k \cdot n} - k^2 \frac{n_{\mu} n_{\nu}}{(k \cdot n)^2}. \quad (2.3)$$

It is easy to check that now $\mathbb{P}_g^s \otimes \mathbb{P}_g^s = \mathbb{P}_g^s$ holds. Also the new outgoing projector is consistent with the collinear case and one can show that in the collinear limit the difference between $y^2 p^{\mu} p^{\nu} / k_{\perp}^2$ and $k_{\perp}^{\mu} k_{\perp}^{\nu} / k_{\perp}^2$ vanishes when contracted into the relevant vertices. More details are provided in [1].

3. Results

Using the elements introduced in the previous section we can compute the transverse momentum dependent gluon-to-gluon splitting function. We present here only the final result:

$$\tilde{P}_{gg}^{(0)}(z, \tilde{\mathbf{q}}, \mathbf{k}) = 2C_A \left\{ \frac{\tilde{\mathbf{q}}^4}{(\tilde{\mathbf{q}} - (1-z)\mathbf{k})^2 [\tilde{\mathbf{q}}^2 + z(1-z)\mathbf{k}^2]} \left[\frac{z}{1-z} + \frac{1-z}{z} + (3-4z) \frac{\tilde{\mathbf{q}} \cdot \mathbf{k}}{\tilde{\mathbf{q}}^2} + z(3-2z) \frac{\mathbf{k}^2}{\tilde{\mathbf{q}}^2} \right] + \frac{(1+\varepsilon)\tilde{\mathbf{q}}^2 z(1-z)[2\tilde{\mathbf{q}} \cdot \mathbf{k} + (2z-1)\mathbf{k}^2]^2}{2\mathbf{k}^2 [\tilde{\mathbf{q}}^2 + z(1-z)\mathbf{k}^2]^2} \right\}, \quad (3.1)$$

or after angular averaging (and setting $\varepsilon = 0$):

$$\begin{aligned} \bar{P}_{gg}^{(0)}\left(z, \frac{\mathbf{k}^2}{\tilde{\mathbf{q}}^2}\right) = C_A \frac{\tilde{\mathbf{q}}^2}{\tilde{\mathbf{q}}^2 + z(1-z)\mathbf{k}^2} & \left[\frac{(2-z)\tilde{\mathbf{q}}^2 + (z^3 - 4z^2 + 3z)\mathbf{k}^2}{z(1-z)|\tilde{\mathbf{q}}^2 - (1-z)^2\mathbf{k}^2|} \right. \\ & \left. + \frac{(2z^3 - 4z^2 + 6z - 3)\tilde{\mathbf{q}}^2 + z(4z^4 - 12z^3 + 9z^2 + z - 2)\mathbf{k}^2}{(1-z)(\tilde{\mathbf{q}}^2 + z(1-z)\mathbf{k}^2)} \right]. \end{aligned} \quad (3.2)$$

Now we can explicitly check the corresponding kinematic limits. In the collinear case this is straightforward, since the transverse integral in Eq. (2.1) is specially adapted for this limit. In particular, one easily obtains the real part of the DGLAP gluon-to-gluon splitting function:

$$\lim_{\mathbf{k}^2 \rightarrow 0} \bar{P}_{ij}^{(0)}\left(z, \frac{\mathbf{k}^2}{\tilde{\mathbf{q}}^2}\right) = 2C_A \left[\frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right]. \quad (3.3)$$

In order to study the high-energy and soft limit it is convenient to change to the following variables: $\tilde{\mathbf{p}} = \frac{\mathbf{k}-\mathbf{q}}{1-z} = \mathbf{k} - \frac{\tilde{\mathbf{q}}}{1-z}$, then in the high-energy limit ($z \rightarrow 0$) we obtain:

$$\lim_{z \rightarrow 0} \hat{K}_{gg}\left(z, \frac{\mathbf{k}^2}{\mu^2}, \varepsilon, \alpha_s\right) = \frac{\alpha_s C_A}{\pi(e^{\gamma_E} \mu^2)^\varepsilon} \int \frac{d^{2+2\varepsilon} \tilde{\mathbf{p}}}{\pi^{1+\varepsilon}} \Theta(\mu_F^2 - (\mathbf{k} - \tilde{\mathbf{p}})^2) \frac{1}{\tilde{\mathbf{p}}^2}$$

where the term under the integral is easily identified as the real part of the LO BFKL kernel. Additionally, one can check that in the angular ordered region of phase space, where $\tilde{\mathbf{p}}^2 \rightarrow 0$, we reproduce the real/unresummed part of the CCFM kernel [34–36]: $\frac{1}{z} + \frac{1}{1-z} + \mathcal{O}\left(\frac{\tilde{\mathbf{p}}^2}{\mathbf{k}^2}\right)$.

4. Summary

The main result of this paper is the calculation of a transverse momentum dependent gluon-to-gluon splitting function. The splitting function reduces both to the conventional gluon-to-gluon DGLAP splitting in the collinear limit as well as to the LO BFKL kernel in the low x /high energy limit; moreover the CCFM gluon-to-gluon splitting function is re-obtained in the limit where the transverse momentum of the emitted gluon vanishes, *i.e.* if the emitted gluon is soft. The derivation of this result is based on the Curci-Furmanski-Petronzio formalism for the calculation of DGLAP splitting functions in axial gauges. To address gauge invariance in the presence of off-shell partons, high energy factorization adapted for axial gauges has been used to derive an effective production vertex which then could be shown to satisfy current conservation. The next step in completing the calculation of TMD splitting functions is the determination of the still missing virtual corrections. With the complete set of splitting functions at hand, it will be finally possible to formulate an evolution equation for the unintegrated (TMD) parton distribution functions including both gluons and quarks.

References

- [1] M. Hentschinski, A. Kusina, K. Kutak, and M. Serino, ‘‘TMD splitting functions in k_T factorization: the real contribution to the gluon-to-gluon splitting,’’ *Eur. Phys. J.* **C78** (2018), no. 3, 174, [1711.04587](https://arxiv.org/abs/1711.04587).
- [2] G. Curci, W. Furmanski, and R. Petronzio, ‘‘Evolution of Parton Densities Beyond Leading Order: The Nonsinglet Case,’’ *Nucl. Phys.* **B175** (1980) 27–92.

- [3] S. Catani and F. Hautmann, “High-energy factorization and small x deep inelastic scattering beyond leading order,” *Nucl. Phys.* **B427** (1994) 475–524, [hep-ph/9405388](#).
- [4] O. Gituliar, M. Hentschinski, and K. Kutak, “Transverse-momentum-dependent quark splitting functions in k_T -factorization: real contributions,” *JHEP* **01** (2016) 181, [1511.08439](#).
- [5] M. Hentschinski, A. Kusina, and K. Kutak, “Transverse momentum dependent splitting functions at work: quark-to-gluon splitting,” *Phys. Rev.* **D94** (2016), no. 11, 114013, [1607.01507](#).
- [6] J. C. Collins, *Renormalization*, vol. 26 of *Cambridge Monographs on Mathematical Physics*. Cambridge University Press, Cambridge, 1986.
- [7] R. K. Ellis, H. Georgi, M. Machacek, H. D. Politzer, and G. G. Ross, “Perturbation Theory and the Parton Model in QCD,” *Nucl. Phys.* **B152** (1979) 285–329.
- [8] J. C. Collins, D. E. Soper, and G. Sterman, “Factorization for short distance hadron-hadron scattering,” *Nucl. Phys. B* **261** (jan, 1985) 104–142.
- [9] G. T. Bodwin, “Factorization of the Drell-Yan Cross-Section in Perturbation Theory,” *Phys.Rev.* **D31** (1985) 2616.
- [10] R. Angeles-Martinez *et al.*, “Transverse Momentum Dependent (TMD) parton distribution functions: status and prospects,” *Acta Phys. Polon.* **B46** (2015), no. 12, 2501–2534, [1507.05267](#).
- [11] V. Fadin, E. Kuraev, and L. Lipatov, “On the pomeranchuk singularity in asymptotically free theories,” *Phys. Lett. B* **60** (dec, 1975) 50–52.
- [12] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, “Multi - Reggeon Processes in the Yang-Mills Theory,” *Sov. Phys. JETP* **44** (1976) 443–450. [*Zh. Eksp. Teor. Fiz.*71,840(1976)].
- [13] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, “The Pomeranchuk Singularity in Nonabelian Gauge Theories,” *Sov.Phys.JETP* **45** (1977) 199–204.
- [14] I. I. Balitsky and L. N. Lipatov, “The Pomeranchuk Singularity in Quantum Chromodynamics,” *Sov.J.Nucl.Phys.* **28** (1978) 822–829.
- [15] S. Catani, M. Ciafaloni, and F. Hautmann, “High-energy factorization and small x heavy flavor production,” *Nucl. Phys.* **B366** (1991) 135–188.
- [16] V. N. Gribov and L. N. Lipatov, “Deep inelastic $e p$ scattering in perturbation theory,” *Sov. J. Nucl. Phys.* **15** (1972) 438–450. [*Yad. Fiz.*15,781(1972)].
- [17] G. Altarelli and G. Parisi, “Asymptotic Freedom in Parton Language,” *Nucl. Phys.* **B126** (1977) 298–318.
- [18] Y. L. Dokshitzer, “Calculation of the Structure Functions for Deep Inelastic Scattering and $e^+ e^-$ Annihilation by Perturbation Theory in Quantum Chromodynamics.,” *Sov. Phys. JETP* **46** (1977) 641–653. [*Zh. Eksp. Teor. Fiz.*73,1216(1977)].
- [19] G. P. Salam, “A Resummation of large subleading corrections at small x ,” *JHEP* **07** (1998) 019, [hep-ph/9806482](#).
- [20] G. Altarelli, R. D. Ball, and S. Forte, “Perturbatively stable resummed small x evolution kernels,” *Nucl. Phys.* **B742** (2006) 1–40, [hep-ph/0512237](#).
- [21] M. Ciafaloni, D. Colferai, G. P. Salam, and A. M. Stasto, “The Gluon splitting function at moderately small x ,” *Phys. Lett.* **B587** (2004) 87–94, [hep-ph/0311325](#).

- [22] A. Sabio Vera, “An “all-poles” approximation to collinear resummations in the Regge limit of perturbative QCD,” *Nucl. Phys. B* **722** (2005), no. 1-2, 65–80, [0505128](#).
- [23] M. Hentschinski, A. Sabio Vera, and C. Salas, “Hard to Soft Pomeron Transition in Small- x Deep Inelastic Scattering Data Using Optimal Renormalization,” *Phys. Rev. Lett.* **110** (2013), no. 4, 041601, [1209.1353](#).
- [24] M. Hentschinski, A. Sabio Vera, and C. Salas, “ F_2 and F_L at small x using a collinearly improved BFKL resummation,” *Phys. Rev.* **D87** (2013), no. 7, 076005, [1301.5283](#).
- [25] J. Kwiecinski, A. D. Martin, and A. M. Stasto, “A Unified BFKL and GLAP description of F_2 data,” *Phys. Rev.* **D56** (1997) 3991–4006, [hep-ph/9703445](#).
- [26] M. Bonvini, S. Marzani, and C. Muselli, “Towards parton distribution functions with small- x resummation: HELL 2.0,” *Submitted to: JHEP* (2017) [1708.07510](#).
- [27] F. Hautmann, M. Hentschinski, and H. Jung, “Forward Z-boson production and the unintegrated sea quark density,” *Nucl. Phys.* **B865** (2012) 54–66, [1205.1759](#).
- [28] A. van Hameren, P. Kotko, and K. Kutak, “Multi-gluon helicity amplitudes with one off-shell leg within high energy factorization,” *JHEP* **12** (2012) 029, [1207.3332](#).
- [29] A. van Hameren, P. Kotko, and K. Kutak, “Helicity amplitudes for high-energy scattering,” *JHEP* **01** (2013) 078, [1211.0961](#).
- [30] A. van Hameren, K. Kutak, and T. Salwa, “Scattering amplitudes with off-shell quarks,” *Phys. Lett.* **B727** (2013) 226–233, [1308.2861](#).
- [31] A. van Hameren, “BCFW recursion for off-shell gluons,” *JHEP* **07** (2014) 138, [1404.7818](#).
- [32] A. van Hameren and M. Serino, “BCFW recursion for TMD parton scattering,” *JHEP* **07** (2015) 010, [1504.00315](#).
- [33] K. Kutak, A. van Hameren, and M. Serino, “QCD amplitudes with 2 initial spacelike legs via generalised BCFW recursion,” *JHEP* **02** (2017) 009, [1611.04380](#).
- [34] M. Ciafaloni, “Coherence Effects in Initial Jets at Small q^2/s ,” *Nucl. Phys.* **B296** (1988) 49–74.
- [35] S. Catani, F. Fiorani, and G. Marchesini, “QCD Coherence in Initial State Radiation,” *Phys. Lett.* **B234** (1990) 339–345.
- [36] G. Marchesini, “QCD coherence in the structure function and associated distributions at small x ,” *Nucl. Phys. B* **445** (jul, 1995) 49–78.