

The full Schwinger-Dyson tower for random tensor models

Carlos Ignacio Pérez Sánchez* [†]

*Mathematisches Institut der Westfälischen Wilhelms-Universität,
Einsteinstraße 62, 48149 Münster, Germany
E-mail: perezsan@uni-muenster.de*

We treat random rank- D tensor models as D -dimensional quantum field theories—tensor field theories (TFT)—and review some of their non-perturbative methods. We classify the correlation functions of complex tensor field theories by boundary graphs, sketch the obtention of the Ward-Takahashi identity and stress its relevance in the derivation of the tower of exact, analytic Schwinger-Dyson equations for all the correlation functions (with connected boundary) of TFTs with quartic pillow-like interactions.

*Corfu Summer Institute 2017 "School and Workshops on Elementary Particle Physics and Gravity"
2-28 September 2017
Corfu, Greece*

*Speaker.

[†]The author's participation in the Corfu 2017 summer school and workshop was thanks to COST Action MP1405 Quantum Structure of Spacetime (QSpace). The author wishes to thank the Corfu Summer Institute 2017 hospitality. The author thanks George Zoupanos and for the invitation to write this note. The Collaborative Research Center "Groups, Geometry & Actions", SFB 878 (University of Münster), is acknowledged for financial support. Discussions with Romain Pascalie, Johannes Thürigen and Raimar Wulkenhaar were fruitful for this review.

- V can be any of $\{c\text{□}c\}_{c=1,2,3}$ or $V = \text{◇}$ for real or $O(N)$ tensor models initiated by Carrozza and Tanasă [17], and used by Klebanov and Tarnopolsky [12] in the context of the SYK-like tensor models
- or $V = c\text{□}c$ for $c = 1, 2, 3$ for complex or $U(N)$ tensor models. The black-white bipartiteness reflects the presence of both the tensor field and its conjugate. Rather $U(N)$ -invariant theories are the TFT we shall deal here with.

Higher-point functions follow an even more complicated schema. Either way, accordingly, the Green’s functions need further specification and, in fact, the classification of the correlation functions for higher-rank theories is the following:

- for real matrix theories, there are as many connected k -point functions [1] as integer partitions of k . Then, there are three 3-point functions, five 4-point functions, seven 5-point functions, and so on.
- for complex tensor models the connected k -point functions are classified by (possibly disconnected) D -colored² graphs in k vertices [16]. In particular $k = 2\ell$ should be even. Each edge in these graphs is of certain color a , and this enforces momentum-transmission³ of this very color. Therefore, the Feynman graph structures with four legs can encode momentum-transmission according to $1\text{□}1$, $2\text{□}2$, $3\text{□}3$ or (as pictured in Fig. 1) $\text{□} \sqcup \text{□}$.

In particular, in order to obtain the analogue of eq. (1.1) in a D -dimensional QFT-context, say, for quartic tensor field theories of rank- D , one needs to specify which of the four 4-point function we are inserting into the 2-point function. The aim of this paper is to explain how to achieve this and to arrive at analytic SDE for every (connected) correlation function. The methods exposed here are based on [16] and [18].




Figure 1: Connected 4-point φ_3^4 -Feynman graph with disconnected boundary. The dashed (or 0 color) transmits simultaneously the three colors. Hence, momentum transmission is encoded by $0a$ -bicolored paths between external legs. This draws a graph, the *boundary graph*, which in this case is disconnected. Thus, this graph contributes to the 4-point function $G_{m|m}^{(4)}$ or $G_{\text{discn.}}^{(4)}$. (see also Table 1 for notation).

Remark. We do not use Einstein’s implicit summation notation.

2. The strategy

The idea of the utilization of a matrix Ward identity (based on the Ward identity [19]) in order to derive the SDE of matrix models is due to Grosse and Wulkenhaar [1, 20].


²This is a common abbreviation in the tensor model jargon, for “vertex-bipartite regularly edge- D -colored graphs”.

³We call the index that is transmitted “momentum” because these models are originated in certain Group Field Theory context, whose Fourier dual has the structure of a TFT; if one interpret the group manifolds as direct space, then the indices are the momenta.

2.1 Complex tensor models

Complex tensor models, colored tensor models and random tensor models [7] study fields $\varphi_{\mathbf{x}} \in \mathbb{C}$ ($\mathbf{x} \in I_1 \times \dots \times I_D \subset \mathbb{Z}^D$) whose indices⁴ transform independently under elements of $U(N)$ of the product group $H_D = U(N)^D$. This means that

$$\varphi_{x_1 \dots x_D} \mapsto \varphi'_{x_1 \dots x_D} = \sum_{y_a} [W_a]_{x_a y_a} \varphi_{x_1 \dots y_a \dots x_D}, \quad \bar{\varphi}_{x_1 \dots x_D} \mapsto \bar{\varphi}'_{x_1 \dots x_D} = \sum_{y_a} [\bar{W}_a]_{x_a y_a} \bar{\varphi}_{x_1 \dots y_a \dots x_D},$$

for each W_a in the a -th factor $U(N)$ of H_D , for any $a = 1, \dots, D$. Each of the factors (and of the location of the tensor indices) is referred to as a *color*⁵. Interactions of this kind of theories are H_D -invariants. We restrict to models for which any (graph)-vertex lies on a subgraph of the type  for certain color a . For $D = 3$, this constrains the interactions of models to the list

$$\text{pillow}_a, \text{pillow}_a^2, \text{pillow}_a^3, \text{pillow}_a^4, \dots \text{ (for any color } a\text{)}. \quad (2.1)$$

Other type of interactions need another methods. The origin of this restriction is technical and will be explained in Section 2.2. Here, we treat models with pillow-like interactions, but otherwise without any restriction in their rank. *Pillows* are melonic⁶ graphs of four vertices. These are $\{1\text{pillow}_1, 2\text{pillow}_2, 3\text{pillow}_3\}$, for rank-3 models, $\{1\text{pillow}_1, 2\text{pillow}_2, 3\text{pillow}_3, 4\text{pillow}_4\}$ for rank-4 models, etc. When the rank is clear, we denote by V_a the pillow with preferred color a (e.g. $1\text{pillow}_1 = V_1$).


2.2 The usefulness of the Ward-Takahashi Identity

We consider the quartic tensor model with interaction $S_{\text{int}} = \lambda \sum_{c=1}^D V_c$ with a kinetic Laplacian-like kernel E , which possibly breaks the H_D -symmetry in the quadratic invariant $\text{Tr}_2(\bar{\varphi}, \varphi) \rightarrow S_0 = \text{Tr}_2(\bar{\varphi}, E\varphi)$. Functional integration of the partition function yields

$$Z[J, \bar{J}; E] = C \exp\left(-\lambda \sum_{a=1}^3 V_a\right) \Bigg|_{(\varphi, \bar{\varphi}) \rightarrow (\frac{\delta}{\delta J}, \frac{\delta}{\delta \bar{J}})} Z^{\text{free}}[J, \bar{J}; E].$$

Any analytic Schwinger-Dyson equation begins by deriving with respect to the sources. By deriving with respect to \bar{J}_s we get [16, 18]

$$\frac{\delta \log Z[J, \bar{J}; E]}{\delta \bar{J}_s} = \frac{1}{E_s} \left\{ J_s - \frac{1}{Z[J, \bar{J}; E]} \left(\frac{\partial S_{\text{int}}(\varphi, \bar{\varphi})}{\partial \bar{\varphi}_s} \right) \Bigg|_{(\varphi, \bar{\varphi}) \rightarrow (\frac{\delta}{\delta J}, \frac{\delta}{\delta \bar{J}})} Z[J, \bar{J}; E] \right\}. \quad (2.2)$$

By assumption (Sec. 2.1), the term in round parenthesis contains, after evaluation of the sources, the subgraph , and thus a derivative of the form

$$\mathcal{L}_{m_a n_a} \equiv \sum_{\mathbf{p}_a \in \mathbb{Z}} \left[\frac{\delta^2 Z[J, \bar{J}; E]}{\delta J_{p_1 \dots p_{a-1} m_a p_{a+1} \dots p_D} \delta \bar{J}_{p_1 \dots p_{a-1} n_a p_{a+1} \dots p_D}} \right] \quad (2.3)$$

⁴We think of the large- N limit, so we write, instead of I_a , directly \mathbb{Z} .

⁵For historical reasons [21].

⁶That is, with vanishing Gurău-degree [7], but this is concept is not essential here because the present results entail no $1/N$ -truncation.

acted on by more derivatives. This term resembles the LHS appearing in the Ward-Takahashi Identity (WTI)

$$\sum_{\mathbf{p}_{\hat{a}} \in \mathbb{Z}} \left[\frac{\delta^2 Z[J, \bar{J}; E]}{\delta J_{p_1 \dots p_{a-1} m_a p_{a+1} \dots p_D} \delta \bar{J}_{p_1 \dots p_{a-1} n_a p_{a+1} \dots p_D}} \times E_{m_a n_a; \mathbf{p}_{\hat{a}}} \right] = D_{J, \bar{J}} Z[J, \bar{J}; E] \quad (2.4)$$

obtained by Ousmane-Samary [22] from the $U(N)$ -invariance (this group being the a -th factor of H_D) of the path integral. Here, $E_{m_a n_a; \mathbf{p}_{\hat{a}}} \equiv E_{p_1 \dots m_a \dots p_D} - E_{p_1 \dots n_a \dots p_D}$, $\mathbf{p}_{\hat{a}} \equiv (p_1, \dots, p_{a-1}, p_{a+1}, \dots, p_D)$ and $D_{J, \bar{J}}$ is a first order differential operator in the sources. It would be useful to reduce the derivatives by using this WTI; however, it also implies the difference of the kernels. We restrict, therefore, to models that satisfy that:

$$\text{for any color } a, E_{m_a n_a; \mathbf{q}_{\hat{a}}} = E_{q_1 \dots m_a \dots q_D} - E_{q_1 \dots n_a \dots q_D} \text{ does not depend on } \mathbf{q}_{\hat{a}}. \quad (\diamond)$$

We thus write only $E_{m_a n_a; \mathbf{p}_{\hat{a}}} =: E_{m_a n_a}$ from now on. The condition \diamond allows one to get the term $E_{m_a n_a}$ out of the sum and solve for $\mathcal{L}_{m_a n_a}$. We remark that the non-triviality of this task relies on the skew-symmetry of the indices of $E_{m_a n_a}$. This means that we need to find the term that is proportional to $\delta_{m_a n_a}$ in $\mathcal{L}_{m_a n_a}$ (see eq. (2.3)). This is a functional that we denote by $Y_{m_a}^{(a)}[J, \bar{J}]$ (and name, sloppily, Y -term). After complete knowledge about this Y -term has been obtained we say the WTI is *full*. The first full WTI was found for $O(N)$ matrix models [1, Sec. 2].

3. The full Ward-Takahashi identity

For arbitrary rank- D $U(N)$ tensor models the full WTI reads [16]:

$$\mathcal{L}_{m_a n_a} = \left(\delta_{m_a n_a} Y_{m_a}^{(a)}[J, \bar{J}] \right) \cdot Z[J, \bar{J}; E] + \sum_{\mathbf{p}_{\hat{a}} \in \mathbb{Z}^{D-1}} \frac{1}{E_{m_a n_a}} \left(\bar{J}_{m_a \mathbf{p}_{\hat{a}}} \frac{\delta}{\delta \bar{J}_{n_a \mathbf{p}_{\hat{a}}}} - J_{n_a \mathbf{p}_{\hat{a}}} \frac{\delta}{\delta J_{m_a \mathbf{p}_{\hat{a}}}} \right) Z[J, \bar{J}; E].$$

In the next subsections, we explain how to define the correlation functions, and, subsequently, how to obtain the Y -term.

3.1 The expansion of the free energy in boundary graphs

The (connected) correlation functions of TFTs will be defined as derivatives with respect to sources, as in usual QFT. Nevertheless, the naive Ansatz

$$\log Z[J, \bar{J}] \stackrel{?}{=} \sum_{\mathbf{x}, \mathbf{y}} G^{(2)}(\mathbf{x}, \mathbf{y}) J_{y_1 y_2 y_3} \bar{J}_{x_1 x_2 x_3} + \sum_{\mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}} G^{(4)}(\mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}) J_{x_1 x_2 x_3} \bar{J}_{w_1 w_2 w_3} J_{y_1 y_2 y_3} \bar{J}_{z_1 z_2 z_3} + \mathcal{O}(4)$$

is, due to the color structure, an oversimplification that impairs the derivation of analytic Schwinger-Dyson equations for the thus defined $G^{(2\ell)}$ -functions. The right expansion takes into account the

transmission of momentum inside classes of Feynman graphs, and is given by

$$\begin{aligned} \log Z[J, \bar{J}] &= \sum_{\mathbf{x}} G^{(2)}(\mathbf{x}) J_{x_1 x_2 x_3} \bar{J}_{x_1 x_2 x_3} + \sum_{\mathbf{x}, \mathbf{y}} G_{\text{disconn.}}^{(4)}(\mathbf{x}, \mathbf{y}) J_{x_1 x_2 x_3} \bar{J}_{x_1 x_2 x_3} J_{y_1 y_2 y_3} \bar{J}_{y_1 y_2 y_3} \\ &+ \sum_{\mathbf{x}, \mathbf{y}} G_1^{(4)}(\mathbf{x}, \mathbf{y}) J_{y_1 y_2 y_3} \bar{J}_{x_1 y_2 y_3} J_{x_1 x_2 x_3} \bar{J}_{y_1 x_2 x_3} + \sum_{\mathbf{x}, \mathbf{y}} G_2^{(4)}(\mathbf{x}, \mathbf{y}) J_{y_1 y_2 y_3} \bar{J}_{y_1 x_2 y_3} J_{x_1 x_2 x_3} \bar{J}_{x_1 y_2 x_3} \\ &+ \sum_{\mathbf{x}, \mathbf{y}} G_3^{(4)}(\mathbf{x}, \mathbf{y}) J_{y_1 y_2 y_3} \bar{J}_{x_1 x_2 y_3} J_{x_1 x_2 x_3} \bar{J}_{y_1 y_2 x_3} + \mathcal{O}(6) \end{aligned} \quad (3.1)$$

The sub-indices of the 4-point functions $G^{(4)}$ will be clear soon. This expansion can be conveniently (compactly) organized by, again, colored graphs. This would yield an algorithmic derivation of the Y -term. Notice the effect of the the double derivative on the source-term

$$\sum_{\mathbf{p}_a} \frac{\delta}{\delta J_{\mathbf{m}_a \mathbf{p}_a}} \frac{\delta}{\delta \bar{J}_{\mathbf{n}_a \mathbf{p}_a}} \sum_{\mathbf{x}^1, \dots, \mathbf{y}, \dots, \mathbf{x}^\ell} G^{(2\ell)}(\mathbf{x}^1, \dots, \mathbf{y}, \dots, \mathbf{x}^\ell) [J_{\mathbf{x}^1} \cdots J_{\mathbf{y}} \cdots J_{\mathbf{x}^\ell}] \cdot [\bar{J}_{\dots} \cdots \bar{J}_{\dots y_a \dots} \cdots \bar{J}_{\dots}]. \quad (3.2)$$

Namely, from this equation it is clear that $Y_{m_a}^{(a)}$ has contributions from ‘hitting’ two sources J and \bar{J} that are connected by an a -colored edge in the boundary graph (in this term, as a matter of fact, $\bar{J}_{\dots y_a \dots}$ and $J_{\mathbf{y}}$). In order to understand this, we find convenient to recall what happens in the for matrix field theories. If this is superfluous for the reader, Sec. 3.1.1 could be skipped.

3.1.1 The free energy for real matrix models

As pointed out in the introduction, the correlation functions (the momenta of the free energy $\log Z_{\text{matrix}}$) of a general real matrix model are classified by integer partitions of k . If these partitions are indexed by $\alpha \in \{1, \dots, P(k)\}$, the free energy is expanded as

$$\log Z_{\text{matrix}}[J] = \sum_{k=1}^{\infty} \sum_{\alpha=1}^{P(k)} \frac{1}{\sigma(\alpha)} G_{\alpha}^{(k)} \star \mathbf{j}(\alpha) = \text{‘sum over triangulation of boundaries (circles)’}. \quad (3.3)$$

This is shorthand but is not a formal expression. An integer partition $\alpha = (n_1, \dots, n_{B(\alpha)}) \in \mathbb{Z}_{\geq 0}^k$ (i.e. $k = \sum_{r=1}^{B(\alpha)} n_r$ with $n_{B(\alpha)} \neq 0$, and $n_i < n_j$ if $i < j$) determines $B(\alpha)$ boundaries, n_r of which carry r sources attached. Thus, $(\mathbf{j}(\alpha))(\mathbf{p}^1, \dots, \mathbf{p}^{B(\alpha)}) := \prod_{r=1}^{B(\alpha)} J_{p_r^1 p_r^2} J_{p_r^2 p_r^3} \cdots J_{p_r^{n_r} p_r^1}$ and $\mathbf{p}^r = (p_r^1, \dots, p_r^{n_r})$. The star, \star , point-wise sums the product $(G_{\alpha}^{(k)} \cdot \mathbf{j}(\alpha))$ is over the arguments $(\mathbf{p}^1, \dots, \mathbf{p}^{B(\alpha)})$. Further, $\sigma(\alpha)$ is a symmetry factor.

The topological significance of this expansion is clear: Equation (3.3) is a sum over triangulations of the boundary of the surfaces that the ribbon graphs triangulate. That is, α determines for n_r circles a precise ‘triangulation by r intervals’. Based on this, one can derive the free energy for complex tensor models. The useful concept there is that of a *boundary graph*.

3.1.2 The free energy of complex tensor models

The expansion of the free energy also has a geometrical meaning. It is an expansion over all triangulations of boundaries, but in higher dimensions. For $D = 3$, these are triangulations of closed, orientable surfaces. In fact, these are triangulable by bipartite 3-colored graphs and the sum turns out to be over all the boundary graph that are triangulated by a particular model S_{int} . The

CORRELATION FUNCTIONS			CORRELATION FUNCTIONS		
Order	Graph notation	Simplified notation	Order	Graph notation	Simplified notation
2-pt function	$G_{\ominus}^{(2)}$	$G^{(2)}$	6-pt functions	$G_{\text{Q}_a}^{(6)}$	$G_{Q_a}^{(6)}$
4-pt functions	$G_{a a}^{(4)}$	$G_{V_a}^{(4)}$		$G_{K_{3,3}}^{(6)}$	$G_{K_{3,3}}^{(6)}$
	$G_{ \ominus \ominus}^{(4)}$	$G_{m m}^{(4)}$		$G_{F_{a;bc}}^{(6)}$	$G_{F_{a;bc}}^{(6)}$
				$G_{ \ominus a a}^{(6)}$	$G_{m V_a}^{(6)}$
				$G_{ \ominus \ominus \ominus}^{(6)}$	$G_{m m m}^{(6)}$

Table 1: Correlation functions of the ϕ_3^4 -theory until order six

boundaries are characterized in [16]. The general expansion for any rank reads:

$$\log Z_{\text{tensors}}[J, \bar{J}; E] = \sum_{\ell=1}^{\infty} \sum_{\substack{\mathcal{B} \text{ is boundary of } S_{\text{int}} \\ \#\text{Vertices}(\mathcal{B})=2\ell}} \frac{1}{|\text{Aut}_c(\mathcal{B})|} G_{\mathcal{B}}^{(2\ell)} \star \mathbf{j}(\mathcal{B}). \quad (3.4)$$

The elements of this formula are:

- to each boundary graph \mathcal{B} , \mathbf{j} associates a function $\mathbf{j}(\mathcal{B}) : M_{D \times \ell}(\mathbb{Z}) \rightarrow \mathbb{C}$ in the sources given by

$$(\mathbf{j}(\mathcal{B}))(\mathbf{X}) \equiv \prod_{\alpha=1}^{\ell} J_{\mathbf{x}^{\alpha}} \bar{J}_{\mathbf{y}^{\alpha}(\mathbf{X})}, \text{ being } \mathbf{x}^{\alpha}, (\alpha = 1, \dots, \ell) \text{ the columns of } \mathbf{X}, \quad (3.5)$$

and $\{\mathbf{y}^{\alpha}(\mathbf{X})\}_{\alpha}$ is the set of (unordered) momenta \mathbf{y}^{α} that one gets in the \bar{J} -sources at the external legs of a Feynman graph \mathcal{G} with $\partial\mathcal{G} = \mathcal{B}$ by ‘injecting momenta the \mathbf{x}^{γ} ’ ($\gamma = 1, \dots, \ell$) at the external legs marked by J -sources (see Fig. 1). We choose the notation $\mathcal{B}_*(\mathbf{X}) = (\mathbf{y}^1, \dots, \mathbf{y}^{\ell})$ (see [16] for the detailed construction). For instance⁷, $\mathbf{j}(1\text{Q}_1)(\mathbf{x}, \mathbf{z}) = J_{z_1 z_2 z_3} \bar{J}_{x_1 z_2 z_3} J_{x_1 x_2 x_3} \bar{J}_{z_1 x_2 x_3}$, since $(1\text{Q}_1)_*(\mathbf{x}, \mathbf{z}) = (x_1, z_2, z_3, z_1, x_2, x_3)$

- one then sums the product $G_{\mathcal{B}}^{(2\ell)}(\mathbf{X}) \cdot (\mathbf{j}(\mathcal{B}))(\mathbf{X})$ over all momenta $\mathbf{X} \in M_{D \times \ell}(\mathbb{Z})$; the star \star abbreviates this sum
- finally, one divides by the order of the automorphism group $\text{Aut}_c(\mathcal{B})$ of the graph \mathcal{B} . The automorphism group will be important in the following section.

Since from now on we work only with TFTs, we omit the subindex ‘tensors’ in the partition function Z_{tensors} . For rank-3 models, the most general expansion is

$$\begin{aligned} \log Z[J, \bar{J}] = & G^{(2)} \star \mathbf{j}(m) + \frac{1}{2!} G_{m|m}^{(4)} \star \mathbf{j}(m|m) + \frac{1}{2} \sum_{c=1}^3 \left[G_{V_c}^{(4)} \star \mathbf{j}(V_c) + \frac{1}{3} G_{Q_c}^{(6)} \star \mathbf{j}(Q_c) + G_{F_c}^{(6)} \star \mathbf{j}(F_c) \right. \\ & \left. + \frac{1}{2} G_{m|V_c}^{(6)} \star \mathbf{j}(m|V_c) \right] + \frac{1}{3} G_{K_{3,3}}^{(6)} \star \mathbf{j}(K_{3,3}) + \frac{1}{3!} G_{m|m|m}^{(6)} \star \mathbf{j}(m|m|m) + \mathcal{O}(8). \end{aligned} \quad (3.6)$$

⁷Recall that we do not use Einstein’s sum convention.

As shown in [16] (relying on [23]) for rank- D models with all⁸ the pillows $S_{\text{int}} = \lambda \sum_{c=1}^D V_c$, for any D -colored graph \mathcal{B} , even if \mathcal{B} is disconnected, $G_{\mathcal{B}}^{(2\ell)} \neq 0$ holds. Table 1 shows the transition from the original source [16] notation to the compact one used here.

3.2 Graph calculus

The free energy is generated by graphs. Since this is not a formal expansion, a tool should be developed in order to read off the coefficients (functions) of the graphs. This is the graph calculus [16], which consists in deriving functionals $\mathcal{A}[J, \bar{J}]$ with respect to $(\mathbf{j}(\mathcal{B}))(\mathbf{X})$, where \mathcal{B} is a boundary graph, and by momenta $\mathbf{X} = (\mathbf{x}^1, \dots, \mathbf{x}^\ell)$ ($2\ell = \#$ vertices of \mathcal{B}). We restrict \mathbf{X} to

$$\mathcal{F}_{D,\ell} := \{(\mathbf{x}^1, \dots, \mathbf{x}^\ell) \in M_{D \times \ell}(\mathbb{Z}) \mid x_c^\alpha \neq x_c^\nu \text{ if } \alpha \neq \nu (\alpha, \nu = 1, \dots, \ell), \text{ for all } c = 1, \dots, D\},$$

the space of momenta away from the ‘colored diagonals’. One then sets $\partial \mathcal{A}[J, \bar{J}] / \partial \mathcal{B}$ to be the function that at $\mathbf{X} \in \mathcal{F}_{D,\ell}$ takes the value

$$\left. \frac{\delta^{2\ell} \mathcal{A}[J, \bar{J}]}{\delta \mathbf{j}(\mathcal{B})(\mathbf{X})} \right|_{J=\bar{J}=0}.$$

An important result is the independence of graphs, meaning that if \mathcal{C} is another graph $\partial \mathcal{C} / \partial \mathcal{B}$ is non-zero only if the graphs \mathcal{B} and \mathcal{C} are isomorphic. If that is the case, the derivative $\partial \mathcal{C} / \partial \mathcal{B}$ is found to be a group action by $\text{Aut}_c(\mathcal{B})$. Concretely, if $\mathbf{C} = (\mathbf{c}^1, \dots, \mathbf{c}^k) \in M_{D \times k}(\mathbb{Z})$ and $\mathbf{X} \in \mathcal{F}_{D,\ell}$,

$$\frac{\partial \mathcal{C}(\mathbf{C})}{\partial \mathcal{B}(\mathbf{X})} = \begin{cases} \delta_{\ell k} \cdot \left(\sum_{\hat{\sigma} \in \text{Aut}_c(\mathcal{B})} \delta_{\mathbf{x}^1, \mathbf{x}^2, \dots, \mathbf{x}^\ell}^{\mathbf{c}^{\sigma(1)}, \dots, \mathbf{c}^{\sigma(k)}} \right) & \text{if } \mathcal{B} \cong \mathcal{C}, \\ 0 & \text{if } \mathcal{B} \not\cong \mathcal{C}. \end{cases}$$

where⁹ $\hat{\sigma} \in \text{Aut}_c(\mathcal{B})$ denotes the lift of a permutation $\sigma \in \text{Sym}(\ell)$ to the¹⁰ corresponding element of the automorphism group $\text{Aut}_c(\mathcal{B})$. Now we are in position to define the correlation functions by

$$G_{\mathcal{B}}^{(2\ell)} \equiv \frac{\partial \log Z[J, \bar{J}; E]}{\partial \mathcal{B}}. \quad (3.7)$$

Example 3.1 (Meaning of $\partial / \partial \heartsuit$). If $\mathcal{A}[J, \bar{J}]$ is a functional and $\mathbf{X} = (\mathbf{x}^1, \mathbf{x}^2, \mathbf{x}^3) \in \mathcal{F}_{3,3}$, one has

$$\frac{\partial \mathcal{A}[J, \bar{J}]}{\partial \heartsuit}(\mathbf{X}) = \frac{\delta^6 \mathcal{A}[J, \bar{J}]}{\delta J_{\mathbf{x}^1} \delta J_{\mathbf{x}^2} \delta J_{\mathbf{x}^3} \delta \bar{J}_{\mathbf{y}^1} \delta \bar{J}_{\mathbf{y}^2} \delta \bar{J}_{\mathbf{y}^3}} \Big|_{J=0=\bar{J}} = \frac{\delta^6 \mathcal{A}[J, \bar{J}]}{\delta J_{\mathbf{x}^1} \delta J_{\mathbf{x}^2} \delta J_{\mathbf{x}^3} \delta \bar{J}_{x_1^1 x_2^2 x_3^3} \delta \bar{J}_{x_1^2 x_2^3 x_3^1} \delta \bar{J}_{x_1^3 x_2^1 x_3^2}} \Big|_{J=0=\bar{J}},$$

since

$$(\heartsuit)_* \{\mathbf{x}^1, \mathbf{x}^2, \mathbf{x}^3\} = \{\mathbf{y}^1, \mathbf{y}^2, \mathbf{y}^3\} = \left\{ \begin{bmatrix} x_1^1 \\ x_2^2 \\ x_3^3 \end{bmatrix}, \begin{bmatrix} x_1^2 \\ x_2^3 \\ x_3^1 \end{bmatrix}, \begin{bmatrix} x_1^3 \\ x_2^1 \\ x_3^2 \end{bmatrix} \right\}.$$

⁸Actually $D - 1$ of the pillows suffice, but that theory is ugly.

⁹Also, the delta $\delta_{\ell k}$ is somehow redundant (as it is a consequence of having isomorphic graphs).

¹⁰The colored automorphisms are rigid enough to be specified by only a permutation of the white (or black) vertices [16].

4. The tower of Schwinger-Dyson equations

With the Y -term known, it is clear that one can express it as a sum over graphs in the form $Y_{m_a}^{(a)}[J, \bar{J}] = \sum_{\mathcal{D}} f_{\mathcal{D}, m_a}^{(a)} \star \mathbf{j}(\mathcal{D})$. The graph calculus allows to compute these f -functions. In order to state the SDE tower, we only need a last graph operation, the swap ζ_a .

Let $\zeta_a(\mathcal{B}; v, w)$ swap of the a -colored edges at two black vertices v, w of a colored graph \mathcal{B} . Examples of this operation are¹²

$$\zeta_a(\text{graph}; v, w) = \text{graph} \quad \text{and} \quad \zeta_b(\text{graph}; \text{left, down}) = \text{graph} \sqcup \text{graph}.$$

In order to derive the SDE for $G_{\mathcal{B}}^{(2\ell)}$ one has to choose a black vertex of \mathcal{B} ; thus, in particular, if \mathcal{B} has no automorphisms (as e.g. F_c in rank 3) there are ℓ independent SDE for $G_{\mathcal{B}}^{(2\ell)}$. Derivatives with respect to the graphs $\zeta_a(\mathcal{B}; i, n) \equiv \zeta_a(\mathcal{B}; \mathbf{y}^i, \mathbf{y}^n)$, $n \neq i$, appear in the SDE.

Let \mathcal{B} be a connected boundary graph of the quartic rank- D model with pillow interactions, $S_{\text{int}} = \lambda \sum_{c=1}^D V_c$. Let \mathcal{B} have 2ℓ vertices. The (2ℓ) -point Schwinger-Dyson equations corresponding to \mathcal{B} are [18, Prop. 3.1]

$$\begin{aligned} & \left(1 + \frac{2\lambda}{E_s} \sum_{a=1}^D \sum_{\mathbf{q}\hat{a}} G_{\text{pillow}}^{(2)}(s_a, \mathbf{q}\hat{a}) \right) G_{\mathcal{B}}^{(2\ell)}(\mathbf{X}) \\ &= \frac{\delta_{\ell,1}}{E_s} + \frac{(-2\lambda)}{E_s} \sum_{a=1}^D \left\{ \sum_{\hat{\sigma} \in \text{Aut}_c(\mathcal{B})} \sigma^* f_{\mathcal{B}, s_a}^{(a)}(\mathbf{X}) + \sum_{n \neq i} \frac{1}{E_{y_a^n, s_a}} Z_0^{-1} \frac{\partial Z[J, \bar{J}]}{\partial \zeta_a(\mathcal{B}; i, n)(\mathbf{X})} \right. \\ & \quad \left. - \sum_{b_a} \frac{1}{E_{s_a b_a}} [G_{\mathcal{B}}^{(2\ell)}(\mathbf{X}) - G_{\mathcal{B}}^{(2\ell)}(\mathbf{X}|_{s_a \rightarrow b_a})] \right\} \quad (\mathbf{X} \in \mathcal{F}_{D, \ell}), \end{aligned} \quad (4.1)$$

with $\mathbf{s} = \mathbf{y}^i$ picked from $\mathcal{B}_*(\mathbf{X}) = (\mathbf{y}^1, \dots, \mathbf{y}^\ell)$, $1 \leq i \leq \ell$. Here $\sigma \in \text{Sym}(\ell)$ acts by permuting the arguments of $f_{\mathcal{B}, s_a}^{(a)}$. More explicit formulæ are given in [18] for ranks three, four and five.

5. Conclusions

The tools leading to the tower of SDE for arbitrary-rank TFTs with pillow interactions have been exposed. The kernel in the kinetic term should satisfy the mild condition \diamond . The scope of this method is boarder than only pillow interactions (e.g. for rank-3 TFTs, the list 2.1). The obtained equations are for (connected) correlation functions with connected boundary graph. The general result for arbitrary, disconnected graphs is work in progress, as is the extension of the present methods to fermionic fields and to $O(N)$ TFTs [17], aiming at SYK-like tensor models.

¹²Notice that in $\zeta_a(\text{graph}; v, w)$ there is no dependence on the choice of the vertices v and w , due to the symmetries of the graph graph .

References

- [1] H. Grosse and R. Wulkenhaar, *Self-Dual Noncommutative ϕ^4 -Theory in Four Dimensions is a Non-Perturbatively Solvable and Non-Trivial Quantum Field Theory*, *Commun. Math. Phys.* **329** (2014) 1069.
- [2] P. Di Francesco, P. H. Ginsparg and J. Zinn-Justin, *2-D Gravity and random matrices*, *Phys. Rept.* **254** (1995) 1.
- [3] J. Ambjørn, B. Durhuus and T. Jonsson, *Three-dimensional simplicial quantum gravity and generalized matrix models*, *Mod. Phys. Lett.* **A6** (1991) 1133.
- [4] R. Gurau and R. G. Gurau, *Random tensors*. Oxford University Press, 2016.
- [5] R. Gurău, *Colored Group Field Theory*, *Commun. Math. Phys.* **304** (2011) 69.
- [6] V. Bonzom, R. Gurău, A. Riello and V. Rivasseau, *Critical behavior of colored tensor models in the large N limit*, *Nucl. Phys.* **B853** (2011) 174.
- [7] R. Gurău and J. P. Ryan, *Colored Tensor Models - a review*, *SIGMA* **8** (2012) 020.
- [8] V. Rivasseau, *Random Tensors and Quantum Gravity*, *SIGMA* **12** (2016) 069.
- [9] A. Kitaev, A simple model of quantum holography (lecture), <http://online.kitp.ucsb.edu/online/entangled15/kitaev/>, 2015.
- [10] E. Witten, *An SYK-Like Model Without Disorder*, 1610.09758.
- [11] J. Maldacena and D. Stanford, *Remarks on the Sachdev-Ye-Kitaev model*, *Phys. Rev.* **D94** (2016) 106002 [1604.07818].
- [12] I. R. Klebanov and G. Tarnopolsky, *Uncolored random tensors, melon diagrams, and the Sachdev-Ye-Kitaev models*, *Phys. Rev.* **D95** (2017) 046004 [1611.08915].
- [13] V. Bonzom, L. Lionni and A. Tanasă, *Diagrammatics of a colored SYK model and of an SYK-like tensor model, leading and next-to-leading orders*, *J. Math. Phys.* **58** (2017) 052301 [1702.06944].
- [14] R. Alkofer, M. Q. Huber and K. Schwenzer, *Algorithmic derivation of Dyson-Schwinger Equations*, *Comput. Phys. Commun.* **180** (2009) 965 [0808.2939].
- [15] M. Q. Huber, Derivation of Dyson-Schwinger equations, physik.uni-graz.at/~mqh/notes/, 2017.
- [16] C. I. Pérez-Sánchez, *The Full Ward-Takahashi Identity for Colored Tensor Models*, *Commun. Math. Phys.* **358** (2018) 589.
- [17] S. Carrozza and A. Tanasă, *$O(N)$ Random Tensor Models*, *Lett. Math. Phys.* **106** (2016) 1531 [1512.06718].
- [18] C. I. Pérez-Sánchez and R. Wulkenhaar, *Correlation functions of coloured tensor models and their Schwinger-Dyson equations*, 1706.07358.
- [19] M. Disertori, R. Gurău, J. Magnen and V. Rivasseau, *Vanishing of Beta Function of Non Commutative Φ_4^4 Theory to all orders*, *Phys. Lett.* **B649** (2007) 95.
- [20] H. Grosse and R. Wulkenhaar, *Progress in solving a noncommutative quantum field theory in four dimensions*, 0909.1389.
- [21] P. Di Francesco, *Rectangular matrix models and combinatorics of colored graphs*, *Nucl. Phys.* **B648** (2003) 461.

- [22] D. O. Samary, *Closed equations of the two-point functions for tensorial group field theory*, *Class. Quant. Grav.* **31** (2014) 185005.
- [23] C. I. Pérez-Sánchez, *Surgery in colored tensor models*, *J. Geom. Phys.* **120** (2017) 262 [1608.00246].

POS(CORFU2017)147