

KLOE/KLOE-2 results and perspectives on dark force search

Elena Pérez del Río* for the KLOE-2 Collaboration

LNF-INFN Via E. Fermi 40, 00044 Frascati, Italy E-mail: eperez@lnf.infn.it

During the last years several Dark Sector Models have been proposed in order to address striking astrophysical observations which fail standard interpretations.

In the minimal case a new vector particle, the so called dark photon or U-boson, is introduced, with small coupling with Standard Model particles. Also, the existence of a dark Higgs boson h' is postulated, in analogy with the Standard Model, to give mass to the U-boson through the Spontaneous Symmetry Breaking mechanism.

The experiment KLOE, working on the DAΦNE $e+e-$ collider in Frascati, searched for the existence of the U-boson in a quite complete way, investigating several different processes and final states. Tight limits on the model parameters have been set at 90%CL. Further improvements are expected in terms of sensitivity and discovery potential with the new KLOE-2 detector working on the improved DAFNE $e+e-$ collider.

*The 39th International Conference on High Energy Physics (ICHEP2018)
4-11 July, 2018
Seoul, Korea*

*Speaker.

1. Introduction

The Standard Model (SM) does not provide a definitive model of all elementary particles. In particular, some astrophysical observations [1, 2, 3, 4, 5, 6, 7, 8] and the muon magnetic discrepancy a_μ are examples of possible physics beyond the SM [9, 10, 11, 12, 13] claim to explain the afore-mentioned anomalies by means of dark matter models, with a Weakly Interacting Massive Particle (WIMP) belonging to a secluded gauge sector. The new gauge interaction would be mediated by a new vector gauge boson, the U boson or dark photon, which could interact with the photon via a kinetic-mixing term ε^2 . In the following, some of the U boson searches, carried out with the KLOE detector, are described.

2. U -boson search in $e^+e^- \rightarrow U\gamma$ with $U \rightarrow e^+e^-$




Figure 1: Exclusion limits on the kinetic-mixing parameter, ε^2 , from KLOE: KLOE₁, KLOE₂ and KLOE₃ (in red) correspond to the combined limits from the analysis of $\phi \rightarrow \eta e^+e^-$, $e^+e^- \rightarrow \mu^+\mu^-\gamma$ and $e^+e^- \rightarrow e^+e^-\gamma$, respectively. The results are compared with the limits from E141, E774 [21], MAMI/A1 [22], APEX [23], WASA [24], HADES [25], NA48/2 [26] and BaBar [27]. The grey band indicates the parameter space favored by the $(g_\mu - 2)$ discrepancy.

The study of the reaction $e^+e^- \rightarrow U\gamma$, $U \rightarrow e^+e^-$ has the characteristic that allows to investigate the low mass region close to the di-electron mass threshold [18]. The signal of the U boson would be then expected as a resonant peak in the di-electron invariant mass. Since no signal was observed, the upper limit of the kinetic-mixing parameter as a function of m_U was evaluated with the CLs technique setting a limit on the U -boson signal at 90% confidence level Fig. 1. The integrated luminosity corresponds to $L_{integrated} = 1.54 \text{ fb}^{-1}$ from the 2004-2005 data campaign.

3. U -boson search in $e^+e^- \rightarrow U\gamma$ with $U \rightarrow \pi^+\pi^-$

The leptonic channels investigated by KLOE loose sensitivity in the $\rho - \omega$ region due to the dominant branching fraction into hadrons. The effective coupling of the U boson is predicted to be given by the product of the virtual-photon coupling and the kinetic-mixing parameter $\varepsilon^2 e F_\pi(q^2)$ [17]. For this search, a total integrated luminosity of 1.93 fb^{-1} was analyzed [28]. No signal was observed and a limit at the 90% CL was set on the coupling factor ε^2 in the energy range between 527 and 987 MeV with a larger sensitivity than previous limits in the $\rho - \omega$ region and above, see Fig. 2.




Figure 2: Left: 90% CL upper limit exclusion plot for ε^2 as a function of the U -boson mass ($KLOE_{(4)}$). **Right:** 90% CL exclusion plot for ε^2 as a function of the U -boson mass for the $e^+e^- \rightarrow U\gamma$ process. The $U \rightarrow \mu^+\mu^-$ limit (dashed line), the $U \rightarrow \pi^+\pi^-$ [28] constraint (solid line), and the $U \rightarrow \mu^+\mu^-, \pi^+\pi^-$ combination (blue area) at full KLOE statistics are presented in comparison with the competitive limits by BaBar [20], NA48/2 [26] and LHCb experiments [30]. The limits are shown together with previous KLOE results as well as other experiments at the moment of publication.

4. Combined limit in the production of U decaying into $\mu^+\mu^-$ and $\pi^+\pi^-$

A previous search for the U boson in the decay into $\mu^+\mu^-$ [31] has been extended by using the full KLOE statistics at $L_{int} = 1.93\text{fb}^{-1}$, updating the analysis with a new estimate of the background, analogous to the one used for the $U \rightarrow \pi^+\pi^-$ search. This new search confirms the non existence of U -boson signal in the di-muon invariant mass spectrum. To increase the sensitivity in the region of the $\rho - \omega$ interference, both results on the 90% upper limit for $\mu\mu$ and $\pi\pi$ have been combined, giving the up-to-date most stringent upper limit for the mixing parameter ε^2 in the U -boson mass region 519-987 MeV. The limit is shown in Fig. 2, together with the other most competitive limits in the region.

5. Conclusions

The KLOE collaboration has extensively contributed to the U -boson searches. Up to now, no evidence for a U boson was found and limits at the 90% confidence level were set on the kinetic-mixing parameter ε^2 in the mass range $5\text{MeV} < m_U < 987\text{MeV}$. In the meantime, a new data campaign has been finalized with the KLOE-2 setup, which has collected more than 5fb^{-1} in the past three years. The new setup and the enlarged statistics could further improve the current limits on the dark coupling constant by at least a factor of two.

6. Acknowledgements

We warmly thank our former KLOE colleagues for the access to the data collected during the KLOE data-taking campaign. We thank the DAΦNE team for their efforts in maintaining low background running conditions and their collaboration during all data taking. We want to thank our technical staff: G.F. Fortugno and F. Sborzacchi for their dedication in ensuring efficient operation of the KLOE computing facilities; M. Anelli for his continuous attention to the gas

system and detector safety; A. Balla, M. Gatta, G. Corradi and G. Papalino for electronics maintenance; C. Piscitelli for his help during major maintenance periods. This work was supported in part by the Polish National Science Centre through the Grants No. 2013/11/B/ST2/04245, 2014/14/E/ST2/00262, 2014/12/S/ST2/00459, 2016/21/N/ST2/01727, 2016/23/N/ST2/01293, 2017/26/M/ST2/00697.

References

- [1] P. Jean et al., *Astrophys.* **407**, L55 (2003).
- [2] C. E. Aalseth, et al., *Phys. Rev. Lett.* **107**, 141301 (2011).
- [3] R. Bernabei, et al., *Int. J. Mod. Phys.* **D13**, 2127 (2004).
- [4] R. Bernabei, et al., *Eur. Phys. J.* **C56**, 333 (2008).
- [5] J. Chang, et al. *Nature* 456, 362 (2008).
- [6] F. Aharonian, et al., *Phys. Rev. Lett.* **101**, 261104 (2008).
- [7] F. Aharonian, et al., *Astron. Astrophys.* **508**, 561 (2009).
- [8] A. A. Abdo, et al., *Phys. Rev. Lett.* **102**, 181101 (2009).
- [9] B. Holdom, *Phys. Lett.* **B166**, 196 (1985).
- [10] C. Boehm, P. Fayet, *Nucl. Phys.* **B683**, 219 (2004).
- [11] P. Fayet, *Phys. Rev.* **D75**, 115017 (2007).
- [12] M. Pospelov, A. Ritz, M.B. Voloshin, *Phys. Lett.* **B662**, 53 (2008).
- [13] Y. Mambrini, *J. Cosmol. Astropart. Phys.* **1009**, 022 (2010).
- [14] A. Gallo, et al., *DAFNE status report, Conf. Proc.* **C060626** (2006) 604-606.
- [15] M. Adinolfi et al., *Nucl. Instrum. Methods* **A488**, 51 (2002)
- [16] M. Adinolfi et al., *Nucl. Instrum. Methods* **A482**, 364 (2002)
- [17] M. Reece, L.T. Wang, *JHEP* **07**, 51 (2009).
- [18] A. Anastasi, et al., *Physics Letters* **B750**, 633-637 (2015).
- [19] L. Barzé et al., *Eur. Phys. J.* **C71**, 1680 (2011).
- [20] J.P. Lees et al., *Phys. Rev. Lett.* **108**, 211801 (2012).
- [21] J. D. Bjorken, et al., *Phys. Rev.* **D 80**, 075018 (2009).
- [22] H. Merkel et al., *Phys. Rev. Lett.* **[112]**, 221802 (2014).
- [23] S. Abrahamyan et al., *Phys. Rev. Lett.* **107**, 191804 (2011).
- [24] P. Adlarson et al., *Phys. Lett.* **B726**, 187 (2013).
- [25] G. Agakishiev et al., *Phys. Lett.* **B731**, 265-271 (2014).
- [26] J.R. Batley et al., *Phys. Lett.* **B746**, 178-185 (2015).
- [27] J.P. Lees et al., *Phys. Rev. Lett.* **113**, 201801 (2014).
- [28] A. Anastasi et al., *Phys. Lett.* **B757** (2016) 356-361
- [29] G. Gounaris, J.J. Sakurai, *Phys. Rev. Lett.* **21** (1968) 244.
- [30] S. Abrahamyan et al., *Phys. Rev. Lett.* **107** (2011) 191804.
- [31] D. Babusci et al., *Phys. Lett.* **B736**, 459-464 (2014).