

## Interferometric Monitoring of Gamma-ray Bright AGNs: J1159+2914

---

Sincheol Kang<sup>\*,ab</sup>, Sang-Sung Lee,<sup>ab</sup> Do-Young Byun,<sup>ab</sup> Jeffrey Hodgson<sup>a</sup> and Jee Won Lee<sup>a</sup>

<sup>a</sup> Korean Astronomy and Space Science Institute

<sup>b</sup> Korea, University of Science and Technology

E-mail: kang87@kasi.re.kr, sslee@kasi.re.kr, bdy@kasi.re.kr, jhodgson@kasi.re.kr, jwlee78@kasi.re.kr

We present the results of multi-epoch monitoring of a blazar J1159+2914, one of the targets of a Very Long Baseline Interferometry (VLBI) monitoring program : Interferometric Monitoring of Gamma-ray Bright AGNs (iMOGABA), as a Korean VLBI Network (KVN) Key Science Program (KSP). The observations were conducted simultaneously at 22, 43, 86, and 129 GHz, during 4 years from December 2012 to December 2016. Obtained total fluxes ranged from 0.26 to 2.88 Jy at all frequencies with a mean rms noise of 26 mJy. We also used the 15 and 230 GHz data observed by Owens Valley Radio Observatory (OVRO) and Sub-Millimeter Array (SMA), respectively. In order to analyze the characteristics of variabilities, we estimated variability time scale from 15 GHz data, using three different functions, such as structure function, Gaussian function, and exponential function. Also, in order to study the multi-frequency correlations, we compared the light curve of 15 GHz with that of 22, 43, and 86 GHz, using cross-correlation analysis. Moreover, we estimated B-field strength using core sizes from VLBA 43 GHz data, turnover frequencies and maximum total fluxes from KVN data, and variability time scales from OVRO data, in order to study the variability of B-field strength nearby the radio emission region.

*14th European VLBI Network Symposium & Users Meeting (EVN 2018)  
8-11 October 2018  
Granada, Spain*

---

\*Speaker.

## 1. Introduction

Blazars are one of the sub-classes of AGNs with one of two jets being launched nearly toward us. Their emissions are highly Doppler boosted and they show dynamic variations. The origins of these drastic variations are usually presumed as source intrinsic or source extrinsic variations.

A blazar J1159+2914 (also known as 1156+295, 4C +29.45, and TON 599) has shown the wide time scale range of variations from hours to years. Many previous studies attempted to prove the mechanism of these variations (Liu et al. 2003, Wang et al. 2014)


The source is one of the targets of the monitoring program of the Korean Very Long Baseline Interferometry (VLBI) Network (KVN), named as Interferometric MONitoring of GAMMA-ray Bright AGNs (iMOGABA). Since December 2012, the iMOGABA observations have been conducted, nearly monthly, for 24 hours at each epoch, targeting on over 30 sources at 22, 43, 86, and 129 GHz, simulataneously (Lee et al. 2016). The purposes of the iMOGABA program as a KVN Key Science Program (KSP) are to study the correlation of behavior between radio and gamma-ray wavelengths in bright AGNs at gamma-ray wavelength and to constrain the characteristics of the gamma-ray flare from the AGNs.

## 2. Observations and Data Reduction

The multi-wavelength VLBI observations had been conducted from December 2012 to December 2016. A blazar J1159+2914 was observed 33, 30, 31, and 22 of 33 epochs at 22, 43, 86, and 129 GHz, respectively. Correlated data were reduced via the Astronomical Image Processing System (AIPS) with the KVN pipeline (Hodgson et al. 2016) We imaged the reduced data from AIPS using DIFMAP program. Before imaging, we averaged the data every 30, 30, 20, and 10 seconds at 22, 43, 86, and 129 GHz, respectively. We assumed that the source is core dominated within KVN VLBI beam size for all epochs, so we CLEANed the images with single CLEAN window at image center, window sizes of 1.5 X 1.5, 0.76 X 0.76, 0.4 X 0.4, and 0.26 X 0.26 mas at 22, 43, 86, and 129 GHz, respectively. After the CLEANing, we obtained beam sizes, total CLEANed flux densities, peak flux densities, rms noises, dynamic ranges, and image qualities. And we fitted the souece with circular Gaussian models. After the model fitting, we obtained modeled total flux densities, peak flux densities, sizes and brightness temperatures of core for each epoch at each frequencies.

## 3. Multi-Wavelength Data

To confirm the correlations of flux variability in wide range of wavelength, we compared our data with the both 15 GHz data obtained from Owens Valley Radio Observatory (OVRO) and 230 GHz data obtained from Sub-Millimeter Array (SMA). Figure 2 shows the multi-wavelength light curves. And to estimate the core sizes at 43 GHz, we re-modeled the published data from Boston University (BU) blazar group observed by Very Long Baseline Array (VLBA). Because, we presumed actual core sizes of J1159+2914 are smaller than the beam size of KVN.



**Figure 1:** Light curves of J1159+2914.

Then we estimated turnover frequencies and maximum flux densities for 17 epochs when the source was detected at all four KVN frequencies with following equation.

$$S = c_1 \left( \frac{\nu}{\nu_r} \right)^{\alpha + \ln(\nu/\nu_r)} \quad (3.1)$$

Here, we assumed flat spectrum so  $\alpha$  is 0. And we estimated  $c_1$ ,  $c_2$ , and  $\nu_r$  by curve fitting. Obtained turnover frequencies ranged from 19 to 37 GHz and maximum flux densities ranged from 0.73 to 2.7 Jy. To estimate B-field strength at turnover frequencies, we re-estimated core sizes at turnover frequencies from core sizes at 43 GHz of BU data ( $R \propto \nu^{-\epsilon}$ , Algaba et al. 2017). And the obtained turnover frequencies were lower than 43 GHz, we assumed that data at 43, 86, and 129 GHz were originated from optically thin region, so we estimated spectral indices of optically thin region with 43, 86, and 129 GHz data. Obtained spectral indices of optically thin region ranged from -0.72 to -0.20.

#### 4. Variability time scale

In order to study the characteristics of time-dependent variations, we fitted the OVRO 15 GHz light curve using the following exponential function (Prince et al. 2017).

$$F(t) = 2F_0 \left[ \exp\left(\frac{t_0 - t}{\tau_r}\right) + \exp\left(\frac{t - t_0}{n\tau_r}\right) \right]^{-1} \quad (4.1)$$

Here,  $F_0$  is local maximum flux density,  $t_0$  is time at a local maximum,  $\tau_r$  is rising variability time scale and  $n$  is ratio between rising and decaying variability time scales.

In this fitting procedure, we combined one decaying (half) flare and six rising and decaying flares with zero base level. Base level of light curve was estimated after summing all flares. Fitting parameters are listed in Figure 3. We defined flares by visual inspection. We note that the rising time scales were longer than decaying time scales in this fitting, unlike previous several studies (Valtaoja & Lainela 1999).

## 5. Variability Brightness Temperature, Doppler Factor, and B-Field Strength


We estimated variability brightness temperatures and Doppler factors using the following equations with above estimated variability time scales.

$$T_b^{var} = 3.47 \times 10^5 \Delta S \left\{ \frac{\lambda D_L}{\tau_{var}(1+z)^2} \right\}^2 \quad \text{and} \quad \delta_{var} = (1+z) \left( \frac{T_b^{var}}{10^{12}} \right)^{1/(3+\alpha)} \quad (5.1)$$

But, we excluded the first flare because it showed only decaying phase. Here,  $\Delta S$  is  $F_0$ ,  $\lambda$  is observing wavelength, and  $D_L$  is luminosity distance. We used  $\lambda$  and  $D_L$  as 2 cm and 4477.8 Mpc, respectively. And  $\alpha$  is the spectral index and we assumed flat spectrum. Applying these  $\delta_{var}$  to estimated core sizes, turnover frequencies, and maximum flux densities, we estimated B-field strengths using the following equation.

$$B = 10^{-5} b(\alpha) S_m^{-2} \theta^4 v_c \left( \frac{\delta}{1+z} \right)^{-1} \quad (5.2)$$

Here,  $b(\alpha)$  is a factor depends on optically thin spectral index. Values of  $b(\alpha)$  were referred from Marscher 1983. Obtained B-field strengths are plotted in Figure 5. And we assumed the Doppler



**Figure 2:** B-field strengths.

factor of 7.10 as the upper limit of other quiet phases. We found that B-field strengths between MJD 56600 and 56900 were significantly higher than other epochs. These high values are might affected by core sizes. Still, we have lots of debates for this result.

## 6. Summary and Future Works

We observed a blazar J1159+2914, one of targets of iMOGABA, from December 2012 to December 2016. We compared our light curves with that of OVRO and SMA data, in order to study the multi-wavelength correlations. To estimate variability time scales, we fitted the light curve of OVRO 15 GHz data, using exponential function. And then, we estimated variability brightness temperatures and Doppler factors, in order to estimate B-field strengths. With estimated B-field strengths, we will discuss the conditions of relativistic jet. Also, we can compare the estimated B-field strengths with B-field strengths of equipartition condition.

## References

- [1] Antonucci, R. 1993, Unified Models for Active Galactic Nuclei and Quasars, *Annu. Rev. Astron. Astrophys.* 31:473-521
- [2] Favre, P., Courvoisier, T. J.-L., and Paltani, S., 2005, AGN variability time scales and the discrete-event model, *A&A* 443, 451-463
- [3] Gurwell, M. A., Peck, A. B., Hostler, S. R., Darrah, M. R., & Katz, C. A., 2007, in *ASP Conf. Ser.* 375, *From Z-Machines to ALMA: (Sub)millimeter Spectroscopy of Galaxies*, ed. A. J. Baker et al. (San Francisco, CA: ASP), 234
- [4] Hodgson, J., Lee, S.-S., Zhao, G.-Y., et al. 2016, The Automatic Calibration of Korean VLBI Network Data, *JKAS*, 49, 137
- [5] Jorstad, S., & Marscher, A., 2016, *The VLBA-BU-Blazar Multi-wavelength Monitoring Program*, *Galaxies*, vol. 4, issue 4, p. 47.
- [6] Lee, S.-S., Lobanov A. P., Krichbaum T. P., et al. 2008, A Global 86 GHz VLBI Survey of Compact Radio Sources. *AJ*, 136, 159
- [7] Lee, S.-S., Wajima, K., Algaba, J.-C., et al. 2016, Interferometric Monitoring of Gamma-Ray Bright AGNs. I. The Results of Single-Epoch Multifrequency Observations, *ApJS*, 227, 8
- [8] Lee, J. W., Lee, S.-S., Hodgson, J., et al. 2017, Interferometric Monitoring of Gamma-Ray Bright AGNs: S5 0716+714. *ApJ*, 841:119
- [9] Liu, B.-R., Liu X., Marchili N., et al. 2013, Two-year monitoring of intra-day variability of quasar 1156+295 at 4.8 GHz, *A&A*, 555, A134
- [10] Marscher, A. P., Jorstad, S. G., D’Arcangelo, F. D., et al. 2008, The inner jet of an active galactic nucleus as revealed by a radio-to- $\gamma$ -ray outburst, *Natur*, 452, 966
- [11] Prince, R., Majumdar, P., & Gupta, N., 2017, Long-term Study of the Light Curve of PKS 1510-089 in GeV Energies, *ApJ*, 844(1), 62.
- [12] Richards, J. L., Max-Moerbeck, W., Pavlidou, V., et al. 2011, Blazars in the Fermi Era: The OVRO 40 m Telescope Monitoring Program, *ApJS*, 194, 29
- [13] Valtaoja, E., and Lainela, M., 1999, Total Flux Density Variations in Extragalactic Radio Sources. I. Decomposition of Variations into Exponential Flares, *ApJS*, 120:95-99
- [14] Wang J.-Y., An T., Baan W. A., et al. 2014, Periodic Radio Variabilities of the Blazar 1156+295: Harmonic Oscillations, *MNRAS*, 443, 58-66
- [15] Zhao W., Hong X.-Y., An T., et al. 2011, Radio Structure of the Blazar 1156+295 with sub-pc Resolution, *A&A*, 529, A113