

Explaining the Flavor Anomalies with a Vector Leptoquark (Moriond 2019 update)

Andreas Crivellin*

Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland

Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

E-mail: andreas.crivellin@cern.ch

Francesco Saturnino^{†‡}

Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics,

University of Bern, CH-3012 Bern, Switzerland

E-mail: saturnino@itp.unibe.ch

Several experiments revealed intriguing hints for lepton flavor universality (LFU) violating new physics (NP) in semi-leptonic B meson decays, mainly in $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell^+\ell^-$ transitions at the $3 - 5\sigma$ level. Leptoquarks (LQ) are prime candidates to address these anomalies as they contribute to semi-leptonic decays already at tree level while effects in other flavor observables, agreeing with the standard model (SM), are loop suppressed.

In these proceedings we review the vector leptoquark $SU(2)_L$ singlet, contained in the famous Pati-Salam model, which is able to address both $b \rightarrow c\tau\nu$ and $b \rightarrow s\mu^+\mu^-$ data simultaneously. Due to the large couplings to tau leptons needed to account for the $b \rightarrow c\tau\nu$ data, sizable loop effects arise which we include in our phenomenological analysis. Updating our result of Ref. [1] with the recent measurements of LHCb [2] and BELLE [3, 4] we find an even better fit to data than before.

*XXVII International Workshop on Deep-Inelastic Scattering and Related Subjects - DIS2019
8-12 April, 2019, Torino, Italy*

*The work of A.C. is supported by a Professorship Grant (PP00P2_176884) of the Swiss National Science Foundation.

[†]Speaker.

[‡]I thank the organizers of the DIS2019 in Turin for giving me the opportunity to present my work, which is supported by the Swiss National Foundation under grant 200020_175449/1. We are very grateful to Joaquim Matias and Bernat Capdevilla for providing us with the fit necessary for the $b \rightarrow s\ell^+\ell^-$ region in Fig. 2. These proceedings match the ones of Andreas Crivellin for Moriond QCD & High Energy Interactions (23-30 March 2019, La Thuile, Italy) and are submitted as a common arxiv version.

1. Introduction

While so far the LHC has not detected any particles beyond the ones present in the Standard Model (SM), intriguing hints for LFU violation in semi-leptonic B -meson decays were accumulated in several (classes of) observables:

$$b \rightarrow s\ell^+\ell^-$$

In these flavor changing neutral current transitions, measurements of the ratios

$$R(K^{(*)}) = \frac{\text{Br}[B \rightarrow K\mu^+\mu^-]}{\text{Br}[B \rightarrow Ke^+e^-]}$$

show sizable deviations from their respective SM prediction. While the newest measurement of $R(K)$ by the LHCb collaboration [2] shows a deviation of 2.5σ from the SM, the Belle result for $R(K^{(*)})$ is consistent with the SM [3]. However, due to the larger errors, this result also agrees with previous LHCb measurements of $R(K^{(*)})$ which deviate from the SM [5] in the same direction as $R(K)$. Taking into account all other $b \rightarrow s\mu^+\mu^-$ observables (like the lepton flavor universal observable P'_5 [6]), the global fit prefers various NP scenarios above the 5σ level [7] compared to the SM, also when the newest measurements are taken into account [8–11].

In order to resolve the discrepancy in the neutral current transitions, an effect of $\mathcal{O}(10\%)$ is required at the amplitude level. Since this flavor changing neutral current (FCNC) is suppressed in the SM as it is only induced at one loop level, a small NP contribution is already sufficient. In a global fit one finds a preference for scenarios like $C_9^{\mu\mu} = -C_{10}^{\mu\mu}$ (i.e. a left-handed current coupling to muons only) [8]. Such an effect is naturally obtained at tree-level with the vector LQ $SU(2)$ singlet [1, 12–32]. However, a $C_9^{\mu\mu} = -C_{10}^{\mu\mu}$ effect complemented by a flavor universal effect in C_9 gives an even better fit to data [8, 33]. As we will see, this is exactly the pattern that arises in our model.

$$b \rightarrow c\tau\nu$$

There are also indications for LFU violation in charged current transitions, namely in the ratios

$$R(D^{(*)}) = \frac{\text{Br}[B \rightarrow D^{(*)}\tau\nu]}{\text{Br}[B \rightarrow D^{(*)}\ell\nu]}$$

where $\ell = \{e, \mu\}$. While the newest measurements from Belle [4] agree with the SM prediction, including previous measurements by BaBar, Belle and LHCb still yield a deviation of 3.1σ [34] from the SM prediction. Furthermore there is also a measurement of the ratio $R(J/\Psi) = \frac{\text{Br}[B_c \rightarrow J/\Psi\tau\nu]}{\text{Br}[B_c \rightarrow J/\Psi\mu\nu]}$ exceeding its SM prediction [35].

Also here a NP effect of $\mathcal{O}(10\%)$ is needed at the amplitude level. However, since $b \rightarrow c\tau\nu$ transitions are mediated at tree level by the exchange of a W boson in the SM, the NP effect needs to be large. This means that NP should contribute at tree level with sizable couplings and at a not too high NP scale. Here, the best single particle solution is the vector LQ $SU(2)$ singlet [1, 12–32] since it does not give a tree-level effect in $b \rightarrow s\nu\nu$ processes and provides a common rescaling of $R(D)$ and $R(D^*)$ with respect to the SM prediction.




Figure 1: Left: Feynman diagram depicting the loop effects induced by the $bc\tau\nu$ operator from $SU(2)$ invariance. Right: $C_{9, sb}^{\ell\ell}$ and $C_7^{sb}(\mu_b)$, generated by these loop effects, as functions of $R(D^{(*)})/R(D^{(*)})_{\text{SM}}$. The solid (dashed) lines correspond to $M = 1$ TeV (5 TeV) while the (dark) blue region is preferred by $b \rightarrow c\tau\nu$ data at the 1σ (2σ) level, taking into account the most recent measurements. From the global fit, taking into account only lepton flavor conserving observables, we have $-1.29 < C_{9, sb}^{\ell\ell} < -0.87$ [38] and $-0.01 < C_7^{sb}(\mu_b) < 0.05$ [7] at the 1σ level. Assuming an explanation of $b \rightarrow c\tau\nu$, our model predicts the right size and sign of the effect in $C_{9, sb}^{\ell\ell}$ and $C_7^{sb}(\mu_b)$ needed to explain $b \rightarrow s\ell^+\ell^-$ data.

2. The Pati Salam vector leptoquark as combined solution to the anomalies

The vector Leptoquark $SU(2)_L$ singlet with hypercharge $-4/3$, arising in the famous Pati-Salam model [36], is a prime candidate to explain both the anomalies in charged current and neutral current B decays simultaneously [12–14, 17–20]. It gives a $C_9 = -C_{10}$ effect in $b \rightarrow s\ell^+\ell^-$ at tree level and at the same time a sizable effect in $b \rightarrow c\tau\nu$ without violating bounds from $b \rightarrow s\nu\nu$ and/or direct searches and does not lead to proton decay. Note that this LQ by itself is not UV complete, however several UV complete models for this LQ have been proposed [15, 16, 21–29, 37].

For the purpose of our phenomenological analysis, let us consider a model where we simply extend the SM by this LQ. Its interaction with the SM particles is given by the Lagrangian

$$\mathcal{L}_{V_1} = \kappa_{fi}^L \bar{Q}_f \gamma_\mu L_i V_\mu^{1\dagger} + h.c. ,$$

where $Q(L)$ is the quark (lepton) $SU(2)_L$ doublet, κ_{fi}^L represents the couplings of the LQ to the left handed quarks (leptons) and f and i are flavor indices. Note that in principle couplings to right-handed SM particles are also allowed, they are however not relevant for this discussion. After electro-weak symmetry breaking, we work in the down basis, meaning that no CKM matrix elements appear in FCNC processes.

We start by taking κ_{23}^L and κ_{33}^L as the only non-zero couplings, as they are necessary to explain $b \rightarrow c\tau\nu$ data. Here, strong effects in $b \rightarrow s\tau^+\tau^-$ transitions [39] are generated which at the 1-loop level affect $b \rightarrow s\ell^+\ell^-$ via the Wilson coefficients $C_{9, sb}^{\ell\ell}$ and C_7^{sb} , as is depicted to the left in Fig. 1. Due to the correlation with $b \rightarrow c\tau\nu$, these Wilson coefficients can be expressed as functions of $R(D^{(*)})/R(D^{(*)})_{\text{SM}}$. The Wilson coefficients' dependency on these ratios is shown in the right plot of Fig. 1, where the RGE evolution of C_7^{sb} from the NP scale down to the b quark scale is also taken into account (see Ref. [40]). Interestingly, assuming an explanation of $b \rightarrow c\tau\nu$ data, the effects




Figure 2: Allowed (colored) regions in the $C_{9, sb}^{\mu\mu} = -C_{10, sb}^{\mu\mu}$ ($\hat{=} 640\kappa_{22}^L \kappa_{32}^{L*}$) $- R(X)/R(X)_{SM}$ plane for $M = 1$ TeV and $X = \{D, D^*\}$ at the 1σ and 2σ level. The region above the black dashed (solid) line is excluded by $\tau \rightarrow \phi\mu$ ($B \rightarrow K\tau\mu$) for $\kappa_{33}^L = 0.5 = 25\kappa_{32}^L$ ($\kappa_{33}^L = 0.5 = 2.5\kappa_{32}^L$). The bound from $\tau \rightarrow \phi\mu$ ($B \rightarrow K\tau\mu$) depends on κ_{33}^L and κ_{32}^L and gets stronger if κ_{32}^L gets smaller (larger). That is, for $\kappa_{33}^L = 0.5$ and $2.7 \lesssim \kappa_{33}^L/\kappa_{32}^L \lesssim 27$, the whole 2σ region preferred by $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell^+\ell^-$ data is consistent with these bounds. Note that we used the most recent experimental results for both the $b \rightarrow c\tau\nu$ and $b \rightarrow s\ell^+\ell^-$ transitions, therefore updating our analysis in Ref. [40].

generated in $C_{9, sb}^{\ell\ell}$ and C_7^{sb} agree with the 1σ ranges of the model independent fit to $b \rightarrow s\mu^+\mu^-$ data excluding LFU violating observables [38, 41].

Now we also allow κ_{32}^L and κ_{22}^L to be non-zero, generating a tree level effect in $b \rightarrow s\mu^+\mu^-$ which is necessary to account for the LFU violating observables as well. In Fig. 2 we show the allowed (colored) regions from $b \rightarrow s\mu^+\mu^-$ and $b \rightarrow c\tau\nu$ as well as the exclusions from $b \rightarrow s\tau\mu$ and $\tau \rightarrow \phi\mu$. A simultaneous explanation of the anomalies is perfectly possible since the colored regions overlap and do not extend to the parameter space excluded by $b \rightarrow s\tau\mu$ and $\tau \rightarrow \phi\mu$. Interestingly, we predict a lepton flavor universal effect in $C_{9, sb}^{\ell\ell}$ and C_7^{sb} in addition to a LFU violating tree-level effect of the form $C_{9, sb}^{\mu\mu} = -C_{10, sb}^{\mu\mu}$ in muonic channels only. This means that the effect of NP compared to the SM is expected to be larger in lepton flavor universal observables like $P5'$ relative to LFU violation observables as $R(K^{(*)})$, which is in perfect agreement with global fit scenarios [8]. In fact, the agreement is even better after the inclusion of the new measurements of BELLE and LHCb.

References

- [1] A. Crivellin, C. Greub, D. Müller and F. Saturnino, Phys. Rev. Lett. **122**, no. 1, 011805 (2019) doi:10.1103/PhysRevLett.122.011805 [arXiv:1807.02068 [hep-ph]].
- [2] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **122**, no. 19, 191801 (2019) doi:10.1103/PhysRevLett.122.191801 [arXiv:1903.09252 [hep-ex]].
- [3] A. Abdesselam *et al.* [Belle Collaboration], arXiv:1904.02440 [hep-ex].

- [4] A. Abdesselam *et al.* [Belle Collaboration], arXiv:1904.08794 [hep-ex].
- [5] R. Aaij *et al.* [LHCb Collaboration], JHEP **1708**, 055 (2017) doi:10.1007/JHEP08(2017)055 [arXiv:1705.05802 [hep-ex]].
- [6] R. Aaij *et al.* [LHCb Collaboration], JHEP **1602**, 104 (2016) doi:10.1007/JHEP02(2016)104 [arXiv:1512.04442 [hep-ex]].
- [7] B. Capdevila, A. Crivellin, S. Descotes-Genon, J. Matias and J. Virto, JHEP **1801**, 093 (2018) doi:10.1007/JHEP01(2018)093 [arXiv:1704.05340 [hep-ph]].
- [8] M. Algueró, B. Capdevila, A. Crivellin, S. Descotes-Genon, P. Masjuan, J. Matias and J. Virto, arXiv:1903.09578 [hep-ph].
- [9] J. Aebischer, W. Altmannshofer, D. Guadagnoli, M. Reboud, P. Stangl and D. M. Straub, arXiv:1903.10434 [hep-ph].
- [10] M. Ciuchini, A. M. Coutinho, M. Fedele, E. Franco, A. Paul, L. Silvestrini and M. Valli, arXiv:1903.09632 [hep-ph].
- [11] A. Arbey, T. Hurth, F. Mahmoudi, D. M. Santos and S. Neshatpour, arXiv:1904.08399 [hep-ph].
- [12] R. Alonso, B. Grinstein and J. Martin Camalich, JHEP **1510**, 184 (2015) doi:10.1007/JHEP10(2015)184 [arXiv:1505.05164 [hep-ph]].
- [13] L. Calibbi, A. Crivellin and T. Ota, Phys. Rev. Lett. **115**, 181801 (2015) doi:10.1103/PhysRevLett.115.181801 [arXiv:1506.02661 [hep-ph]].
- [14] S. Fajfer and N. Konik, Phys. Lett. B **755**, 270 (2016) doi:10.1016/j.physletb.2016.02.018 [arXiv:1511.06024 [hep-ph]].
- [15] R. Barbieri, G. Isidori, A. Pattori and F. Senia, Eur. Phys. J. C **76**, no. 2, 67 (2016) doi:10.1140/epjc/s10052-016-3905-3 [arXiv:1512.01560 [hep-ph]].
- [16] R. Barbieri, C. W. Murphy and F. Senia, Eur. Phys. J. C **77**, no. 1, 8 (2017) doi:10.1140/epjc/s10052-016-4578-7 [arXiv:1611.04930 [hep-ph]].
- [17] G. Hiller, D. Loose and K. Schönwald, JHEP **1612**, 027 (2016) doi:10.1007/JHEP12(2016)027 [arXiv:1609.08895 [hep-ph]].
- [18] B. Bhattacharya, A. Datta, J. P. Guévin, D. London and R. Watanabe, JHEP **1701**, 015 (2017) doi:10.1007/JHEP01(2017)015 [arXiv:1609.09078 [hep-ph]].
- [19] D. Buttazzo, A. Greljo, G. Isidori and D. Marzocca, JHEP **1711**, 044 (2017) doi:10.1007/JHEP11(2017)044 [arXiv:1706.07808 [hep-ph]].
- [20] J. Kumar, D. London and R. Watanabe, Phys. Rev. D **99**, no. 1, 015007 (2019) doi:10.1103/PhysRevD.99.015007 [arXiv:1806.07403 [hep-ph]].
- [21] N. Assad, B. Fornal and B. Grinstein, Phys. Lett. B **777**, 324 (2018) doi:10.1016/j.physletb.2017.12.042 [arXiv:1708.06350 [hep-ph]].
- [22] L. Di Luzio, A. Greljo and M. Nardecchia, Phys. Rev. D **96**, no. 11, 115011 (2017) doi:10.1103/PhysRevD.96.115011 [arXiv:1708.08450 [hep-ph]].
- [23] L. Calibbi, A. Crivellin and T. Li, Phys. Rev. D **98**, no. 11, 115002 (2018) doi:10.1103/PhysRevD.98.115002 [arXiv:1709.00692 [hep-ph]].

- [24] M. Bordone, C. Cornella, J. Fuentes-Martin and G. Isidori, *Phys. Lett. B* **779**, 317 (2018) doi:10.1016/j.physletb.2018.02.011 [arXiv:1712.01368 [hep-ph]].
- [25] R. Barbieri and A. Tesi, *Eur. Phys. J. C* **78**, no. 3, 193 (2018) doi:10.1140/epjc/s10052-018-5680-9 [arXiv:1712.06844 [hep-ph]].
- [26] M. Blanke and A. Crivellin, *Phys. Rev. Lett.* **121**, no. 1, 011801 (2018) doi:10.1103/PhysRevLett.121.011801 [arXiv:1801.07256 [hep-ph]].
- [27] A. Greljo and B. A. Stefanek, *Phys. Lett. B* **782**, 131 (2018) doi:10.1016/j.physletb.2018.05.033 [arXiv:1802.04274 [hep-ph]].
- [28] M. Bordone, C. Cornella, J. Fuentes-Martín and G. Isidori, *JHEP* **1810**, 148 (2018) doi:10.1007/JHEP10(2018)148 [arXiv:1805.09328 [hep-ph]].
- [29] S. Matsuzaki, K. Nishiwaki and K. Yamamoto, *JHEP* **1811**, 164 (2018) doi:10.1007/JHEP11(2018)164 [arXiv:1806.02312 [hep-ph]].
- [30] L. Di Luzio, J. Fuentes-Martin, A. Greljo, M. Nardecchia and S. Renner, *JHEP* **1811**, 081 (2018) doi:10.1007/JHEP11(2018)081 [arXiv:1808.00942 [hep-ph]].
- [31] A. Biswas, D. Kumar Ghosh, N. Ghosh, A. Shaw and A. K. Swain, arXiv:1808.04169 [hep-ph].
- [32] A. Angelescu, D. Beirevi, D. A. Faroughy and O. Sumensari, *JHEP* **1810**, 183 (2018) doi:10.1007/JHEP10(2018)183 [arXiv:1808.08179 [hep-ph]].
- [33] M. Algueró, B. Capdevila, S. Descotes-Genon, P. Masjuan and J. Matias, *Phys. Rev. D* **99**, no. 7, 075017 (2019) doi:10.1103/PhysRevD.99.075017 [arXiv:1809.08447 [hep-ph]].
- [34] Y. Amhis *et al.* [HFLAV Collaboration], *Eur. Phys. J. C* **77**, no. 12, 895 (2017) doi:10.1140/epjc/s10052-017-5058-4 [arXiv:1612.07233 [hep-ex]].
- [35] R. Aaij *et al.* [LHCb Collaboration], *Phys. Rev. Lett.* **120**, no. 12, 121801 (2018) doi:10.1103/PhysRevLett.120.121801 [arXiv:1711.05623 [hep-ex]].
- [36] J. C. Pati and A. Salam, *Phys. Rev. D* **10**, 275 (1974) Erratum: [*Phys. Rev. D* **11**, 703 (1975)]. doi:10.1103/PhysRevD.10.275, 10.1103/PhysRevD.11.703.2
- [37] C. Cornella, J. Fuentes-Martin and G. Isidori, arXiv:1903.11517 [hep-ph].
- [38] S. Descotes-Genon, L. Hofer, J. Matias and J. Virto, *JHEP* **1606**, 092 (2016) doi:10.1007/JHEP06(2016)092 [arXiv:1510.04239 [hep-ph]].
- [39] B. Capdevila, A. Crivellin, S. Descotes-Genon, L. Hofer and J. Matias, *Phys. Rev. Lett.* **120**, no. 18, 181802 (2018) doi:10.1103/PhysRevLett.120.181802 [arXiv:1712.01919 [hep-ph]].
- [40] A. Crivellin and F. Saturnino, arXiv:1905.08257 [hep-ph].
- [41] W. Altmannshofer and D. M. Straub, *Eur. Phys. J. C* **75**, no. 8, 382 (2015) doi:10.1140/epjc/s10052-015-3602-7 [arXiv:1411.3161 [hep-ph]].