

# High School Students' Muon Underground Shielding Experiment

---

**Nathan A. Unterman<sup>1</sup>**

*New Trier High School*

*7 North Happ Road, Northfield, IL 60093 USA*

*E-mail: nunterman@gmail.com*

**Mark Adams**

*University of Illinois at Chicago*

*Physics Department, 845 W. Taylor Street, Chicago, IL 60607, USA*

*E-mail: adams@uic.edu*

**Tom Blackmore, Emmanuelle Copeland, Marybeth Senser**

*Downers Grove South High School*

*Science Department, 1436 Norfolk Street, Downers Grove, IL 60516, USA*

*E-mail: msenser@csd099.org*

**Henry Seiden, Anthony Valsamis**

*Glenbrook North High School*

*Science Department, 2300 Shermer Road, Northbrook, IL 60062, USA*

*E-mail: avalsamis@glenbrook225.org*

**Benjamin Grey, Jacob Miller, Max Miller, Allen Sears**

*Ida Crown Jewish Academy*

*Science Department, 8233 North Central Park Avenue, Skokie, IL 60076, USA*

*E-mail: asears@icja.org*

**Paul Graham, Ellie Winkler**

*New Trier High School*

*Science Department, 7 North Happ Road, Northfield, IL 60093, USA*

*E-mail: nunterman@gmail.com*

**Alex Bernat, Shoshana Frank, Joshua Simon, Shira Eliaser**

*Rochelle Zell Jewish High School*

*Science Department, 1095 Lake Cook Road, Deerfield, IL 60015, USA*

*E-mail: seliaser@rzjhs.org*

---

<sup>1</sup>Speaker

## Abstract

Students and teachers from high schools near Chicago measured the rate of cosmic ray muons 100 meters underground in the Fermilab MINOS neutrino tunnel. The effect of the overburden was studied as a function of distance away from the access tunnel. A surface detector provided normalization over time and a third detector monitored possible background from the neutrino beam.

### 1. Aim

Students measured the change in cosmic ray flux in the neutrino tunnel as they move downstream from the access shaft. The change in burden is anticipated to affect the flux.

A measure of the change in flux during descent to the tunnel in the elevator was made.

Angle of acceptance will be factored into the results.

To guide high school students and their teachers in an authentic cosmic ray experiment.

### 2. Setup and Reasoning



**Figure 1** Moving tunnel arrays from assembly area to tunnel.



**Figure 2** Surface control detector.

Three Detectors Assembled:


- One to monitor cosmic rays above ground as a control.
- An identical detector (TC) to measure the rate underground.
- A third detector (TB) to measure the muon background in the neutrino beam.

The cosmic ray detectors simultaneously record data in large (21 degrees) and small (6 degrees) angular acceptance modes as well as data from the full sky.

### 3. Materials and Methods

- Tunnel Cosmic Ray (TC) stack is vertical.
- Tunnel Beam (TB) stack looks for horizontal (beamline origin) muons as a veto for TC.

**Tunnel Beam = TB**  
**Tunnel Cosmic Ray = TC**  
**TB inputs: 1,2,3,4**  
**muon trig beam = 124**  
**muon trig shaft = 13; 23**  
**TC inputs: 1,2,3,4**  
**muon trig large acceptance = 12**  
**cosmic trig medium acceptance = 123**  
**cosmic trig large acceptance = 124 or 1234**



POS(ICRC2019)170


### 4. Expectations

- As the detector below ground is moved along the tunnel, further away from the vertical access shaft, the ratio of muon flux below ground compared to the rate on the surface will decrease. This is due to the increase in the proportion of burden within the cone of acceptance of the underground detector.
- The muon flux will constantly decrease as the detector descends in the elevator because of the increased burden in the angles of acceptance.

## 5. Preliminary Results

- A possible decrease of flux was found as the distance from the access shaft. Further analysis is needed to subtract the effect of the neutrino beamline to verify preliminary results.
- Once the effect of the neutrino beamline is subtracted, angles of acceptance, TB1-TB3 and TB2-TB3 will be analyzed for any measureable effects when aimed toward and away from the shaft.

## 6. Elevator Results



A 200 fold decrease in flux was measured in the elevator while descending into the tunnel. At -95 meters, we notice a bump in the data. We have determined that the neutrino beam was on. Further analysis is being done to eliminate this noise.

## 7. Conclusion

High school students proposed, designed, built, and analyzed an authentic cosmic ray experience. The teachers also learned how to guide students in developing specific science skills and to shepherd a multi-school collaboration. The level of excitement and learning was a result of the outreach effort of the high energy physics community through QuarkNet.

## 8. Next Steps

After the analysis is completed, the students plan to submit an article for publication and present at the AAPT Winter Meeting.

## **9. Acknowledgments**

QuarkNet for supplying the teacher training and equipment.

FermiLab for providing the facility, guidance, safety training, and expert support during the experiment.

The science departments and administrations of Downers Grove South High School, Glenbrook High Schools, Ida Crown Jewish Academy, New Trier High School, and Rochelle Zell Jewish High School.