

Searches for lepton flavour and lepton number violation in K^+ decays at NA62

Joel Swallow^{†,*}

*The University of Birmingham,
School of Physics and Astronomy, Edgbaston B15 2TT, UK*

E-mail: joel.christopher.swallow@cern.ch

A search for the lepton number violating $K^+ \rightarrow \pi^- \mu^+ e^+$ and lepton flavour violating $K^+ \rightarrow \pi^+ \mu^- e^+$ decays is reported using NA62 data collected in 2017 and 2018. No significant excess is observed above the background expectation and therefore upper limits are established on the branching ratios at 90% confidence level: $\mathcal{B}(K^+ \rightarrow \pi^- \mu^+ e^+) < 4.2 \times 10^{-11}$ and $\mathcal{B}(K^+ \rightarrow \pi^+ \mu^- e^+) < 6.6 \times 10^{-11}$. These results improve over the previous limits by factors of 12 and 8 respectively.

40th International Conference on High Energy physics - ICHEP2020

July 28 - August 6, 2020

Prague, Czech Republic (virtual meeting)

*Speaker

[†]for the NA62 Collaboration:

R. Aliberti, F. Ambrosino, R. Ammendola, B. Angelucci, A. Antonelli, G. Anzivino, R. Arcidiacono, T. Bache, A. Baeva, D. Baigarashev, M. Barbanera, J. Bernhard, A. Biagioni, L. Bician, C. Biino, A. Bizzeti, T. Blazek, B. Bloch-Devaux, V. Bonaiuto, M. Boretto, M. Bragadireanu, D. Britton, F. Brizioli, M.B. Brunetti, D. Bryman, F. Buccini, T. Capussela, J. Carmignani, A. Ceccucci, P. Cenci, V. Cerny, C. Cerri, B. Checcucci, A. Conovaloff, P. Cooper, E. Cortina Gil, M. Corvino, F. Costantini, A. Cotta Ramusino, D. Coward, G. D'Agostini, J. Dainton, P. Dalpiaz, H. Danielsson, N. De Simone, D. Di Filippo, L. Di Lella, N. Doble, B. Dobrich, F. Duval, V. Duk, D. Emelyanov, J. Engelfried, T. Enik, N. Estrada-Tristan, V. Falaleev, R. Fantechi, V. Fascianelli, L. Federici, S. Fedotov, A. Filippi, M. Fiorini, J. Fry, J. Fu, A. Fucci, L. Fulton, E. Gamberini, L. Gatignon, G. Georgiev, S. Ghinescu, A. Gianoli, M. Giorgi, S. Giudici, F. Gonnella, E. Goudzovski, C. Graham, R. Guida, E. Gushchin, F. Hahn, H. Heath, J. Henshaw, E.B. Holzer, T. Husek, O. Hutanu, D. Hutchcroft, L. Iacobuzio, E. Iacopini, E. Imbergamo, B. Jenninger, J. Jerhot, R.W. Jones, K. Kampf, V. Kekelidze, S. Kholodenko, G. Khorauli, A. Khotyantsev, A. Kleimenova, A. Korotkova, M. Koval, V. Kozhuharov, Z. Kucerova, Y. Kudenko, J. Kunze, V. Kurochka, V. Kurshetsov, G. Lanfranchi, G. Lamanna, E. Lari, G. Latino, P. Laycock, C. Lazzeroni, M. Lenti, G. Lehmann Miotto, E. Leonardi, P. Lichard, L. Litov, R. Lollini, D. Lomidze, A. Lonardo, P. Lubrano, M. Lupi, N. Lurkin, D. Madigozhin, I. Mannelli, A. Mapelli, F. Marchetto, R. Marchevski, S. Martellotti, P. Massarotti, K. Massri, E. Maurice, M. Medvedeva, A. Mefodev, E. Menichetti, E. Migliore, E. Minucci, M. Mirra, M. Misheva, N. Molokanova, M. Moulson, S. Movchan, M. Napolitano, I. Neri, F. Newson, A. Norton, M. Noy, T. Numao, V. Obraztsov, A. Ostankov, S. Padolski, R. Page, V. Palladino, A. Parenti, C. Parkinson, E. Pedreschi, M. Pepe, M. Perrin-Terrin, L. Peruzzo, P. Petrov, Y. Petrov, F. Petrucci, R. Piandani, M. Piccini, J. Pinzino, I. Polenkevich, L. Pontisso, Yu. Potrebenikov, D. Protopopescu, M. Raggi, A. Romano, P. Rubin, G. Ruggiero, V. Ryjov, A. Salamon, C. Santoni, G. Saracino, F. Sargeni, S. Schuchmann, V. Semenov, A. Sergi, A. Shaikhiev, S. Shkarovskiy, D. Soldi, V. Sugonyaev, M. Sozzi, T. Spadaro, F. Spinella, A. Sturgess, J. Swallow, S. Trilov, P. Valente, B. Velghe, S. Venditti, P. Vicini, R. Volpe, M. Vormstein, H. Wahl, R. Wanke, B. Wrona, O. Yushchenko, M. Zamkovsky, A. Zinchenko.

1. Introduction

In the standard model (SM) the lepton number L and lepton flavour numbers L_ℓ , $\ell \in [e, \mu, \tau]$ are conserved due to apparent approximate global symmetries. However, this is an emergent property of the SM not required during its construction and evidence for violation of lepton flavour numbers (LFV) has been demonstrated by the observation of neutrino oscillations. No observation of LFV for charged leptons or any lepton number violating (LNV) process has been observed, and either would be a clear indication of physics beyond the SM (BSM).

A number of postulated BSM scenarios include mechanisms resulting in LFV and LNV. Seesaw models constructed to explain the mass of neutrinos, orders of magnitude lower than other fundamental fermions, introduce Majorana neutrino mass terms and exchange of Majorana neutrinos can facilitate LNV ($\Delta L = 2$) processes [1] [2]. In other BSM scenarios the existence of leptoquarks is postulated which can couple to both leptons and quarks, exchange of these particles can then allow LFV processes ($\Delta L_e = 1$, $\Delta L_\mu = 1$) [3] [4].

The NA62 experiment at CERN has collected the largest sample of K^+ decays to date and this data has been analysed to search for LNV and LFV processes. NA62 has set new upper limits on LNV $K^+ \rightarrow \pi^- \ell^+ \ell^+$ decays, of 2.2×10^{-10} and 4.2×10^{-11} (at 90% confidence level) in the cases of $\ell = e$ and $\ell = \mu$ respectively using a sub-set of 2017 data, representing 30% of the data collected to date (Run 1) [5]. In this proceedings contribution the search for $K^+ \rightarrow \pi^\mp \mu^\pm e^+$ decays using the full Run 1 data-set is reported.

2. The NA62 beamline and detector

The NA62 experiment, beamline and detector are described in detail in [6] and a schematic diagram is shown in figure 1. A high intensity 400 GeV/ c proton beam is delivered to the NA62 experiment by the CERN SPS in spills of 3 s effective duration containing $\sim 1.8 \times 10^{13}$ protons. The beam impinges onto a beryllium target producing a secondary hadron beam composed of π^+ (70%) protons (23%) and K^+ (6%), with mean momentum of 75 GeV/ c and a 1% rms spread. The KTAG, a differential Cherenkov counter, positively tags K^+ with a time resolution better than 70 ps. Kaon decays in a 75 m fiducial volume (FV), contained in a large vacuum tank, are studied. A magnetic spectrometer (STRAW), formed of four straw tracking chambers and a dipole magnet (M), is used to reconstruct the momenta of charged particles produced in K^+ decays in the FV. The CHOD, constructed from two sets of scintillator hodoscope planes, provides time measurements for charged tracks with 200 ps precision. For three-charged-track final states a vertex can be reconstructed in the FV, and associated to a K^+ decay if the total final state momentum is consistent with the (measured) average beam momentum, and is coincident in time with a KTAG K^+ tag signal. This means studies of three-charged-track decays do not necessarily require measurements of the beam K^+ momentum, provided by the gigatracker (GTK) and crucial to the study of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay [7]. Information for particle identification (PID), to distinguish π^\pm , μ^\pm and e^\pm , is provided by a quasi-homogeneous liquid krypton calorimeter (LKr) and muon detector (MUV3). A photon veto system is formed from twelve large angle veto (LAV) stations, the LKr and small angle vetos (IRC and SAC). The ring imaging Cherenkov detector (RICH), CHOD, LKr and MUV3 provide signals used for triggering.




Figure 1: Schematic top view of the NA62 beamline and detector.

3. Analysis Strategy

A search for $K^+ \rightarrow \pi^\mp \mu^\pm e^+$ decays has been performed using 2017 and 2018 NA62 data, equivalent to the full Run 1 data-set for these analyses. Potential signal events may be identified if the three-charged-track $\pi^\mp \mu^\pm e^+$ candidate final state has an invariant mass, $M_{\pi\mu e}$, consistent with the K^+ mass. A blind analysis strategy is adopted and the $M_{\pi\mu e}$ range 478–510 MeV/ c^2 was masked until the analysis was finalised. The signal region, 490–498 MeV/ c , was chosen, centred on the K^+ mass and with width approximately equal to 6 times the invariant mass resolution. Two independent analyses were performed with results cross-validated. The most common three-track K^+ decay, $K^+ \rightarrow \pi^+\pi^+\pi^-$, is used for normalisation. The core of the selection for normalisation and signal events is identical, requiring a three-track final state with vertex in the FV and consistent with a K^+ decay (see section 2) and vetoing photons detected by the LAVs. Following the common selection criteria normalisation events are identified with a requirement of invariant mass under the 3π hypothesis consistent with the K^+ mass, and PID conditions are applied to identify signal candidates with $\pi^\mp \mu^\pm e^+$ final states. For the $K^+ \rightarrow \pi^- \mu^+ e^+$ search it is additionally required that the invariant mass of the identified $\pi^- e^+$ pair under the $e^- e^+$ hypothesis $M_{\pi e} > 140$ MeV/ c^2 , rejecting backgrounds from decay chains involving $\pi^0 \rightarrow e^+ e^- \gamma$ (for example $K^+ \rightarrow \pi^+ [e^+ e^- \gamma]_{\pi^0}$) and misidentification.

Data from three triggers were used for these searches: the minimum bias three-track ‘Multi-Track’ trigger, and specialised ‘Multi-Track μ ’ and ‘Multi-Track e ’ triggers. The Multi-Track μ trigger saves events with three charged tracks, 10 GeV of energy deposition in the LKr and at least one time-coincident signal in the MUV3. The Multi-Track e trigger saves three-charged-track events with 20 GeV energy deposition in the LKr. These triggers are run concurrently with the main ‘PNN’ trigger, used for the study of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay [7], but downscaled by factors of approximately 100, 8 and 8 respectively.

All three triggers are used to collect candidate signal $K^+ \rightarrow \pi^\mp \mu^\pm e^+$ events while normalisation $K^+ \rightarrow \pi^+\pi^+\pi^-$ decays are collected with only the Multi-Track trigger. The effective number of K^+ decays in the FV that can be used for these searches, measured using the number of selected normalisation $K^+ \rightarrow \pi^+\pi^+\pi^-$ decays, the branching ratio of this decay and the downscaling factors




Figure 2: Misidentification probabilities between π^\pm and e^\pm as a function of track charge times momentum.

and inefficiencies of the triggers, is $(1.32 \pm 0.01) \times 10^{12}$.

4. Background Studies

There are two primary processes which can lead to background mechanisms: misidentification (misID) and decays in flight (DIF). For example, in both $K^+ \rightarrow \pi^\mp \mu^\pm e^+$ searches the most significant background at low $M_{\pi\mu e}$ arises from $K^+ \rightarrow \pi^+ \pi^+ \pi^-$ decays followed by one $\pi^\pm \rightarrow \mu^\pm \nu_\mu$ DIF and one misID of a π^+ as an e^+ . Simulations reproduce the effect of the DIF processes observed in data, however for misID data-driven models must be used. The probability of misID of a π^+ as an e^+ is measured using a control sample of $K^+ \rightarrow \pi^+ \pi^+ \pi^-$ decays collected with the Multi-Track trigger. A model is constructed to describe the misID probability as a function of track momentum as shown in figure 2.

Similarly, $K^+ \rightarrow \pi^+ e^+ e^-$ decays with an e^- misidentified as a π^- (and DIF or misID of the π^+ to produce a μ^+) can lead to a background in the signal region for the $K^+ \rightarrow \pi^- \mu^+ e^+$ search. In this case a control sample of e^\pm from $K^+ \rightarrow \pi^+ [e^+ e^- \gamma]_{\pi^0}$ decays is used to measure the misID probability of e^\pm as a π^\pm , as shown in figure 2.


Background expectations for the $K^+ \rightarrow \pi^\mp \mu^\pm e^+$ searches are, in general, evaluated using simulations with significant data-driven corrections, such as the data-driven misID models discussed above. Results are shown in tables 1 and 2 for the $K^+ \rightarrow \pi^- \mu^+ e^+$ and $K^+ \rightarrow \pi^+ \mu^- e^+$ searches respectively. Before unblinding the expected number of background events at high $M_{\pi\mu e}$, above the blinded region, is found to be compatible with the observed number of events. To validate the background predictions for the signal region the side-bands within the blinded region but outside the signal region were used as control regions, again good agreement is obtained between the expected and observed number of events. The predicted numbers of background events in the signal regions of the $K^+ \rightarrow \pi^- \mu^+ e^+$ and $K^+ \rightarrow \pi^+ \mu^- e^+$ searches are 1.06 ± 0.20 and 0.92 ± 0.34 respectively.

Table 1: Number of expected background and observed events for $K^+ \rightarrow \pi^- \mu^+ e^+$ search.

Region $M_{\pi\mu e}$ range [MeV/ c^2]	High mass Region	Control Regions (Sidebands)		Signal Region
	> 510	478–490	498–510	490–498
Total background expected	5.50 ± 0.53	1.68 ± 0.20	1.66 ± 0.26	1.06 ± 0.20
Data	8	2	4	0

Table 2: Number of expected background and observed events for $K^+ \rightarrow \pi^+ \mu^- e^+$ search.

Region $M_{\pi\mu e}$ range [MeV/ c^2]	High mass Region	Control Regions (Sidebands)		Signal Region
	> 510	478–490	498–510	490–498
Total background expected	1.95 ± 0.48	3.41 ± 0.54	1.27 ± 0.40	0.92 ± 0.34
Data	4	2	0	2

**Figure 3:** Reconstructed mass spectra for selected events in searches for $K^+ \rightarrow \pi^- \mu^+ e^+$ (left) and $K^+ \rightarrow \pi^+ \mu^- e^+$ (right) for data and simulated samples.

5. Results

After unmasking the signal region 0 and 2 events are found for the $K^+ \rightarrow \pi^- \mu^+ e^+$ and $K^+ \rightarrow \pi^+ \mu^- e^+$ searches respectively, with full invariant mass spectra shown in figure 3. This is compatible with the background expectations and therefore upper limits are established on the branching ratios, using the CLs method [8], at:

$$\mathcal{B}(K^+ \rightarrow \pi^- \mu^+ e^+) < 4.2 \times 10^{-11} @ 90\% CL, \quad (1)$$

$$\mathcal{B}(K^+ \rightarrow \pi^+ \mu^- e^+) < 6.6 \times 10^{-11} @ 90\% CL. \quad (2)$$

These represent an improvement on the previous limits, set by the BNL E865 experiment [9], by factors of 12 and 8 respectively.

6. Conclusion and Outlook

Searches for LNV/LFV decays $K^+ \rightarrow \pi^\mp \mu^\pm e^+$ have been performed using 2017 and 2018 (Run 1) NA62 data. New upper limits on the branching ratios are established, improving over previous constraints by approximately an order of magnitude. In general searches for LNV/LFV K^+ decays at NA62 are not limited by background and with data-taking resuming in 2021 NA62 has strong prospects to improve sensitivities for these searches.

References

- [1] L.S. Littenberg and R. Shrock, *Implications of improved upper bounds on $|\Delta L| = 2$ processes*, *Phys. Lett. B* **491** (2000) 285 [[hep-ph/0005285](#)].
- [2] A. Atre, T. Han, S. Pascoli and B. Zhang, *The Search for Heavy Majorana Neutrinos*, *JHEP* **05** (2009) 030 [[0901.3589](#)].
- [3] R.N. Cahn and H. Harari, *Bounds on the masses of neutral generation-changing gauge bosons*, *Nuclear Physics B* **176** (1980) 135 .
- [4] R. Mandal and A. Pich, *Constraints on scalar leptoquarks from lepton and kaon physics*, *JHEP* **12** (2019) 089 [[1908.11155](#)].
- [5] NA62 collaboration, *Searches for lepton number violating K^+ decays*, *Phys. Lett. B* **797** (2019) 134794 [[1905.07770](#)].
- [6] NA62 collaboration, *The Beam and detector of the NA62 experiment at CERN*, *JINST* **12** (2017) P05025 [[1703.08501](#)].
- [7] NA62 collaboration, *An investigation of the very rare $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay*, [2007.08218](#).
- [8] A.L. Read, *Presentation of search results: the CLs technique*, *Journal of Physics G: Nuclear and Particle Physics* **28** (2002) 2693.
- [9] R. Appel et al., *Search for lepton flavor violation in K^+ decays*, *Phys. Rev. Lett.* **85** (2000) 2877 [[hep-ex/0006003](#)].