

Tetraquark Interpretation and Production Mechanism of the Belle $Y_b(10750)$ -Resonance

Ahmed Ali,^a Luciano Maiani,^{b,c} Alexander Parkhomenko^{d,*} and Wei Wang^{b,e}

^aDeutsches Elektronen-Synchrotron DESY,
Street number, D-22607 Hamburg, Germany

^bT. D. Lee Institute, Shanghai Jiao Tong University,
Shanghai 200240, China

^cDipartimento di Fisica and INFN, Sapienza Università di Roma,
Piazzale Aldo Moro 2, I-00185 Roma, Italy

^dDepartment of Theoretical Physics, P. G. Demidov Yaroslavl State University,
Sovietskaya 14, 150003 Yaroslavl, Russia

^eINPAC, SKLPPC, MOE KLPPC, School of Physics and Astronomy, Shanghai Jiao Tong University,
Shanghai 200240, China

E-mail: ahmed.ali@desy.de, luciano.maiani@cern.ch, parkh@uniyar.ac.ru,
wei.wang@sjtu.edu.cn

Recently, the Belle Collaboration has updated the analysis of the cross sections for the processes $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) in the e^+e^- center-of-mass energy range from 10.52 to 11.02 GeV. A new structure, called $Y_b(10750)$, with the mass $M(Y_b) = (10752.7 \pm 5.9_{-1.1}^{+0.7})$ MeV and the Breit-Wigner width $\Gamma(Y_b) = (35.5_{-11.3-3.3}^{+17.6+3.9})$ MeV was observed. We interpret $Y_b(10750)$ as a compact $J^{PC} = 1^{--}$ state with a dominant tetraquark component. The mass eigenstate $Y_b(10750)$ is treated as a linear combination of the diquark-antidiquark and $b\bar{b}$ components due to the mixing via gluonic exchanges shown recently to arise in the limit of large number of quark colors. The mixing angle between Y_b and $\Upsilon(5S)$ can be estimated from the electronic width, recently determined to be $\Gamma_{ee}(Y_b) = (13.7 \pm 1.8)$ eV. The mixing provides a plausible mechanism for $Y_b(10750)$ production in high energy collisions from its $b\bar{b}$ component and we work out the Drell-Yan and prompt production cross sections for $pp \rightarrow Y_b(10750) \rightarrow \Upsilon(nS)\pi^+\pi^-$ at the LHC. The resonant part of the dipion invariant mass spectrum in $Y_b(10750) \rightarrow \Upsilon(1S)\pi^+\pi^-$ and the corresponding angular distribution of π^+ -meson in the dipion rest frame are presented.

40th International Conference on High Energy physics - ICHEP2020
July 28 - August 6, 2020
Prague, Czech Republic (virtual meeting)

*Speaker

1. Introduction

In 2019, the Belle Collaboration has presented the updated analysis of the cross sections for $e^+e^- \rightarrow \Upsilon(nS) \pi^+\pi^-$ ($n = 1, 2, 3$) in the electron-positron center-of-mass energy range from 10.52 to 11.02 GeV [1]. In addition to the already known structures $\Upsilon(10860)$ and $\Upsilon(11020)$, they reported the observation of a lower-mass resonance called $Y_b(10750)$ with the global significance of 5.2σ . The measured masses and decay widths (in MeV), and ranges of $\Gamma_{ee} \times \mathcal{B}$ (in eV) of all three structures are borrowed from [1] and presented in Table 1.

The combined analysis of the BaBar and Belle data on the ratio $R_{b\bar{b}}$ has been recently undertaken [2] with an account of the coherent sum of $\Upsilon(10860)$, $\Upsilon(11020)$, and $Y_b(10750)$ and a continuum amplitude, proportional to $1/\sqrt{s}$, where \sqrt{s} is the center-of-mass energy of the e^+e^- pair. Masses, Breit-Wigner decay widths, leptonic partial decay widths, and relative phases are fit parameters of the $R_{b\bar{b}}$ lineshape. One gets a number of solutions for the partial electronic widths (mathematically 8 solutions are expected), which differ in other parameters [2]. Most of these solutions are likely unphysical except the solution, in which the electronic width of Y_b is given as¹:

$$\Gamma_{ee}(Y_b(10750)) = (13.7 \pm 1.8) \text{ eV}. \quad (1)$$

The resulting masses and decay widths of the three resonances are found to be in agreement with the ones obtained from the $R_{\Upsilon \pi^+\pi^-}$ scan [1]. However, $Y_b(10750)$ is open to a number of interpretations to be tested in the existing and future experiments.

The conventional interpretation of $Y_b(10750)$ is that $Y_b(10750)$ and $\Upsilon(10680)$ are a mixture of $\Upsilon(5S)$ and $\Upsilon(4D)$ [4]. In [3], we interpret $Y_b(10750)$ as a $J^{PC} = 1^{--}$ tetraquark candidate, whose dominant component Y_b^0 consists of a colored diquark-antidiquark pair $[bq]_{\bar{3}_c} [\bar{b}\bar{q}]_{3_c}$, bound in the $SU(3)$ antitriplet-triplet representation [5, 6]. However, it can have a small $b\bar{b}$ component due to the mixing via gluonic exchanges. On the other hand, $\Upsilon(10860)$ and $\Upsilon(11020)$, which are dominantly radial $b\bar{b}$ excitations, $\Upsilon(5S)$ and $\Upsilon(6S)$, respectively, also have a small diquark-antidiquark component Y_b^0 in their content. Due to the proximity of the mass eigenstates $Y_b(10750)$ and $\Upsilon(10860)$, we consider that the mixing is dominantly between Y_b^0 and $\Upsilon(5S)$. This also provides a plausible interpretation of some anomalous features measured in the $\Upsilon(10860)$ -decays.

Mixing between a bottomonium and hidden-beauty tetraquark, anticipated in [7], was shown in [8], to be induced at the level of non-planar diagrams in the large- N_c limit. Albeit suppressed at large N_c by the exponential factor $e^{-N_c/2}$, when extrapolated back to $N_c = 3$ one finds a result not dissimilar from $f \sim N_c^{-3/2}$. Thus, a production in the e^+e^- -annihilation of resonances such as $Y_b(10750)$, in addition to the bottomonium spectral lines and with a small Γ_{ee} , is a significant signature of tetraquarks.

2. Bottomonium-Tetraquark Mixing Formalism

Let us define the tetraquark states Y_b^I ($I = 0, 1$) in the isospin basis as $Y_b^0 \equiv (Y_{[bu]} + Y_{[bd]})/\sqrt{2}$ and $Y_b^1 \equiv (Y_{[bu]} - Y_{[bd]})/\sqrt{2}$. Their mass difference due to the isospin breaking can be ignored. Their production is possible via the isosinglet $b\bar{b}$ -component, so only the Y_b^0 -state is considered.

¹In the analysis [3] on which this talk is based we used the preliminary results of X.-K. Dang et al. [2] presented by Chang-Zheng Yuan in XVth Rencontres du Vietnam (September 22nd – 28th, 2019, Quy Nhon, Vietnam).

Table 1: Measured masses and decay widths (in MeV), and ranges of $\Gamma_{ee} \times \mathcal{B}$ (in eV) of the $\Upsilon(10860)$, $\Upsilon(11020)$, and the new structure $Y_b(10750)$. The first uncertainty is statistical and the second is systematic.

State	$\Upsilon(10860)$	$\Upsilon(11020)$	$Y_b(10750)$
Mass	$10885.3 \pm 1.5^{+2.2}_{-0.9}$	$11000.0^{+4.0+1.0}_{-4.5-1.3}$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
Width	$36.6^{+4.5+0.5}_{-3.9-1.1}$	$23.8^{+8.0+0.7}_{-6.8-1.8}$	$35.5^{+17.6+3.9}_{-11.3-3.3}$
$\Upsilon(1S)\pi^+\pi^-$	0.75 – 1.43	0.38 – 0.54	0.12 – 0.47
$\Upsilon(2S)\pi^+\pi^-$	1.35 – 3.80	0.13 – 1.16	0.53 – 1.22
$\Upsilon(3S)\pi^+\pi^-$	0.43 – 1.03	0.17 – 0.49	0.21 – 0.26

Experimentally observed mass differences are $M[\Upsilon(10860)] - M[Y_b(10750)] \simeq 133$ MeV and $M[\Upsilon(11020)] - M[Y_b(10750)] \simeq 247$ MeV, so the mixing between $\Upsilon(10860)$ and $Y_b(10750)$ states should be more pronounced:

$$\begin{pmatrix} Y_b(10750) \\ \Upsilon(10860) \end{pmatrix} = \begin{pmatrix} \cos \tilde{\theta} & \sin \tilde{\theta} \\ -\sin \tilde{\theta} & \cos \tilde{\theta} \end{pmatrix} \begin{pmatrix} Y_b^0 \\ \Upsilon(5S) \end{pmatrix}. \quad (2)$$

The mixing angle $\tilde{\theta}$ is estimated phenomenologically. In general, the mixing can be easily generalized to the case of all three states. This mixing relates $\Gamma_{ee}[Y_b(10750)]$ and $\Gamma_{ee}[\Upsilon(5S)]$:

$$\frac{\Gamma_{ee}[Y_b(10750)]}{\Gamma_{ee}[\Upsilon(10860)]} = \tan^2 \tilde{\theta} \left[\frac{M[\Upsilon(10860)]}{M[Y_b(10750)]} \right]^4 \simeq 1.04 \tan^2 \tilde{\theta}. \quad (3)$$

LHS of this equation can be determined numerically, using $\Gamma_{ee}[\Upsilon(10860)] = (310 \pm 70)$ eV [9] and $\Gamma_{ee}[Y_b(10750)] = (13.7 \pm 1.8)$ eV. The estimate of the mixing angle is $\tan^2 \tilde{\theta} = 0.044 \pm 0.006$ and $\tilde{\theta} \simeq 12^\circ$ [3]. This supports the prediction from the large- N_c approach that the mixing angle between a pure bottomonium and hidden-bottom tetraquark state is relatively large.

3. Production Cross Sections at the LHC

Hadroproduction cross sections for $\Upsilon(5S)$ and $\Upsilon(6S)$ in $p\bar{p}$ (Tevatron) and pp (LHC) collisions were calculated using the NRQCD framework [10], assuming a factorization ansatz to separate the short- and long-distance effects. Cross-sections for $Y_b(10750)$ are scaled from the ones for $\Upsilon(5S)$, since in both cases the production takes place via the $b\bar{b}$ -component of $Y_b(10750)$. The cross-section ratio is determined by the mixing angle [3]:

$$\begin{aligned} & \frac{\sigma(pp \rightarrow Y_b(10750) + X) \mathcal{B}_f(Y_b(10750))}{\sigma(pp \rightarrow \Upsilon(10860) + X) \mathcal{B}_f(\Upsilon(10860))} \\ & \simeq \frac{\Gamma_{ee}(Y_b(10750)) \mathcal{B}_f(Y_b(10750))}{\Gamma_{ee}(\Upsilon(10860)) \mathcal{B}_f(\Upsilon(10860))} \simeq 1.04 \tan^2 \tilde{\theta} \frac{\mathcal{B}_f(Y_b(10750))}{\mathcal{B}_f(\Upsilon(10860))}, \end{aligned} \quad (4)$$

where $\mathcal{B}_f(Y_b(10750)) = \mathcal{B}(Y_b(10750) \rightarrow \Upsilon(nS)\pi^+\pi^-)$ and $\mathcal{B}_f(\Upsilon(10860)) = \mathcal{B}(\Upsilon(10860) \rightarrow \Upsilon(nS)\pi^+\pi^-)$ with $n = 1, 2, 3$. RHS of this equation has been measured by Belle [1]. Absolute

Table 2: Total cross sections (in pb) for the processes $pp \rightarrow Y_b(10750) \rightarrow (Y(nS) \rightarrow \mu^+\mu^-)\pi^+\pi^-$ ($n = 1, 2, 3$) at the LHC ($\sqrt{s} = 14$ TeV), assuming the transverse momentum range $3 \text{ GeV} < p_T < 50 \text{ GeV}$.

	QCD (gg)			Drell-Yan DY
	$n = 1$	$n = 2$	$n = 3$	
LHC 14	[0.29, 3.85]	[0.70, 4.78]	[0.45, 3.10]	[0.002, 0.004]
LHCb 14	[0.08, 1.21]	[0.20, 1.51]	[0.13, 0.99]	[0.001, 0.002]

cross sections for $Y(10860)$ are estimated in NRQCD [10]:

$$\sigma(pp \rightarrow Y(10860)+X) = \sum_Q \sigma_Q = \sum_Q \int dx_1 dx_2 \sum_{i,j} f_i(x_1) f_j(x_2) \hat{\sigma}(ij \rightarrow \langle \bar{b}b \rangle_Q + X) \langle O[Q] \rangle, \quad (5)$$

where $f_i(x_1)$ is the parton distribution function (PDF) of a generic i -th parton inside a proton, the label $Q = {}^{2S+1}L_J^c$ denotes the $b\bar{b}$ -pair quantum number (color c , spin S , and orbital L and total J angular momenta), $\langle O[Q] \rangle$ are long-distance matrix elements (LDMEs), and $\hat{\sigma} = \sigma / \langle O[Q] \rangle$ is a normalized partonic cross section. Leading-order partonic processes for the S -wave configurations are $g + g \rightarrow Y[{}^3S_1^1] + g$, $g + g \rightarrow Y[{}^1S_0^8, {}^3S_1^8] + g$, $g + q \rightarrow Y[{}^1S_0^8, {}^3S_1^8] + q$, and $q + \bar{q} \rightarrow Y[{}^1S_0^8, {}^3S_1^8] + g$. Details of the calculations are presented in [3].

Numerical estimates for hadroproduction and Drell-Yan cross sections at the LHC are collected in Table 2. Total cross sections (in pb) at $\sqrt{s} = 14$ TeV are calculated for $pp \rightarrow Y_b(10750) + X \rightarrow Y(nS)(\rightarrow \mu^+\mu^-)\pi^+\pi^- + X$ ($n = 1, 2, 3$) at the LHC, assuming the transverse momentum range $3 \text{ GeV} < p_T < 50 \text{ GeV}$. Rapidity ranges are $|y| < 2.5$ for ATLAS and CMS (called LHC 14) and $2.0 < y < 4.5$ for the LHCb. Error estimates in the QCD production are from the variation of the central values of the Color-Octet LDMEs and the various decay branching ratios, as discussed in [10]. Contributions from $Y(1S, 2S, 3S)$ are added together in the Drell-Yan production mechanism [11]. To estimate the expected number of events, we use 1 pb for the cross section, which lies in the middle of the indicated ranges, yielding $O(10^4)$ signal events at the LHCb, and an order of magnitude larger for the other two experiments, ATLAS and CMS. In addition to the production due to the mixing mechanism, there may be direct production of tetraquarks, which would add incoherently. Thus, the numbers in Table 2 give lower bounds to the expected $Y_b(10750)$ production in proton collisions.

4. Dipion Invariant Mass and Angular Distributions

The total amplitude $\mathcal{M} = \sum_I \mathcal{M}_I$ of the $Y_b(10750) \rightarrow Y(nS) PP'$ decay (here, $P^{(\prime)}$ denotes a pseudoscalar meson), and is the sum over possible isospin states $\mathcal{M}_I = \mathcal{S}, \mathcal{D}, \mathcal{D}', \mathcal{D}''$ [12]. Here, \mathcal{S} is the S -wave amplitude for PP' system and $\mathcal{D}, \mathcal{D}'$ and \mathcal{D}'' are the D -wave amplitudes. For the $\pi^+\pi^-$ -pair in the $Y_b(10750) \rightarrow Y(1S)\pi^+\pi^-$ decay, the isospin is $I = 0$ and the scalar $f_0(500)$ - and $f_0(980)$ - and the tensor $f_2(1270)$ -resonances contribute. The isospin-0 amplitudes are the combinations of the resonance amplitudes, $\mathcal{M}_0^{\mathcal{S}}$ and $\mathcal{M}_0^{f_2}$, and the non-resonating continuum amplitudes, \mathcal{M}_0^{1C} and \mathcal{M}_0^{2C} [10, 12]. The differential cross section for the $Y_b(10750) \rightarrow Y(nS)\pi^+\pi^-$ decay can be found in [10, 12].




Figure 1: The normalized resonant $m_{\pi^+\pi^-}$ (left plot) and $\cos\theta$ (right plot) distributions for $e^+e^- \rightarrow Y_b(10750) \rightarrow \Upsilon(1S)\pi^+\pi^-$ are shown using the coupling constants obtained in [12] (green histogram). The contributions from $f_0(500)$ and $f_0(980)$ scalars (left red curve) and $f_2(1270)$ (right black curve) are indicated in the left plot.

The $\pi^+\pi^-$ invariant mass and angular distributions in the $Y_b(10750) \rightarrow \Upsilon(1S)\pi^+\pi^-$ are presented in Fig. 1. They are normalized by the cross section $\sigma_{\Upsilon(1S)\pi^+\pi^-}^{\text{Belle}} = (1.61 \pm 0.16)$ pb measured by the Belle Collaboration [13]. Only resonant contributions are plotted, using the relevant input parameters [12]. Spectral shapes will be modified after a realistic non-resonant contribution is included. A fit can only be undertaken as updated experimental measurements become available.

We also obtain estimates of the branching fractions of $Y_b(10750) \rightarrow \Upsilon(nS)\pi^+\pi^-$ decays [10]. The products $\Gamma_{ee} \times \mathcal{B}$ are measured by the Belle Collaboration [1], while $\Gamma_{ee}[Y_b(10750)] = (13.7 \pm 1.8)$ eV is known from the Belle and BaBar data analysis on $R_{b\bar{b}}$ [2]. Corresponding ranges of the branching fractions are as follows [10]: $\mathcal{B}_{\Upsilon(1S)\pi^+\pi^-} = (0.9 - 3.4)\%$, $\mathcal{B}_{\Upsilon(2S)\pi^+\pi^-} = (3.9 - 8.9)\%$, and $\mathcal{B}_{\Upsilon(3S)\pi^+\pi^-} = (1.5 - 1.9)\%$. Note that due to the dominant tetraquark nature of $Y_b(10750)$, and its quark content, $Y_b(10750) \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$ decays are not anticipated, in agreement with the data from the Belle Collaboration [14].

5. Summary

The tetraquark-based interpretation of the new structure $Y_b(10750)$ found in the e^+e^- annihilation by the Belle Collaboration is presented. The $Y_b(10750)$ - and $\Upsilon(10860)$ -states are assumed to be the tetraquark- $b\bar{b}$ -mixed states, anticipated in the large- N_c limit. The $b\bar{b}$ -component is used to predict the hadroproduction and Drell-Yan cross sections at the LHC. A crucial test of this interpretation is in the $m_{\pi^+\pi^-}$ and $\cos\theta$ distributions in the $Y_b(10750) \rightarrow \Upsilon(nS)\pi^+\pi^-$ decays, whose resonant contribution is worked out. They are not expected in other dynamical schemes such as $Y_b(10750)$ interpreted as a D -wave $b\bar{b}$ -state, with a very large $S - D$ mixing. The tetraquark- $Q\bar{Q}$ mixing scheme suggested has wider implications.

Acknowledgments

The work of W. W. is supported in part the National Natural Science Foundation of China under Grant Nos. 11575110, 11735010, 11911530088, and the Natural Science Foundation of Shanghai under Grant No. 15DZ2272100. A. P. and W. W. acknowledge financial support by the Russian

Foundation for Basic Research and National Natural Science Foundation of China according to the joint research project (Nos. 19-52-53041 and 1181101282). This research is partially supported by the “YSU Initiative Scientific Research Activity” (Project No. AAAA-A16-116070610023-3).

References

- [1] BELLE collaboration, *Observation of a new structure near 10.75 GeV in the energy dependence of the $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) cross sections*, *JHEP* **10** (2019) 220 [1905.05521].
- [2] X.-K. Dong, X.-H. Mo, P. Wang and C.-Z. Yuan, *Hadronic cross section of e^+e^- annihilation at bottomonium energy region*, *Chin. Phys. C* **44** (2020) 083001 [2002.09838].
- [3] A. Ali, L. Maiani, A. Y. Parkhomenko and W. Wang, *Interpretation of $Y_b(10753)$ as a tetraquark and its production mechanism*, *Phys. Lett. B* **802** (2020) 135217 [1910.07671].
- [4] A. Badalian, B. Bakker and I. Danilkin, *Dielectron widths of the S -, D -vector bottomonium states*, *Phys. Atom. Nucl.* **73** (2010) 138 [0903.3643].
- [5] R. L. Jaffe and F. Wilczek, *Diquarks and exotic spectroscopy*, *Phys. Rev. Lett.* **91** (2003) 232003 [hep-ph/0307341].
- [6] L. Maiani, F. Piccinini, A. Polosa and V. Riquer, *Diquark-antidiquarks with hidden or open charm and the nature of $X(3872)$* , *Phys. Rev. D* **71** (2005) 014028 [hep-ph/0412098].
- [7] M. Knecht and S. Peris, *Narrow Tetraquarks at Large N* , *Phys. Rev. D* **88** (2013) 036016 [1307.1273].
- [8] L. Maiani, A. D. Polosa and V. Riquer, *Tetraquarks in the $1/N$ Expansion: a New Appraisal*, *Phys. Rev. D* **98** (2018) 054023 [1803.06883].
- [9] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2020** (2020) 083C01.
- [10] A. Ali, C. Hambrock and W. Wang, *Hadroproduction of $\Upsilon(nS)$ above $B\bar{B}$ Thresholds and Implications for $Y_b(10890)$* , *Phys. Rev. D* **88** (2013) 054026 [1306.4470].
- [11] A. Ali and W. Wang, *Production of the Exotic 1^{--} Hadrons $\phi(2170)$, $X(4260)$ and $Y_b(10890)$ at the LHC and Tevatron via the Drell-Yan Mechanism*, *Phys. Rev. Lett.* **106** (2011) 192001 [1103.4587].
- [12] A. Ali, C. Hambrock and S. Mishima, *Tetraquark-based analysis and predictions of the cross sections and distributions for the processes $e^+e^- \rightarrow \Upsilon(1S)(\pi^+\pi^-, K^+K^-, \eta\pi^0)$ near $\Upsilon(5S)$* , *Phys. Rev. Lett.* **106** (2011) 092002 [1011.4856].
- [13] BELLE collaboration, *Observation of anomalous $\Upsilon(1S)\pi^+\pi^-$ and $\Upsilon(2S)\pi^+\pi^-$ production near the $\Upsilon(5S)$ resonance*, *Phys. Rev. Lett.* **100** (2008) 112001 [0710.2577].
- [14] A. Abdesselam et al., *Study of Two-Body $e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$ Production in the Energy Range from 10.77 to 11.02 GeV*, **1609.08749**.