

Carpet-2 observation of $E > 300$ TeV photons accompanying a 150-TeV neutrino from the Cygnus Cocoon

Viktor Romanenko^{a,*} on behalf of the Carpet-3 Collaboration

(a complete list of authors can be found at the end of the proceedings)

^a*Institute for Nuclear Research of the Russian Academy of Sciences,
prospekt 60-letiya Oktyabrya 7a, Moscow, Russia*

E-mail: vsrom94@gmail.com

We report on the observation of an excess of $E > 300$ TeV gamma-ray candidate events in temporal and spatial coincidence with the IceCube high-energy neutrino alert consistent with the origin in the Cygnus Cocoon. The Cygnus Cocoon is a prospective Galactic source of high-energy neutrinos and photons. The observations have been performed with Carpet-2, a surface air-shower detector equipped with a large-area muon detector at the Baksan Neutrino Observatory in the Northern Caucasus.

37th International Cosmic Ray Conference (ICRC 2021)

July 12th – 23rd, 2021

Online – Berlin, Germany

*Presenter

1. Introduction

The observation of astrophysical gamma rays has confirmed the existence of Galactic sources with energies above 100 TeV. In particular, there was offered a good candidate for a Galactic PeVatron [1] from the HAWC collaboration results [2, 3]. This is the site where particles are accelerating up to PeV energies and gamma radiations may be produced by interactions of energetic hadrons with ambient matter and radiation. Recent observations of the LHAASO collaboration discovered the existence of such Galactics PeVatrons [4]. They detected 530 photons in the energy range of 0.1 up to 1.4 PeV from 12 Galactic sources. But at the moment we cannot unambiguously say about the generation mechanism of these photons and their cross observations with high-energy neutrinos produced in the same sources should help to establish their origin.

The first mentioned of Cygnus Cocoon was made in the work [5] by the Fermi-LAT collaboration. It is a superbubble around a region of active formation of massive stars, about 180 light-years across, located in the constellation Cygnus and containing two young star clusters, Cygnus OB2 and NGC 6910. According to the work [6], it may be a site for cosmic-rays acceleration and as a result a source for neutrino and gamma radiations of very high energy. However, the Cygnus Cocoon was associated with Galactic PeVatron in work [4] which makes it a motivated very high energy neutrino source.

On November 20, 2020, IceCube reported [7] about a candidate track-like neutrino event with an estimated energy of 154 TeV. The event was selected by the Bronze alert stream [8] with an average astrophysical neutrino purity of 30%. Inside the 90% of the localization region were several Fermi-LAT 4FGL sources [9] with the closest source is 4FGL J2028.6+4110e (Cygnus Cocoon). The arrival direction of the neutrino and the position of possible sources are shown in the figure 1. These alerts are routinely followed up by numerous instruments, in particular by the Carpet-2 gamma-ray telescope at the Baksan Neutrino Observatory [10]. This gives a chance to detect sub-PeV gamma rays co-produced with neutrinos, which cannot reach us from extragalactic sources because of pair production on cosmic microwave background radiation [11]. Standard Carpet-2 alert analysis revealed two gamma-ray candidate events in one-month intervals centered at the alert time [12]. Here, we present results of a more detailed study of possible sub-PeV gamma-rays from the Cygnus Cocoon associated with the IceCube neutrino event.

2. The Carpet-2 EAS array and the Dataset

Carpet-2 is an extensive air shower array located at the Baksan Neutrino Observatory, the North Caucasus. This array includes the Carpet array which consists of 196 m² large square detector with 400 scintillator counters at the center. Also, it is separated into 25 units with 16 scintillators in each and used for the reconstruction of primary-particle energy from the shower size N_e . Five detectors, each of 9 m² areas with 18 scintillators. Four detectors located at 30 m distance and one is at 40 m distance from the array center. They are used for the arrival direction reconstruction from the time delay of the outer detectors. All those detectors use the standard liquid scintillator counters of 0.49 m² area and 30 cm thickness. An underground muon detector (MD) is located at the distance of 47 meters from the center of the Carpet array. The MD was equipped with a 175 (5 × 35) plastic scintillator with an area of 1 m² each (the scintillator thickness is 5 cm), attached to the ceiling of




Figure 1: Arrival direction of the photon candidates (blue stars) coinciding in time of arrival with an IceCube-201120A neutrino event. The neutrino event (red rhombus) and some gamma-ray sources from the 4FGL catalog (stars of different colors).

the MD tunnel. The rock absorber above detectors is equal to 500 g/cm^2 , which corresponds to the threshold energy of 1 GeV for vertical muons. The muon detector is used for selecting candidate gamma-ray showers which are muon-poor. A description of the facility and some results can be found in [13–15].

For the present study, we used Carpet–2 data recorded between April 7, 2018 and April 25, 2021, total 829 days of data acquisition. Standard quality cuts require that ≥ 500 GeV air-shower energy is deposited in Carpet; four outer stations participate in the determination of the arrival direction; the reconstructed shower axis is at least 0.7 m within the Carpet boundary; the reconstructed zenith angle is $\leq 40^\circ$. In total, 65703 events passed these cuts in this time period.

3. Data analysis and Results

The Monte Carlo simulation used by us is described in [16]. These simulations are used to relate the reconstructed shower size N_e to the primary gamma-ray energy E , to estimate the detection and reconstruction efficiency, and to develop criteria for separation between events induced by primary photons and by cosmic rays. The effective area of the installation as a function of energy is presented in [10].




Figure 2: Dependence of the probability (p-value) on the width of the window centered on the neutrino arrival time (full line: all events, dashed line: photon candidates).




Figure 3: Number of events in $d = 82$ -days bins centered at a given day of data acquisition for all events. Blue dots correspond to the time of arrival of gamma candidate events. The vertical dashed line indicates the neutrino arrival time.

In what follows, we concentrate on the circular region in the sky of 4.7° angular radius (86% CL angular resolution) centered at the 4FGL J2028.6+4110e, associated with the Cygnus Cocoon, and call this region the Cygnus-Cocoon Circle (CCC). There is no excess from CCC over the expected background in the whole present Carpet-2 dataset. Therefore an upper limit on its integral gamma-ray flux was obtained: $I_\gamma(E_\gamma > 300 \text{ TeV}) < 2.6 \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}$ (95% CL).

But we got a hint of a possible flare if we consider the events around the IceCube neutrino alert. In total, 346 events from the Cygnus Cocoon were registered, five of them are associated with photons candidates and three photons coincide in time of arrival with the IceCube alert, Figure 1. To estimate the significance of the possible flare, an evaluation of the probability was carried out as a function of the window width with centering at the Ice Cube alert. Figure 2 shows those probabilities for all events and photon candidate events. Figure 3 presents the number of events in a sliding window of the width $d = 82$ days of data acquisition centered at a certain date, as a function of this date for all events. One can clearly see the enhancement around the neutrino event, consistent between all events.

For the full set $p_{\text{pre}} = 2.4 \times 10^{-4}$ (3.67σ pre-trial) at the $d = 82$, and for photon candidate are $p_{\text{pre}} = 5.5 \times 10^{-3}$ (2.78σ pre-trial) at the $d = 70$. Also, an introduced correction for window-width trials based on the Monte-Carlo simulation reduces the statistical significance of the probable flare. After correction, we have $p = 1.5 \times 10^{-3}$ (3.17σ post-trial) for all events and $p = 1.1 \times 10^{-2}$ (2.55σ post-trial) for the photon candidates respectively.

4. Conclusion

An excess of events was observed by Carpet-2 from the direction of the Cygnus Cocoon region in temporal coincidence with the IceCube neutrino alert from the same direction. The statistical significance of the excess is 3.1σ post-trial. For the first time, rare sub-PeV neutrino and gamma rays from the direction of a prospective Galactic PeVatron were observed in directional and temporal coincidence. This observation supports previously proposed scenarios of the origin of a part of observed high-energy neutrinos in pi-meson decays in Galactic sources. Additional details and discussion of the results can be found in [17].

Acknowledgement

The reported study was funded by RFBR, project numbers 20-32-90213 and 19-29-11027.

References

- [1] Albert, A., et al. The Astrophysical Journal Letters 896.2 (2020): L29.
- [2] Abeysekara, A. U., et al. The Astrophysical Journal 881.2 (2019): L134.
- [3] Albert, A., et al. The Astrophysical Journal Letters 907.2 (2021): L30.
- [4] Cao, Zhen, et al. Nature (2021): L1.
- [5] Ackermann, Markus, et al. science 334.6059 (2011): L1103-1107.

- [6] Bykov, Andrei M., et al. *Space Science Reviews* 216.3 (2020): L1-37.
- [7] IceCube Collaboration. *GRB Coordinates Network* 28927 (2020): 1.
- [8] Blaufuss, Erik, et al. arXiv preprint arXiv:1908.04884 (2019).
- [9] Abdollahi, Soheila, et al. *The Astrophysical Journal Supplement Series* 247.1 (2020): 33.
- [10] Dzhappuev, D. D., et al. *JETP Letters* 112.12 (2020): L753-756.
- [11] Nikishov, A. I. *JETP*. 41 (1961).
- [12] Dzhappuev, D., et al. *The Astronomer's Telegram* 14255 (2020).
- [13] Dzhappuev, D. D., et al. *Bulletin of the Russian Academy of Sciences: Physics* 71.4 (2007): 525-527.
- [14] Szabelski, J., and Carpet-3 Collaboration. *Nuclear Physics B-Proceedings Supplements* 196 (2009): 371-374.
- [15] Dzhappuev, D. D., et al. *EPJ Web of Conferences*. Vol. 207. EDP Sciences, 2019.
- [16] Dzhappuev, D. D., et al. *JETP Letters* 109.4 (2019): 226-231.
- [17] Dzhappuev, D. D., et al. arXiv preprint arXiv:2105.07242 (2021), *The Astrophysical Journal Letters* (2021) in press.

Full Authors List: Carpet-3 Collaboration

D. D. Dzhappuev¹, Yu. Z. Afashokov¹, I. M. Dzaparova^{1,2}, T. A. Dzhatdov^{3,1}, E. A. Gorbacheva¹, I. S. Karpikov¹, M. M. Khadzhiev¹, N. F. Klimenko¹, A. U. Kudzhaev¹, A. N. Kurenya¹, A. S. Lidvansky¹, O. I. Mikhailova¹, V. B. Petkov^{1,2}, E. I. Podlesnyi^{4,3,1}, G. I. Rubtsov¹, S. V. Troitsky¹, I. B. Unatlov¹, I. A. Vaiman^{4,3}, A. F. Yanin¹, Ya. V. Zhezher¹ and K. V. Zhuravleva¹

¹Institute for Nuclear Research of the Russian Academy of Sciences, 60th October Anniversary Prospect 7a, Moscow 117312, Russia.

²Institute of Astronomy, Russian Academy of Sciences, 119017, Moscow, Russia.

³D.V. Skobeltsyn Institute of Nuclear Physics, M. V. Lomonosov Moscow State University, Moscow 119991, Russia.

⁴Physics Department, M. V. Lomonosov Moscow State University, Moscow 119991, Russia.