

Phase diagram of the 2 + 1-dimensional Gross-Neveu model with chiral imbalance

Marc Winstel,^{a,*} Laurin Pannullo^a and Marc Wagner^{a,b}

^a*Institut für Theoretische Physik, Goethe Universität Frankfurt am Main
Max-von-Laue-Str. 1, 60438 Frankfurt am Main, Germany*

^b*Helmholtz Research Academy Hesse for FAIR,
Campus Riedberg, Max-von-Laue-Straße 12, D-60438 Frankfurt am Main, Germany
E-mail: winstel@itp.uni-frankfurt.de, pannullo@itp.uni-frankfurt.de,
mwagner@itp.uni-frankfurt.de*

In this work, the phase diagram of the 2 + 1-dimensional Gross-Neveu model is investigated with baryon chemical potential as well as chiral chemical potential in the mean-field approximation. We study the theory using two lattice discretizations, which are both based on naive fermions. An inhomogeneous chiral phase is observed only for one of the two discretizations. Our results suggest that this phase disappears in the continuum limit.

*The 38th International Symposium on Lattice Field Theory, LATTICE2021 26th-30th July, 2021
Zoom/Gather@Massachusetts Institute of Technology*

*Speaker

1. Introduction

The Gross-Neveu (GN) model describes N_f fermion flavors with a quartic interaction in the scalar channel. It is frequently used as a toy model for spontaneous breaking of chiral symmetry [1]. In general, four-Fermi theories are of interest in many applications in high energy physics [2–7], condensed matter physics¹ [8–18] but also of theoretical aspects of quantum field theory (e.g., renormalizability in the $1/N$ expansion or in a perturbative approach) [19–21]. In the mean-field approximation, the 2 + 1-dimensional GN model exhibits a symmetric phase, a homogeneous symmetry-broken phase and an inhomogeneous phase, where the chiral condensate is an oscillating function of the spatial coordinate [22–24].

Recently, two independent mean-field studies of the μ - T phase diagram (μ denotes the baryon chemical potential) of the 2 + 1-dimensional GN model were published [25–27]. An important result of these works is that the existence of an inhomogeneous phase depends on the details of the regularization. When the regulator (lattice spacing or Pauli-Villars cutoff) is removed, the inhomogeneous phase disappears.

In this work, we continue our previous lattice field theory investigations [25, 27] by studying the phase diagram of the 2 + 1-dimensional GN model with an additional chiral chemical potential μ_{45} .

2. The 2 + 1-dimensional GN model

The action of the 2 + 1-dimensional GN model with baryon chemical potential μ as well as chiral chemical potential μ_{45} is given by

$$S[\bar{\psi}, \psi] = \int d^3x \left(\sum_{n=1}^{N_f} \bar{\psi}_n (\gamma_\nu \partial_\nu + \gamma_0 \mu + \gamma_{45} \gamma_0 \mu_{45}) \psi_n - \frac{g^2}{2} \left(\sum_{n=1}^{N_f} \bar{\psi}_n \psi_n \right)^2 \right), \quad (1)$$

where ψ_n is a four-component massless fermion field, N_f is the number of flavors and g^2 is the coupling of the four-fermion interaction. A reducible representation of the Clifford algebra is chosen, where γ_0, γ_1 and γ_2 are block-diagonal. The upper two components of ψ_n can be interpreted as being “left-handed”, the lower two components as “right-handed”. There are two linearly independent matrices γ_4 and γ_5 , which anti-commute with γ_0, γ_1 and γ_2 and, thus, both fulfill the necessary properties of a suitable γ_5 matrix [7, 31]. Continuous symmetry transformations correspond to the generators \mathbb{I}_4 and γ_{45} ², while discrete chiral symmetry transformations are given by either γ_4 or γ_5 [6, 30]. Since $\gamma_{45} = \text{diag}(+\mathbb{I}_2, -\mathbb{I}_2)$, the chemical potential μ_{45} generates a chiral imbalance. More details can be found in Ref. [27] and will also be discussed in an upcoming publication.

By introducing an auxiliary boson field σ and integrating over the fermion fields, one obtains the effective action

$$S_{\text{eff}}[\sigma] = N_f \left(\frac{1}{2\lambda} \int d^3x \sigma^2 - \ln \left(\det(Q) \right) \right), \quad Z = \int D\sigma e^{-S_{\text{eff}}[\sigma]}, \quad (2)$$

¹Especially 2 + 1-dimensional models are often used to describe superconducting electrons in high-temperature superconductors, which are confined to planes.

²For N_f fermion fields one needs to consider tensor products with the generators of flavor rotations to get the full set of symmetry transformations.

where

$$Q = \gamma_v \partial_v + \gamma_0 \mu + \gamma_{45} \gamma_0 \mu_{45} + \sigma \quad (3)$$

is the Dirac operator and $\lambda = N_f g^2$. The effective action is real-valued, if the boson field σ is restricted to be constant in temporal direction (see appendix A of Ref. [27] for a proof). As $S_{\text{eff}} \propto N_f$, the limit $N_f \rightarrow \infty$ reduces the relevant configurations in the path integral to the global minima of S_{eff} , i.e., the mean-field approximation becomes exact in that limit. A Ward-Takahashi identity causes σ to be proportional to the chiral condensate $\langle \bar{\psi}_n \psi_n \rangle$. Moreover, the effective action Eq. (2) is invariant under $\sigma \rightarrow -\sigma$. Thus, σ is an order parameter for the discrete chiral symmetry.

The Dirac operator can be expressed in terms of irreducible representations of the Clifford algebra [28],

$$Q[\mu, \mu_{45}, \sigma] = \text{diag} \left(Q^{(2)}[\mu + \mu_{45}, \sigma], \tilde{Q}^{(2)}[\mu - \mu_{45}, \sigma] \right), \quad (4)$$

where $Q^{(2)}[\mu, \sigma]$ and $\tilde{Q}^{(2)}[\mu, \sigma]$ are defined as

$$Q^{(2)}[\mu, \sigma] = +\tau_2(\partial_0 + \mu) + \tau_3 \partial_1 + \tau_1 \partial_2 + \sigma, \quad (5)$$

$$\tilde{Q}^{(2)}[\mu, \sigma] = -\tau_2(\partial_0 + \mu) - \tau_3 \partial_1 - \tau_1 \partial_2 + \sigma. \quad (6)$$

The Dirac operators $Q^{(2)}$ and $\tilde{Q}^{(2)}$ contain two inequivalent 2×2 representations, $(\gamma_0, \gamma_1, \gamma_2) = (\tau_2, \tau_3, \tau_1)$ and $(\gamma_0, \gamma_1, \gamma_2) = -(\tau_2, \tau_3, \tau_1)$, with τ_j denoting the Pauli matrices. They act on the “left-handed” and “right-handed” components introduced above. For $\mu_{45} = 0$ their chemical potentials are the same, but for $\mu_{45} \neq 0$ they differ, $\mu + \mu_{45}$ and $\mu - \mu_{45}$, indicating chiral imbalance. The determinants of $Q^{(2)}$ and $\tilde{Q}^{(2)}$ are invariant under $\sigma \rightarrow -\sigma$ as well as under $\mu \rightarrow -\mu$. A consequence of the latter is that $\det Q$ as well as effective action (2) are invariant under the exchange of μ and μ_{45} . This property is used to check our numerical results presented in section 4.

One can show that

$$\det Q^{(4)}[\mu, \sigma] = \left(\det Q^{(2)}[\mu, \sigma] \right)^2 = \left(\det \tilde{Q}^{(2)}[\mu, \sigma] \right)^2 \quad (7)$$

with $Q^{(4)}[\mu, \sigma] = Q[\mu, 0, \sigma]$ (see Ref. [27] for the proof). In consequence, the same phase diagram would be obtained if a Dirac operator

$$Q'[\mu, \mu_{45}, \sigma] = \text{diag} \left(Q^{(4)}[\mu + \mu_{45}, \sigma], Q^{(4)}[\mu - \mu_{45}, \sigma] \right) \quad (8)$$

would be used in the action. Then, the spinor fields would obtain an additional degree of freedom, independent of chirality, that one can interpret as “isospin” and μ_{45} would serve as an isospin chemical potential. We will explore this concept further in an upcoming publication.

3. Lattice discretizations

We use lattice discretizations of Eq. (2) similar to that discussed in section 4 of Ref. [27]. The main difference is that we use a lattice also in temporal direction instead of a superposition of plane waves. The cubic space-time volume is $\beta \times V = \beta \times L \times L$ and the corresponding lattice has spacing a and $N_t \times N_s \times N_s$ sites, i.e., $\beta = a N_t$ and $L = a N_s$. The boundary conditions are antiperiodic in temporal direction and periodic in the two spatial directions.

For numerical computations it is convenient to work in momentum space. When using naive fermions, a weight function \tilde{W}_2 with specific properties has to be included in the interaction term of the GN model, to ensure the correct continuum limit (see Refs. [19, 27, 32], in particular appendix 2 of Ref. [19]). The Dirac operator is then

$$Q(p, q) = i\delta_{p,q} \sum_{\nu=0}^2 \gamma_\nu \sin(p_\nu - \delta_{\nu,0} i(\mu + \gamma_{45}\mu_{45})) + \delta_{p_0,q_0} \tilde{W}_2(\mathbf{p} - \mathbf{q}) \tilde{\sigma}(\mathbf{p} - \mathbf{q}), \quad (9)$$

where all quantities are expressed in units of a , i.e., dimensionless. In this work as well as in our previous work [27] we use and compare two weight functions corresponding to two different lattice discretizations,

$$\tilde{W}'_2(\mathbf{p}) = \prod_{\nu=1,2} \tilde{W}'_1(p_\nu) \quad , \quad \tilde{W}'_1(p_\nu) = \frac{\cos(p_\nu) + 1}{2} \quad (10)$$

and

$$\tilde{W}''_2(\mathbf{p}) = \prod_{\nu=1,2} \tilde{W}''_1(p_\nu) \quad , \quad \tilde{W}''_1(p_\nu) = \Theta(\pi/2 - |p_\nu|), \quad (11)$$

where $\Theta(p)$ is the Heaviside function.

4. Numerical results

The phase diagram at $\mu_{45} = 0$ is known from Refs. [26, 27]. There is a symmetric phase ($\sigma = 0$) at large μ and/or T and a homogeneous symmetry-broken phase with a spatially constant field $\sigma = \bar{\sigma} \neq 0$ at small μ and small T . For $W_2 = W'_2$ and finite lattice spacing there is also an inhomogeneous phase, where σ is a varying function of the spatial coordinates ($\sigma = \sigma(\mathbf{x})$), at small T between the symmetric and the homogeneous symmetry-broken phase. This phase, however, vanishes in the continuum limit. There is no inhomogeneous phase for $W_2 = W''_2$, neither at finite lattice spacing nor in the continuum limit.

In this section we discuss numerical results for the two lattice discretizations (10) and (11) with the aim to extend the phase diagram to $\mu_{45} \neq 0$. Technical aspects like scale setting and tuning of the coupling constant are explained in Ref. [27]. As usual, all dimensionful quantities are expressed in units of $\sigma_0 = \sigma|_{\mu=0, \mu_{45}=0, T \approx 0}$, the vacuum expectation value of σ .

4.1 The homogeneous phase diagram

When only allowing a homogeneous condensate, i.e., $\tilde{\sigma}(\mathbf{p}) = \bar{\sigma} \delta_{\mathbf{p},0}$, the lattice discretizations with W'_2 and W''_2 are identical (cf. Eq. (10) and Eq. (11) for $\mathbf{p} = 0$). The effective action at given μ , μ_{45} and T is a function of just one variable and can be minimized numerically in a straightforward way to determine the physically preferred value of $\bar{\sigma}$.

Fig. 1 shows the phase diagram in 3-dimensional μ - μ_{45} - T space at finite lattice spacing and space-time volume. We note that these results reflect the previously discussed $\mu \leftrightarrow \mu_{45}$ symmetry of the effective action. The observed phase boundary for $\mu_{45} = 0$ is close to the continuum result from Ref. [33], where small deviations are due to discretization errors and finite volume




Figure 1: Phase diagram of the chirally imbalanced 2 + 1-dimensional GN model with restriction to a homogeneous condensate in 3-dimensional μ - μ_{45} - T space for $a\sigma_0 = 0.2327$ and $L\sigma_0 = 27.92$.




Figure 2: “Phase diagram” of the chirally imbalanced 2 + 1-dimensional GN model obtained via stability analyses in 3-dimensional μ - μ_{45} - T space for $a\sigma_0 = 0.2327$ and $L\sigma_0 = 23.27$ and the discretization with $W_2 = W_2''$. The two plots show the same data from different angles.

corrections. There are also slight differences to our lattice results from Ref. [27], because of different discretizations in temporal direction.

For large temperature, $T/\sigma_0 \gtrsim 0.4$, the phase boundary exhibits an approximate rotational symmetry in the μ - μ_{45} plane, i.e., the phase boundary is crudely described by $\mu^2 + \mu_{45}^2 = \text{const.}$ The behavior is completely different at low temperatures, $T/\sigma_0 \lesssim 0.1$, where the phase boundary resembles a square in the μ - μ_{45} plane.

4.2 Instabilities with respect to spatially inhomogeneous perturbations

To study possibly existing inhomogeneous phases, one has to allow for arbitrary modulations, i.e., non-vanishing $\sigma(\mathbf{p})$, and to find the global minima of the effective action. In principle, such a minimization is possible within our lattice field theory approach, but it is a computationally extremely challenging task. For 1-dimensional modulations and $\mu_{45} = 0$ we were successfully using a local minimization algorithm with several randomly chosen initial field configurations [27]. In the near future we plan to use similar techniques to search for 2-dimensional modulations that are global minima, both at $\mu_{45} = 0$ and $\mu_{45} \neq 0$.

In this work, a simpler strategy is applied and the existence of inhomogeneous phases is studied via stability analyses. For given μ , μ_{45} and T we check the stability of the homogeneous condensate $\bar{\sigma}$, as determined in the previous subsection, with respect to spatially inhomogeneous perturbations $\delta\tilde{\sigma}(\mathbf{p})$ (for details we refer to Ref. [27]). If there is a perturbation $\delta\tilde{\sigma}(\mathbf{p})$, which lowers the effective action compared to the homogeneous condensate, an inhomogeneous phase is indicated. In this way regions in the phase diagram can be mapped, which correspond to or are part of an inhomogeneous phase.

Since there is no inhomogeneous phase in the continuum limit for chirally balanced matter [26, 27], we now explore, whether $\mu_{45} \neq 0$ might favor the existence of such a phase. We start with the discretization with $W_2 = W_2'$ and show in Fig. 2 the boundaries of the instability region in 3-dimensional μ - μ_{45} - T space at finite lattice spacing and space-time volume. The instability region resembles a tetrahedron and is adjacent to the homogeneously broken phase depicted in Fig. 1. Increasing $|\mu_{45}|$ seems to disfavor an inhomogeneous phase, because the instability region shrinks.

Computations at several values of the lattice spacing suggest that the instability region vanishes in the continuum limit also at $\mu_{45} \neq 0$. This is supported by computations with the discretization with $W_2 = W_2'$, where no instability region is observed, neither at $\mu_{45} = 0$ (as discussed in Ref. [27]) nor at $\mu_{45} \neq 0$. These results will be shown and discussed in more detail in an upcoming publication.

Acknowledgements

We acknowledge useful discussions with L. Kurth, A. Königstein, A. Sciarra, M. Thies and A. Wipf. We thank A. Königstein for useful comments on the manuscript and for bringing Ref. [31] to our attention.

We acknowledge support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the CRC-TR 211 “Strong interaction matter under extreme conditions” – project number 315477589-TRR 211. M. Wagner acknowledges support by the Heisenberg Programme of the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 399217702. M. Winstel acknowledges support by the GSI Forschungs- und Entwicklungsvereinbarungen (GSI F&E). Calculations on the GOETHE-HLR and on the on the FUCHS-CSC high-performance computers of the Frankfurt University were conducted for this research. We would like to thank HPC-Hessen, funded by the State Ministry of Higher Education, Research and the Arts, for programming advice.

References

- [1] D. J. Gross and A. Neveu, “Dynamical symmetry breaking in asymptotically free field theories”, *Phys. Rev. D* **10**, 3235 (1974).
- [2] V. V. Braguta and A. Y. Kotov, “Catalysis of Dynamical Chiral Symmetry Breaking by Chiral Chemical Potential,” *Phys. Rev. D* **93**, no.10, 105025 (2016) [arXiv:1601.04957 [hep-th]].
- [3] R. Narayanan, “Relevance of the three-dimensional Thirring coupling at finite temperature and density,” *Phys. Rev. D* **102**, no.1, 016014 (2020) [arXiv:2006.00608 [hep-th]].
- [4] S. Hands, A. Kocic and J. B. Kogut, “The Four Fermi model in three-dimensions at nonzero density and temperature,” *Nucl. Phys. B* **390**, 355-378 (1993) [arXiv:hep-lat/9206024 [hep-lat]].
- [5] S. J. Hands, J. B. Kogut and C. G. Strouthos, “The (2+1)-dimensional Gross-Neveu model with a U(1) chiral symmetry at nonzero temperature,” *Phys. Lett. B* **515**, 407-413 (2001) [arXiv:hep-lat/0107004 [hep-lat]].
- [6] H. Gies and L. Janssen, “UV fixed-point structure of the three-dimensional Thirring model”, *Phys. Rev. D* **82**, 085018 (2010) [arXiv:1006.3747 [hep-th]].
- [7] D. D. Scherer and H. Gies, “Renormalization group study of magnetic catalysis in the 3d Gross-Neveu model”, *Phys. Rev. B* **85** no. 19, 195417 (2012) [arXiv:1201.3746 [cond-mat.str-el]].
- [8] A. Chodos and H. Minakata, “The Gross-Neveu model as an effective theory for polyacetylene,” *Phys. Lett. A* **191**, 39 (1994).
- [9] T. Ohsaku, “Relativistic model for two-band superconductivity,” [arXiv:cond-mat/0306472 [cond-mat]].
- [10] R. MacKenzie, P. K. Panigrahi and S. Sakhi, “Superconductivity in a planar field theory through Kosterlitz-Thouless mechanism,” *Phys. Rev. B* **48** no. 6, 3892 (1993).
- [11] A. Kalinkin and V. M. Skorikov, “Phase transitions in four fermion models,” *Inorg. Mater.* **39**, 765-779 (2003).
- [12] D. Ebert, V. C. Zhukovsky, P. B. Kolmakov, K. G. Klimenko and R. N. Zhokhov, “Magnetization in (2+1)-dimensional Gross–neveu Model With Application to Carbon Nanotubes” *Particle Physics at the Year of Light*, World Scientific, 510-513 (2017).
- [13] D. Ebert and D. Blaschke, “Thermodynamics of a generalized graphene-motivated (2+1) D Gross–Neveu model beyond the mean field within the Beth–Uhlenbeck approach,” *PTEP* **2019**, no.12, 123I01 (2019) [arXiv:1811.07109 [cond-mat.mes-hall]].
- [14] K. G. Klimenko, R. N. Zhokhov and V. C. Zhukovsky, “Superconducting phase transitions induced by chemical potential in (2+1)-dimensional four-fermion quantum field theory,” *Phys. Rev. D* **86**, 105010 (2012) [arXiv:1210.7934 [hep-th]].

- [15] K. G. Klimenko, R. N. Zhokhov and V. C. Zhukovsky, “Superconductivity phenomenon induced by external in-plane magnetic field in (2+1)-dimensional Gross-Neveu type model,” *Mod. Phys. Lett. A* **28**, 1350096 (2013) [arXiv:1211.0148 [hep-th]].
- [16] V. C. Zhukovsky, K. G. Klimenko and T. G. Khunjua, “The effect of chiral density waves on the superconducting phase in the two-dimensional Gross-Neveu model,” *Moscow Univ. Phys. Bull.* **68**, 105-109 (2013).
- [17] S. Hands and A. S. Sehra, “Supercurrent flow in NJL(2+1) at high baryon density,” *Phys. Lett. B* **637**, 229-234 (2006) [arXiv:hep-lat/0510105 [hep-lat]].
- [18] S. Hands, B. Lucini and S. Morrison, “Numerical portrait of a relativistic thin film BCS superfluid,” *Phys. Rev. D* **65**, 036004 (2002) [arXiv:hep-lat/0109001 [hep-lat]].
- [19] J. Lenz, L. Pannullo, M. Wagner, B. Wellegehausen and A. Wipf, “Inhomogeneous phases in the Gross-Neveu model in 1+1 dimensions at finite number of flavors”, *Phys. Rev. D* **101**, no. 9, 094512 (2020) [arXiv:2004.00295 [hep-lat]].
- [20] S. Hands, A. Kocic and J. B. Kogut, “Compositeness, anomalous dimensions and renormalizability in four Fermi theories,” *Phys. Lett. B* **273**, 111-117 (1991).
- [21] J. Stoll, N. Zorbach, A. Koenigstein, M. J. Steil and S. Rechenberger, “Bosonic fluctuations in the (1 + 1)-dimensional Gross-Neveu(-Yukawa) model at varying μ and T and finite N ,” [arXiv:2108.10616 [hep-ph]].
- [22] M. Thies and K. Urlichs, “Revised phase diagram of the Gross-Neveu model”, *Phys. Rev. D* **67**, 125015 (2003) [arXiv:hep-th/0302092].
- [23] O. Schnetz, M. Thies and K. Urlichs, “Phase diagram of the Gross-Neveu model: exact results and condensed matter precursors”, *Annals Phys.* **314**, 425 (2004) [arXiv:hep-th/0402014].
- [24] M. Thies, “From relativistic quantum fields to condensed matter and back again: updating the Gross-Neveu phase diagram”, *J. Phys. A* **39**, 12707 (2006) [arXiv:hep-th/0601049].
- [25] M. Winstel, J. Stoll and M. Wagner, “Lattice investigation of an inhomogeneous phase of the 2 + 1-dimensional Gross-Neveu model in the limit of infinitely many flavors”, *J. Phys.: Conf. Ser.* **1667**, 012044 (2020) [arXiv:1909.00064 [hep-lat]].
- [26] R. Narayanan, “Phase diagram of the large N Gross-Neveu model in a finite periodic box”, *Phys. Rev. D* **101**, no. 9, 096001 (2020) [arXiv:2001.09200 [hep-th]].
- [27] M. Buballa, L. Kurth, M. Wagner and M. Winstel, “Regulator dependence of inhomogeneous phases in the 2 + 1-dimensional Gross-Neveu model,” *Phys. Rev. D* **103** no. 3, 034503 (2021) [arXiv:2012.09588 [hep-lat]].
- [28] K. Urlichs, “Baryons and baryonic matter in four-fermion interaction models”, doctoral thesis at the University of Erlangen-Nürnberg (2007).

- [29] B. Rosenstein, B. J. Warr and S. H. Park, “Thermodynamics of (2+1)-dimensional four Fermi models”, *Phys. Rev. D* **39**, 3088 (1989).
- [30] J. Braun, H. Gies and D. D. Scherer, “Asymptotic safety: a simple example,” *Phys. Rev. D* **83**, 085012 (2011) [arXiv:1011.1456 [hep-th]].
- [31] H. Gies, L. Janssen, S. Rechenberger and M. M. Scherer, “Phase transition and critical behavior of d=3 chiral fermion models with left/right asymmetry,” *Phys. Rev. D* **81**, 025009 (2010) [arXiv:0910.0764 [hep-th]].
- [32] Y. Cohen, S. Elitzur and E. Rabinovici, “A Monte Carlo study of the Gross-Neveu model,” *Nucl. Phys. B* **220**, 102-118 (1983).
- [33] K. G. Klimenko, “Phase structure of generalized Gross-Neveu models”, *Z. Phys. C* **37**, 457 (1988).