

Hypercubic symmetry restoration of Boriçi – Creutz fermions: costs and effects[†]

Rudina Osmanaj (Zeqirllari)^{a,*} and Dafina Xhako^b

^a *Physics Department, Faculty of Natural Sciences, University of Tirana
Bvd Zog I, Tirana, Albania*

^b *Department of Physics Engineering, Faculty of Mathematical Engineering and Physical Engineering,
Polytechnic University of Tirana
Street Sulejman Delvina, Tirana, Albania*

E-mail: rudina.zeqirllari@fshn.edu.al, dafinaxhako@yahoo.com

Minimally doubled fermions, have been proposed as lattice fermions which preserve chiral symmetry and being strictly local too. In our previous works, we have studied how the broken hypercubic symmetry of Boriçi - Creutz fermions affect the light hadrons spectrum, have proposed a method of how to restore it and also made a simple test on the pion mass using the corrected action. In this work we calculate the light hadrons spectrum, after have restored the hypercubic symmetry, in two different lattice directions, and see how the spectrum is affected. Also we investigate the computational cost of the restoration and how effective the use of BC fermions can be.

*The 38th International Symposium on Lattice Field Theory
LATTICE 2021, 26th-30th July, 2021
Zoom/Gather@Massachusetts Institute of Technology*

*Speaker

[†]This talk is dedicated to the memory of Prof. Artan Boriçi (19.4.1965 - 25.3.2021), who passed away after a long battle with Covid - 19

1. Introduction

Minimally doubled fermions have been proposed as a strictly local discretization of the QCD fermionic action, which preserves chiral symmetry at finite cut-off, while break the hyper-cubic symmetry. Being strictly local and chiral, they present the possibility of fast simulations. Anyway the broken hyper-cubic symmetry has to be restore, possibly with a minimal numerical cost. There are several types of MDF fermions, and each one of them breaks the hyper - cubic symmetry in a specific direction [8, 16]. One kind of minimally doubled fermions are Boriçi - Creutz fermions, proposed by Creutz and Boriçi, in 2008 [1, 5]. The Dirac operator for these fermions in the momentum space:

$$D(p) = \sum_{\mu} i \gamma_{\mu} \sin p_{\mu} + \sum_{\mu} i \gamma'_{\mu} \cos p_{\mu} - 2i \Gamma$$

where $\gamma'_{\mu} = \Gamma \gamma_{\mu} \Gamma$ $\Gamma = \sum_{\mu} \gamma_{\mu} / 2$ has two poles: $p_1 = (0,0,0,0)$ and $p_2 = (\pi/2, \pi/2, \pi/2, \pi/2)$.

These poles, that correspond to the two physical flavours, have a preferred direction in the euclidean space - time, defined by the line that connect them, which correspond to the major hyper - cube diagonal. Eventually, the hyper - cubic symmetry is broken [8,16]. In our previous publications we have shown the effects of this broken symmetry in the light hadrons spectrum [9, 10, 13, 14], have proposed a method of how to restore it partially for BC fermions using the chiral condensate [12, 13, 15] and also have presented some preliminary results of the calculations of the charged pion mass, in two different lattice directions, using the same conditions in which we have restored the hyper - cubic symmetry [14]. We have found that there a reduction of the difference of the masses calculated in different directions, which means that partially the broken hyper cubic symmetry is corrected. Regardless of these results, we have to see if this effect is represented even in the other light mesons and baryons masses. Also, the total computational cost (for defining the value of the counter-term in each case and for the calculation of the hadrons spectrum) has to be defined, in order to understand if this kind of fermions is suitable for the hadrons spectrum calculation.

2. Materials and methods

What we have done in this work, is to calculate the masses of the rho meson, nucleon and delta baryon in two different directions: in the hypercube diagonal direction and in one edge of the hypercube x_1 with the original Boriçi – Creutz action and then with the corrected Boriçi – Creutz action, where is added one of the counter-terms, dimension-3 counter-term. Let's remember that because of the broken hyper - cubic symmetry, the counter - terms are necessary for a renormalized theory. The allowed counter - terms for the BC action are dimension-4 counter-term $c_4(g_0)\psi\Gamma_{\mu}D_{\mu}\psi$ and dimension-3 counter-term $ic_3(g_0)\psi\Gamma\psi$ [16]. The coefficients of the dimension 3 and 4 counter-terms, c_3 and c_4 , respectively, have been evaluated in one-loop lattice perturbation theory [16].

The first step was to write the proper hadrons operators using the point splitting method, [6,11] as $\psi(p)$ contains two degenerate flavors. Details are described in the reference [11,13]. Sure, writing the operators using two flavours isn't very simple, and for complicated operators, this step will be challenging. Then we studied the dependence of the dimension-3 counter-term

from the lattice spacing. The methodology of how the counter-term value is defined for each lattice spacing, is presented in ref [12,15]. What was interesting was the fact that c_3 is almost independent from the lattice spacing. After this, we have repeated exactly the same procedure for the calculation of the light hadron masses, as described in the references [13, 14], but now even for the modified Boriçi – Creutz fermion action, where is added a counter-term $c_3 = 0.4$, that partially restore the hyper-cubic broken symmetry.

Boriçi – Creutz action in the second case (the corrected action) is:

$$D(p) = \sum_{\mu} [i\gamma_{\mu} \sin p_{\mu} + i(\Gamma - \gamma_{\mu}) \cos p_{\mu} + i(c_3 - 2)\Gamma]$$

Simulations are carried out on QCDCALAB (Tirana, Albania) in the quenched approximation for the Wilson gauge action, for the original BC action and the modified BC actions for five different bare mass quarks, five lattice spacings, using FermiQCD C++ package [18]. The codes written for the Boriçi – Creutz action and for the hadrons propagators are implemented in this package.

The gauge configurations are generated using the Wilson gauge action:

$$S_g[U] = -\beta \sum_{\mu \partial i} \frac{1}{3} \Re \text{Tr} U_{\mu, i} U_{\partial, i+\mu} U_{\mu, i+\partial}^{-1} U_{\partial, i}^{-1}$$

Let's remember that the quarks propagators or the Green functions of the Dirac operator are defined as:

$$\sum_{i, \alpha' \alpha} D_{\alpha \alpha', ii'}^{aa'} [U] S_{\alpha', \beta, ij}^{a'b} [U] = \delta_{ij} \delta_{\alpha \beta} \delta^{ab}$$

where $\alpha\beta$ are the Dirac indices, while ab - the color indices.

The Euclidean hadron propagators on the lattice with periodic boundary conditions can be written as:

$$S_{t, t_0} \cong \frac{1}{2} c_1 \cosh a m_1 \left(t - t_0 - \frac{L}{2} \right) + \frac{1}{2} c_2 \cosh a m_2 \left(t - t_0 - \frac{L}{2} \right)$$

where L is the lattice extension in time direction. To compute the hadron propagators we have used static point sources located at $t_0 = (0, 0, 0, 1)$ on 500 configurations. As a solver we have used an improved variant of the Stabilized Biconjugate Gradient method, BICGStab [17].

Point splitting method is used for defining each quark field and hadrons operators [11,6], whereas mass errors are computed using jackknife [7, 3].

3. Results

In the first figure, we have presented one of our previous results, the square mass of the neutral pion, using BC action with the added dimension 3 counter-term, as a function of the bare quark masses. As it expected, it goes to zero, in the chiral limit.




Figure 1. Pion mass as a function of the bare quark masses with the corrected BC action.

In the figures below are presented the results of the rho meson, nucleon and delta baryon masses, calculated in two different directions. In figure 2, are presented the results using the original BC action, while in figure 3 the case when the corrected BC action, with the dimension 3 counter-term is added.




Figure 2. Hadron masses of the light hadrons, in two different directions using the original BC action




Figure 3. Hadron masses of the light hadrons, in two different directions using the corrected BC action, where c_3 is added

In every case we have extrapolated the results in the continuum limit, and the taken results are presented in Table 1 (BC original action) and Table 2 (corrected BC actions)

Hadron	Masses in the diagonal direction (MeV/c ²)	Masses in the x_1 direction (MeV/c ²)
Rho meson	741	690
Nucleon	849	805
Delta baryon	1240	1187

Table 1. Hadron masses calculated with the original BC action

Hadron	Masses in the diagonal direction (MeV/c ²)	Masses in the x_1 direction (MeV/c ²)
Rho meson	741	730
Nucleon	860	852
Delta baryon	1240	1236

Table 2. Hadron masses calculated with the corrected BC action, where c_3 is added

What can be seen easily is the fact that for each hadron, the difference of masses calculated in different directions, begin to decrease when we have used BC action with the dimension 3 counter - term added, a sign that confirm that the hyper-cubic symmetry is partially restored, considering the errors too. Anyway the dimension 4 counter-term has to be added, but it needs

further studies, because this counter-term depends on the lattice spacing, and this increase the total computational cost and complexity.

4. Summary

Boriçi - Creutz fermions (minimally doubled fermions) are known to preserve chiral symmetry for a degenerate quark doublet and are strictly local. These fermions are ideal for studying QCD with u and d quarks, can be helpful for $N_f = 2$ lattice simulations, relatively simpler and possibly faster than Ginsparg-Wilson fermions. The chiral condensate can be used as an order parameter for BC fermions, and help us to find the proper counter-term that restore partially the broken hyper-cubic symmetry. Tuning of counter-term coefficient c_3 doesn't seem difficult and expensive, because apparently it is almost independent from the lattice spacing and coupling constant (further studies). Tuning c_4 will require more care and work so as to properly restore the desired symmetry, because it depends on the lattice and the coupling constant we use on simulations. (further studies) Using BC fermions with the added counter-term c_3 , the difference between the light hadrons masses calculated in different directions decrease, so we can think that partially have restore the broken hyper - cubic symmetry. Anyway....BC fermions, seem to be not the most suitable fermions for spectroscopy in full QCD

Acknowledgements

This work makes use of the endless contribution of Artan Boriçi in Lattice QCD. Artan, passed away on March 2021, after a long battle with COVID-19, and the authors of the paper dedicate this work to him.

References

- [1] A. Boriçi, "Minimally doubled fermion revival", arXiv preprint arXiv:0812.0092 (2008).
- [2] C. Lanczos, "Solution of systems of linear equations by minimized iterations." J. Res. Nat. Bur. Standards 49.1, 33-53, (1952).
- [3] J. Tukey, "Bias and confidence in not quite large samples", Ann. Math. Statist. 29, 614 (1958).
- [4] L. H. Karsten. "Lattice fermions in euclidean space-time". In: Physics Letters B 104.4, 315–319 (1981).
- [5] M. Creutz, "Four-dimensional graphene and chiral fermions", Journal of High Energy Physics 2008.04, 017, (2008).
- [6] M. Creutz, "Minimal doubling and point splitting", arXiv preprint arXiv:1009.3154, (2010).
- [7] M. Quenouille, "Problems in Plane Sampling," The Annals of Mathematical Statistics, 20, 3, 355–375, (1949).
- [8] P. Bedaque et al. "Broken symmetries from minimally doubled fermions", Physics Letters B 662.5, 449 - 455 (2008).
- [9] R. Zeqirllari, I. Buzi, and A. Boriçi, "Hypercubic Symmetry Breaking Effects On Pi And Rho Meson masses from simulations with Minimally Doubled Actions," AKTET, 4, 3, 392–396 (2011).

- [10] R. Zeqirllari, A. Boriçi, “Numerical Study of the Breaking Mass of Hyper - Cubic Symmetry for the Minimally Doubled Fermions Boriçi – Creutz”, *Open Journal Of Modern Physics* 1, 2, 96–104 (2014).
- [11] R. Zeqirllari, A. Boriçi, “Fermionet Rrjetore Boriçi - Creutz dhe ndarja pikësore e aromave”, *Buletini I Shkencave Natyrore, Universiteti i Tiranës* 19, 145-150 (2015), (in Albanian).
- [12] R. Osmanaj and D. Hyka. "Minimally doubled fermions and spontaneous chiral symmetry breaking", in *EPJ Web of Conferences* 175, EDP Sciences, (2018).
- [13] R. Osmanaj (Zeqirllari), D. Hyka (Xhako), A. Boriçi, "Rivendosja e simetrisë hiperkubike tek fermionet Boriçi - Creutz", *Konferenca e Shkencave të Natyrës dhe Teknologjisë*, (2017), (in Albanian).
- [14] R. Osmanaj – Zeqirllari, D. Hyka-Xhako, D., and A. Boriçi , “Preliminary results of the hypercubic symmetry restoration of Boriçi - Creutz fermions in the light hadrons spectrum”, (BPU10 – August 2018) in "10th Jubilee International Conference of the Balkan Physical Union", 2019, vol. 2075, no. 1. doi:10.1063/1.5091257
- [15] R. Osmanaj, D. Xhako, & N. Hyka “Application of Gauss - Lanczos Algorithm to Determine Low Modes Density of Dirac Operator”, *International Journal of Innovative Technology and Interdisciplinary Sciences*, 3(2), 443-450., (2020).
<https://doi.org/10.15157/IJITIS.2020.3.2.443>
- [16] S. Capitani, et al. "Renormalization of minimally doubled fermions", *Journal of High Energy Physics* 2010.9, 27, (2010).
- [17] Van der Vorst, H. A, "Bi-CGSTAB: A Fast and Smoothly Converging Variant of Bi-CG for the Solution of Nonsymmetric Linear Systems". *SIAM J. Sci. Stat. Comput.* 13 (2): 631–644. (1992), doi:10.1137/0913035. hdl:10338.dmlcz/104566.
- [18] www.fermiqcd.net