

Reactor antineutrino anomaly revisited

Z. Xin^{a,b,*}

^a*Institute of High Energy Physics, Chinese Academy of Sciences,
19B Yuquan Road, Beijing, China*

^b*School of Physical Sciences, University of Chinese Academy of Sciences,
19A Yuquan Road, Beijing, China*

E-mail: xinzhao@ihep.ac.cn

In our recent work [1], we studied the status of the reactor antineutrino anomaly in light of new reactor flux conversion and summation models. We present a new improved calculation of the IBD yields of the standard Huber-Mueller (HM) model and those of the new models. We show that the reactor rates and the fuel evolution data are consistent with the predictions of the Kurchatov Institute (KI) conversion model and with those of the Estienne-Fallot (EF) summation model, leading to a plausible robust demise of the reactor antineutrino anomaly. We also show that the results of several goodness of fit tests favor the KI and EF models over other models under consideration.

*** *The 22nd International Workshop on Neutrinos from Accelerators (NuFact2021)* ***

*** *6–11 Sep 2021* ***

*** *Cagliari, Italy* ***

*Speaker

1. Introduction

Reactor antineutrinos have been widely used to study the fundamental properties of neutrinos [2], which are mainly from beta decays of neutron-rich fission fragments generated by the heavy fissionable isotopes ^{235}U , ^{238}U , ^{239}Pu , and ^{241}Pu [3–5]. In 2011 improved calculations by Mueller *et al.* [6] and Huber [7] (HM model) predicted reactor antineutrino fluxes which are about 5% larger than the fluxes measured in several short-baseline reactor neutrino experiments. This discrepancy is known as the “reactor antineutrino anomaly” (RAA) [8].

There are two basic methods to predict reactor antineutrino fluxes: the summation method and the conversion method [4, 5]. The summation method is based on fission and decay information provided by the nuclear databases. The conversion method utilizes virtual branches to convert measured β spectra to corresponding antineutrino spectra. The converted ^{235}U , ^{239}Pu , and ^{241}Pu antineutrino spectra are based on the measurement at the Institut Laue-Langevin (ILL) in the 1980’s [9]. The converted ^{238}U antineutrino spectrum can be obtained based on the measured β spectrum at FRM II in Garching [10] in 2013. In addition to the HM model, we consider other three conversion models: HKSS [11], KI [12] and HKSS-KI models, and one summation model: EF [13] model.

Our updated calculation of IBD yields in all models is presented in Section 2, and then the methods of analysis are introduced as well. Our results of the fits of reactor rates and evolution data are also shown in Section 3. In Section 4, we further discuss which the best-fit model is. At last, we will summarize our conclusions.

2. Model predictions and method of analysis

The event rates are usually expressed as a physical quantity called “cross section per fission” $\sigma_{f,a}$ as known as “inverse beta decay (IBD) yield”:

$$\sigma_{f,a} = \sum_i f_i^a \sigma_i, \quad \text{with} \quad \sigma_i = \int_{E_\nu^{\text{thr}}}^{E_\nu^{\text{max}}} dE_\nu \Phi_i(E_\nu) \sigma_{\text{IBD}}(E_\nu) \quad (1)$$

where a is the experiment label, σ_i is the IBD yield for the fissionable isotope i , and f_i^a is the effective fission fraction of the isotope i . $\Phi_i(E_\nu)$ is the neutrino flux generated by the fissionable isotope i , and $\sigma_{\text{IBD}}(E_\nu)$ is the IBD cross section considered as the Strumia and Vissani cross section [14] including the radiative corrections with PDG2020 [2] here. The neutrino energy is integrated from $E_\nu^{\text{thr}} = 1.806$ MeV to $E_\nu^{\text{max}} = 10$ MeV using the EF model high-energy spectra with 100% uncertainties. The off-equilibrium corrections are also taken into account, which is based on Table VII of Ref. [6] assuming the 450 days approximation of the spectrum at equilibrium. Our updated results of IBD yields for these five models are shown in Table III in Ref. [1].

A χ^2 function based on Wilks’ theorem is usually applied to analyze the reactor antineutrino data. There are three main approaches when choosing different methods to deal with systematic theoretical uncertainties: **(A)** consider a covariance matrix with experimental and theoretical uncertainties added in quadrature [8]; **(B)** calculate the fit results considering only the experimental uncertainties and add by hand a global theoretical uncertainty [15]; **(C)** take into account the theoretical uncertainties with appropriate pull terms [16], which is chosen in this work:

$$\chi^2 = \sum_{a,b} \left(\sigma_{f,a}^{\text{exp}} - R_{\text{NP}}^a \sigma_{f,a}^{\text{th}} \right) (V^{\text{exp}})^{-1}_{ab} \left(\sigma_{f,b}^{\text{exp}} - R_{\text{NP}}^b \sigma_{f,b}^{\text{th}} \right) + \sum_{i,j} (r_i - 1) \left(\tilde{V}^{\text{mod}} \right)^{-1}_{ij} (r_j - 1), \quad (2)$$

$$\sigma_{f,a}^{\text{th}} = \sum_i r_i f_i^a \sigma_i^{\text{mod}}. \quad (3)$$

Here σ_i^{mod} denotes the IBD yield of the antineutrino flux generated by the fissionable isotope i and $\tilde{V}_{ij}^{\text{mod}} = V_{ij}^{\text{mod}} / (\sigma_i^{\text{mod}} \sigma_j^{\text{mod}})$, where V^{mod} is the covariance matrix. The coefficient R_{NP}^a is a possible suppression factor of the IBD yield in the experiment a .

3. Fit of reactor rates and reactor fuel evolution data

In this section, we firstly consider the reactor rates listed in Table IV of Ref. [1], and the results are presented in Table 1. For HM model, the fit of reactor rates $\bar{R}_{\text{HM}} = 0.936_{-0.023}^{+0.024}$, which indicates a RAA with 2.5σ .

Model	Rates		Evolution		Rates + Evolution	
	\bar{R}_{mod}	RAA	\bar{R}_{mod}	RAA	\bar{R}_{mod}	RAA
HM	$0.936_{-0.023}^{+0.024}$	2.5σ	$0.933_{-0.024}^{+0.025}$	2.6σ	$0.930_{-0.023}^{+0.024}$	2.8σ
EF	$0.960_{-0.031}^{+0.033}$	1.2σ	$0.975_{-0.030}^{+0.032}$	0.8σ	$0.975_{-0.030}^{+0.032}$	0.8σ
HKSS	$0.925_{-0.023}^{+0.025}$	2.9σ	$0.925_{-0.024}^{+0.026}$	2.8σ	$0.922_{-0.023}^{+0.024}$	3.0σ
KI	$0.975_{-0.021}^{+0.022}$	1.1σ	$0.973_{-0.022}^{+0.023}$	1.2σ	0.970 ± 0.021	1.4σ
HKSS-KI	$0.964_{-0.022}^{+0.023}$	1.5σ	$0.955_{-0.023}^{+0.024}$	1.9σ	$0.960_{-0.021}^{+0.022}$	1.8σ

Table 1: Average ratio \bar{R}_{mod} obtained from the least-squares analysis of the reactor rates in Table IV of Ref. [1] and of the Daya Bay and RENO evolution data.

There is practically no RAA for the EF model, since \bar{R}_{EF} differs from unity by only 1.2σ . Also the KI corrections lead to the practical disappearance of RAA, especially without the HKSS corrections. However, as for the 5 MeV bump, it is not fitted well by any of the models.

Then the reactor evolution data provided by Daya Bay and RENO collaborations are taken into account. To compare these flux evolution data with the different model predictions, we fit the evolution data with a linear function describing the IBD yield as a function of f_{239} , as done by the Daya Bay and RENO collaborations:

$$\sigma_{f,a}^{\text{lin}} = \bar{\sigma}_f + \frac{d\sigma_f}{df_{239}} (f_{239}^a - \bar{f}_{239}), \quad (4)$$

where $\bar{\sigma}_f$ is the average IBD yield and $d\sigma_f/df_{239}$ is the change in the IBD yield. The discrepancies of $d\sigma_f/df_{239}$ shown in Figure 1(b) are 3.5σ and 3.6σ for the HM model and HKSS model, respectively. However, other models give consistent values of $\bar{\sigma}_f$ and $d\sigma_f/df_{239}$.

Then, we further fit the evolution data and the combined data using Eq. 2 as done to reactor rates. In Table 1, the inclusion of the evolution data confirms the existence of a reactor antineutrino anomaly for the HM (2.8σ) and HKSS (3.0σ) models and the absence of the anomaly for the EF model (only 0.8σ). For KI and HKSS-KI models, the resulting 1.4σ (KI) and 1.8σ (HKSS-KI) are still too small to claim an anomaly.




Figure 1: Results of the linear fits of the Daya Bay evolution data.

4. Best-fit reactor flux model

In this section we apply goodness of fit tests to the reactor rates to select which the best-fit model is. Table VI in Ref. [1] shows that the standard χ^2 goodness of fit test rejects none of the five models if we consider the usual minimum p -value of 5% corresponding to a confidence level of 95%. However, the χ^2 test is only sensitive to the sizes not the signs of the deviations. Then, we applied the following tests: sign, Kolmogorov-Smirnov (KS), Cramer-von Mises (CVM), Anderson-Darling (AD), Z_K , Z_C , and Z_A tests, which are sensitive to the sign and size of the deviations of the transformed data with respect to the standard Gaussian distribution. More details can be seen in Ref. [1].

Since the EF and KI models are preferred by different tests and sets of data, it is fair to consider both as favorite. Therefore, we conclude that the KI model is the best among the conversion models and the only summation model that we considered, the EF model, is practically equally good.

5. Summary and conclusions

In Ref. [1] we revisited the reactor antineutrino anomaly based on the recent reactor antineutrino flux calculations. We first performed an improved calculation of the IBD yields of five reactor antineutrino flux models in Section 2. Then, in Section 3, based on the proper statistical method for the analysis, we calculated the suppression of the reactor antineutrino flux predicted by the 5 models by fitting the measured rates listed in Table IV of Ref. [1]. We found there is practically no anomaly for the EF, KI, and HKSS-KI models. The addition of the Daya Bay and RENO evolution data confirms these conclusions. In Section 4, we further explored the question of which is the best-fit model by applying several goodness of fit tests. We can consider EF as the best summation model and KI as the best conversion model, leaving the decision of a clear preference between the two models to future studies with more data.

References

- [1] C. Giunti, Y.F. Li, C.A. Ternes and Z. Xin, *Reactor antineutrino anomaly in light of recent flux model refinements*, [2110.06820](#).
- [2] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2020** (2020) 083C01.
- [3] C. Bemporad, G. Gratta and P. Vogel, *Reactor-based neutrino oscillation experiments*, *Rev. Mod. Phys.* **74** (2002) 297 [[hep-ph/0107277](#)].
- [4] P. Huber, *Reactor antineutrino fluxes - status and challenges*, *Nucl. Phys.* **B908** (2016) 268 [[arXiv:1602.01499](#)].
- [5] A.C. Hayes and P. Vogel, *Reactor Neutrino Spectra*, *Ann.Rev.Nucl.Part.Sci.* **66** (2016) 219 [[arXiv:1605.02047](#)].
- [6] T.A. Mueller et al., *Improved Predictions of Reactor Antineutrino Spectra*, *Phys. Rev.* **C83** (2011) 054615 [[arXiv:1101.2663](#)].
- [7] P. Huber, *On the determination of anti-neutrino spectra from nuclear reactors*, *Phys. Rev.* **C84** (2011) 024617 [[arXiv:1106.0687](#)].
- [8] G. Mention et al., *The Reactor Antineutrino Anomaly*, *Phys. Rev.* **D83** (2011) 073006 [[arXiv:1101.2755](#)].
- [9] F. Von Feilitzsch, A.A. Hahn and K. Schreckenbach, *Experimental beta spectra from Pu-239 and U-235 thermal neutron fission products and their correlated anti- neutrinos spectra*, *Phys. Lett.* **B118** (1982) 162.
- [10] N. Haag, A. Gutlein et al., *Experimental Determination of the Antineutrino Spectrum of the Fission Products of ^{238}U* , *Phys. Rev. Lett.* **112** (2014) 122501 [[arXiv:1312.5601](#)].
- [11] L. Hayen et al., *First-forbidden transitions in the reactor anomaly*, *Phys.Rev.* **C100** (2019) 054323 [[arXiv:1908.08302](#)].
- [12] V. Kopeikin, M. Skorokhvatov and O. Titov, *Reevaluating reactor antineutrino spectra with new measurements of the ratio between ^{235}U and ^{239}Pu β spectra*, [arXiv:2103.01684](#).
- [13] M. Estienne, M. Fallot et al., *Updated Summation Model: An Improved Agreement with the Daya Bay Antineutrino Fluxes*, *Phys. Rev. Lett.* **123** (2019) 022502 [[arXiv:1904.09358](#)].
- [14] A. Strumia and F. Vissani, *Precise quasielastic neutrino nucleon cross section*, *Phys. Lett.* **B564** (2003) 42 [[astro-ph/0302055](#)].
- [15] DAYA BAY collaboration, *Improved Measurement of the Reactor Antineutrino Flux and Spectrum at Daya Bay*, *Chin.Phys.* **C41** (2017) 013002 [[arXiv:1607.05378](#)].
- [16] C. Giunti, X.P. Ji, M. Laveder, Y.F. Li and B.R. Littlejohn, *Reactor Fuel Fraction Information on the Antineutrino Anomaly*, *JHEP* **1710** (2017) 143 [[arXiv:1708.01133](#)].